

Direct product of automorphism groups of digraphs

Mariusz Grech *

*Mathematical Institute, University of Wrocław, pl. Grunwaldzki 2/4,
50-384 Wrocław, Poland*

Wilfried Imrich

Montanuniversität Leoben, Franz Josef-Straße 18, 8700 Leoben, Austria

Anna Dorota Krystek †

*Faculty of Mathematics, Wrocław University of Science and Technology,
wyb. Wyspiańskiego 27, 50-370 Wrocław, Poland*

Łukasz Jan Wojakowski ‡

Nokia Networks, ul. Lotnicza 12, 54-155 Wrocław, Poland

Received 5 October 2017, accepted 24 January 2019, published online 22 June 2019

Abstract

We study the direct product of automorphism groups of digraphs, where automorphism groups are considered as permutation groups acting on the sets of vertices. By a direct product of permutation groups $(A, V) \times (B, W)$ we mean the group $(A \times B, V \times W)$ acting on the Cartesian product of the respective sets of vertices. We show that, except for the infinite family of permutation groups $S_n \times S_n$, $n \geq 2$, and four other permutation groups, namely $D_4 \times S_2$, $D_4 \times D_4$, $S_4 \times S_2 \times S_2$, and $C_3 \times C_3$, the direct product of automorphism groups of two digraphs is itself the automorphism group of a digraph. In the course of the proof, for each set of conditions on the groups A and B that we consider, we indicate or build a specific digraph product that, when applied to the digraphs representing A and B , yields a digraph whose automorphism group is the direct product of A and B .

Keywords: Digraph, automorphism group, permutation group, direct product.

Math. Subj. Class.: 05E18, 05C20, 20B25

*Supported by the Polish National Science Centre grant No. 2012/07/B/ST1/03318.

†Sponsored by the Polish National Science Centre grant No. 2012/05/B/ST1/00626 and by the travel grant PL 08/2016 of the OEAD/DWM.ZWB.183.1.2016.

‡Corresponding author. Sponsored by the Polish National Science Centre grant No. 2012/05/B/ST1/00626 and by the travel grant PL 08/2016 of the OEAD/DWM.ZWB.183.1.2016.

The original problem of König [20], to describe finite abstract groups that are isomorphic to automorphism groups of simple graphs, quickly found an answer due to Frucht [5], namely, each finite group is isomorphic to the automorphism group of some simple graph. A related question, asking which permutation groups on a given set are automorphism groups of graphs on that set of vertices, proved to be much more difficult.

The simplest example of a permutation group that has no graph representation in this sense is the trivial group on two elements. Both simple graphs on two vertices admit the full permutation group S_2 as automorphisms.

In the present paper, we deal with a generalization of the original problem. We study permutation group representability on directed simple graphs (digraphs). Note that the trivial group of the above example, while having no graph representation, obviously does have a digraph representation.

There are, however, groups that have neither graph nor digraph representations. The smallest example is the Klein four group $S_2 \times S_2$ (even symmetries of a square), and that is despite the fact that both factors do have graph representations. This observation led us to study the representability of direct products of representable groups.

Our main result is Theorem 2.1 that says that, given two permutation groups (A, V) and (B, W) that have digraph representations, their direct product $(A \times B, V \times W)$ also has a digraph representation, unless $A \times B$ is one of the four exceptional groups $D_4 \times S_2$, $D_4 \times D_4$, $S_4 \times S_2 \times S_2$, $C_3 \times C_3$, or a member of the infinite family of groups $S_n \times S_n$, $n \geq 2$. It is a digraph counterpart of Theorem 2.10 of [8] by Grech for undirected graphs.

Although it might seem that this generalization should be straightforward, it turns out that we are in need, in addition to the conclusions of the aforementioned paper, of a whole collection of new techniques. The reason is that, as we have already seen in the introduction, there are plenty of permutation groups that are not the automorphism groups of a graph but are the automorphism groups of a digraph with at least one directed edge.

Research on the problem of representability of a permutation group $A = (A, V)$ as the full automorphism group of a digraph (graph) $G = (V, E)$ started with studies of regular permutation groups (see [15, 16, 18, 23, 24, 25, 29, 30], for instance). In particular, it was established that abelian groups and generalized dihedral groups have no simple graph representation. Moreover, 13 other groups with this property were found. The solution of the problem for undirected graphs was completed by Godsil [7] in 1979. He proved that with the exception of the groups mentioned above, all other regular permutation groups are automorphism groups of graphs. For digraphs, L. Babai [1] in 1980 used the result of Godsil, and proved that, except for the groups S_2^2 , S_2^3 , S_2^4 , C_3^2 and the eight element quaternion group Q , each regular permutation group is the automorphism group of a digraph.

The fact that all digraphs and graphs can be interpreted as complete digraphs (graphs) in which the edges and non-edges are distinguished by assigning them one of two colors provides motivation for working with edge-colored digraphs (or graphs) rather than with plain digraphs (graphs). This subject was introduced by H. Wielandt in [32], where permutation groups that are automorphism groups of edge-colored digraphs were called 2-closed, and those that are automorphism groups of edge-colored graphs were referred to as 2*-closed. In [19] Kisielewicz introduced the notion of graphical complexity of permutation groups and suggested studying products of permutation groups in this context. We

denote by $\text{DGR}(k)$ ($\text{GR}(k)$) the class of automorphism groups of k -edge-colored digraphs (graphs), and by DGR (GR), the union of all classes $\text{DGR}(k)$ ($\text{GR}(k)$). A k -edge-colored digraph (graph) is a complete digraph (graph) with every edge colored in one of k colors. It is obvious that $\text{GR}(k) \subseteq \text{DGR}(k)$, for every k . Note that the class $\text{DGR}(2)$ ($\text{GR}(2)$) is the class of automorphism groups of digraphs (graphs).

The most general open question in this field is to find all permutation groups that belong to the class DGR . Another problem is to describe all the classes $\text{DGR}(k)$. Several results on $\text{DGR}(k)$ membership for basic classes of permutation groups are known, see for instance [1, 12, 34].

A closely connected topic is research on factorization of digraphs, see [3, 6, 22] and the bibliography given there. The same problem as before is considered, but from a slightly different point of view. Special attention is devoted to homogeneous factorization of complete digraphs [12, 21].

Also, various products of automorphism groups of digraphs were considered, see for instance [10, 11, 14, 28, 31]. In particular, in [10], the direct product of automorphism groups of edge-colored digraphs was studied. One of the results, worked out there, is that, for $k \geq 2$, the direct product $(A \times B, V \times W)$ of two permutation groups (A, V) and (B, W) from the class $\text{DGR}(k)$ belongs to the class $\text{DGR}(k + 1)$.

In [9] the study of the direct product was carried on and gave an improvement of the result from [10]. It was shown that for $k \geq 3$, the direct product of two groups from $\text{DGR}(k)$ is either in $\text{DGR}(k)$ or is equal to S_2^3 . The same holds for the case of automorphism groups of edge-colored graphs. The result of the present paper can be seen as an extension of the above result for the case $k = 2$.

1 Preliminaries

We assume that the reader has basic knowledge in the areas of graphs and permutation groups, so we omit an introduction to standard terminology. If necessary, additional details can be found in [2, 11, 33].

We recall the most important definitions. A *digraph* G is a pair (V, E) , where V is the set of vertices. The set of oriented edges, E , is a subset of $V \times V \setminus \{(v, v) : v \in V\}$ (the set of ordered pairs of different elements of V). By $\overline{G}$ we denote the complement of G . A complete digraph with n vertices is denoted by K_n .

An *undirected edge* is a pair $\{v, w\}$ such that both (v, w) and (w, v) belong to E . By $d_G^1(v)$ we mean the number of undirected edges of the form $\{v, w\}$, $w \in V$ in a digraph G (the number of 1-neighbors of the vertex v). We define the number of non-neighbors (or 0-neighbors) of a vertex v by $d_G^0(v) = d_G^1(v)$. If a digraph G is regular, then we denote these numbers $d^1(G)$ and $d^0(G)$, respectively. A *directed edge* is an edge $(v, w) \in E$ such that $(w, v) \notin E$. For every $v \in V$, by $d_G^f(v)$, we denote the number of its forward-neighbors, that is, of directed edges of the form (v, w) , $w \in V$ (with $(w, v) \notin E$).

In the case when a digraph G has no directed edges, we say that G is an *undirected graph* (a graph). For a digraph G we let $s(G)$ denote the undirected graph (shadow graph) that is obtained from G by replacing all directed edges by undirected ones. We will also use the notion of *weak neighbors* of a vertex v in a digraph G , that is, of vertices that are neighbors of v in $s(G)$. Similarly, a digraph is said to be *weakly connected* if $s(G)$ is connected.

We define two products of digraphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$. Their

Cartesian product $G_1 \square G_2$ is a digraph $G_1 \square G_2 = (V, E)$, where $V = V_1 \times V_2$, and $((v_1, v_2), (w_1, w_2)) \in E$ if either $(v_1, w_1) \in E_1$ and $v_2 = w_2$, or $v_1 = w_1$ and $(v_2, w_2) \in E_2$. We say that a digraph is *prime* if it is not the Cartesian product of two nontrivial digraphs. It is not hard to show that Cartesian multiplication of graphs is commutative, associative, and that K_1 is a unit.

The second product $G_1 * G_2 = (V, E)$, first studied by Watkins [28], is a digraph where $V = V_1 \times V_2$ and $((v_1, v_2), (w_1, w_2)) \in E$ if and only if either $(v_1, w_1) \in E_1$ and $v_2 = w_2$, or $v_1 \neq w_1$ and $(v_2, w_2) \in E_2$.

For a digraph G with vertex set $V \times W$, the subdigraphs of G induced by sets $V \times \{w\}$ will be called *rows*, and the subdigraphs induced by sets $\{v\} \times W$ will be called *columns*. An edge that belongs neither to a row nor to a column will be called a *slant edge*. When $G = G_1 \square G_2$, for given $v \in V(G)$ and $i \in \{1, 2\}$ we will use the notation *layer* for the row or column (image of G_i) containing v and denote it G_i^v .

A permutation σ of the set V is an automorphism of a digraph $G = (V, E)$ ($\sigma \in \text{Aut}(G)$) if, for $v, w \in V$, a pair $(v, w) \in E$ if and only if $(\sigma(v), \sigma(w)) \in E$. It is obvious that $\text{Aut}(G)$ is a group and that $\text{Aut}(G) = \text{Aut}(\overline{G})$.

All groups considered here are groups of permutations. They are considered up to permutation group isomorphism. S_n denotes the full group of permutations of an n -element set. By $C_n, n > 2$, we denote the cyclic group on n elements (i.e. the group generated by the cycle $(1, 2, \dots, n)$). And finally, by $D_n, n > 2$, we denote the dihedral group acting on an n -element set (i.e. the group generated by $(1, 2, \dots, n)$ and $(1, n)(2, n-1) \dots ([n/2], n - [n/2] + 1)$).

We define two kinds of products of permutation groups. Let A and B be permutation groups acting on the sets V and W , respectively. The *direct product* $A \times B$ is the permutation group consisting of the elements $\{(a, b) : a \in A, b \in B\}$ acting on the set $V \times W$ as follows: $(a, b)((v, w)) = (a(v), b(w))$, for $v \in V, w \in W$. The group $A \times A$ is denoted A^2 . A *wreath product* $A \wr B$ acting imprimitively on the set $V \times W$ is the permutation group consisting of the elements $\{(a, b_1, \dots, b_n) : a \in A, b_i \in B, n = |V|\}$ acting on the set $V \times W$ as follows: $(a, b_1, \dots, b_n)(i, w) = (a(i), b_i(w))$, where $i \in \{1, \dots, n\} = V, w \in W$. (A acts on the set of columns, B acts on each column independently.)

The class of groups which are the automorphism groups of digraphs with at least one directed edge will be denoted by EDGR.

Lemma 1.1. *Let G be a digraph and $v, w, x, y \in V(G)$, such that the only edges joining any two of them are $(v, w), (y, x) \in E(G)$ and $\{w, y\}, \{v, x\} \in E(G)$. Then, for every cartesian decomposition of the digraph $G = G_1 \square G_2$, there is an $i \in \{1, 2\}$ such that all the arcs between v, w, x, y belong to G_i^v .*

Proof. Without loss of generality assume that the layer G_1^v contains w . Vertex y can now be in the layer $G_1^v = G_1^w$ or in the layer G_2^w . Assume the latter. Then, x has to be at the intersection of G_1^y and G_2^v , as there are no slant arcs in G , but then the orientations of (v, w) and (y, x) are inconsistent with the definition of the cartesian product. Hence, vertex y must be in the layer $G_1^v = G_1^w$. Since the vertex x is a weak neighbor of both y and v which are in a single layer, it also must belong to that layer, because there are no slant arcs. $\square$

In contrast to the undirected case, where Imrich [14] found a short list of exceptional graphs for which both the graph and its complement are connected and not prime, for

digraphs with at least one directed edge there are no exceptions, as the following theorem shows:

Theorem 1.2. *For every digraph G with at least one directed edge either G or $\overline{G}$ is weakly connected and prime.*

Proof. Assume the digraph G with at least one directed edge is not prime, that is $G = G_1 \square G_2$. We have to show that $\overline{G}$ is weakly connected and prime.

Let $(v, w) = ((v_1, v_2), (w_1, w_2)) \in E(G)$ be one of the directed edges of G . Without loss of generality, assume that $(v_1, w_1) \in E(G_1)$ and $v_2 = w_2$. Since the cartesian decomposition is not trivial, there exists a vertex $v'_2 \in V(G_2)$, $v'_2 \neq v_2$. Then $((v_1, v'_2), (w_1, v'_2))$ is also a directed edge in $E(G)$. If between (v_1, v'_2) and (v_1, v_2) there is no edge or there is a directed edge, then it is easy to see that the subdigraph of $\overline{G}$ induced by the vertices (w_1, v_2) , (v_1, v_2) , (w_1, v'_2) , (v_1, v'_2) contains edges (directed or undirected) between every pair of vertices, and therefore belongs to a single layer of $\overline{G}$. If there is an undirected edge between (v_1, v'_2) and (v_1, v_2) then the same holds by Lemma 1.1. Now, all other vertices of $\overline{G}$ can be split into three categories according to their adjacency in $\overline{G}$ to the vertices (w_1, v_2) , (v_1, v_2) , (w_1, v'_2) , (v_1, v'_2) . First, those in G_1^v are neighbors of both (v_1, v'_2) and (w_1, v'_2) , and those in $G_1^{(v_1, v'_2)}$ are neighbors of both (v_1, v_2) and (w_1, v_2) . Second, those in G_2^v are neighbors of both w and (w_1, v'_2) and those in G_2^w are neighbors of both v and (v_1, v'_2) . Third, all other vertices are neighbors of all four vertices (w_1, v_2) , (v_1, v_2) , (w_1, v'_2) , (v_1, v'_2) .

Because a vertex can be a neighbor of two vertices in one and the same layer only if it also belongs to that layer, we conclude that all vertices in $\overline{G}$ belong to a single layer, so $\overline{G}$ is prime. It is easy to see that it also is weakly connected.

Assume now that G is prime and not weakly connected. Its complement $\overline{G}$ is connected. If $\overline{G}$ were not prime, then, by the previous paragraph, $G = \overline{\overline{G}}$ would have to be weakly connected, contrary to assumption. Thus $\overline{G}$ is weakly connected and prime. $\square$

In what follows we need a result analogous to the Sabidussi-Vizing [26, 27] theorem about the automorphism group of the Cartesian product of connected coprime graphs. To prove it, we use a result on unique prime factorization of digraphs with respect to the Cartesian product. This result can be traced back to Feigenbaum [4], but for an easy proof in a more general setting we refer to the recent paper by Imrich and Peterin [17]:

Theorem 1.3. *Every weakly connected digraph has a unique prime factor decomposition with respect to the Cartesian product.*

We can now state our two simplified versions of the Sabidussi-Vizing theorem for digraphs.

Theorem 1.4. *Let G, H be non-isomorphic weakly connected digraphs, where $|V(G)| \geq |V(H)|$ and G is prime. Then $\text{Aut}(G \square H) = \text{Aut}(G) \times \text{Aut}(H)$.*

Proof. It is clear that $\text{Aut}(G) \times \text{Aut}(H) \subset \text{Aut}(G \square H)$. We shall prove the opposite inclusion. To that end, it suffices to show that every $a \in \text{Aut}(G \square H)$ maps G -layers to G -layers and H -layers to H -layers in $G \square H$.

We know that $\text{Aut}(G \square H) \subset \text{Aut}(s(G \square H))$ and, in general, the factors of the shadow graph $s(G \square H) = s(G) \square s(H)$ need not be prime. Take $a \in \text{Aut}(G \square H)$. A G -layer in $G \square H$ has the form $G \square \{h\}$ for $h \in V(H)$. Consider $s(G \square \{h\})$, a

cartesian product of subgraphs of $s(G)$ and $s(H)$. Using the terms defined in Chapter 6 of [13], it is a convex subgraph of the shadow graph $s(G \square H)$, and so, by a corollary that leverages the convexity preserving property of automorphisms, obtained as a step in the proof of Theorem 6.8 therein (first paragraph on page 69), the image of $s(G \square \{h\})$ under the automorphism a is again a cartesian product of subgraphs of $s(G)$ and $s(H)$, that is, $a(s(G \square \{h\})) = s(G_1) \square s(H_1)$, where $G_1 \subset G$ and $H_1 \subset H$. But, since the vertex sets of the shadows are the same as those of the digraphs, we also have that $a(G \square \{h\}) = G_1 \square H_1$. Suppose $|V(G_1)| = 1$, that would imply that $H_1 = H$ with $|V(H)| = |V(G)|$ and that G is isomorphic to H , which is contrary to assumption. Now suppose that $1 < |V(G_1)| < |V(G)|$. This would imply that the digraph G has a nontrivial cartesian product decomposition, which is also contrary to assumption. We are, thus, left with the case $|V(G_1)| = |V(G)|$, which proves that a maps G -layers to G -layers.

Because we have no slant arcs and H is weakly connected this means that a maps H -layers into H -layers. $\square$

Theorem 1.5. *Let G be a weakly connected, prime digraph with at least one directed edge. Let H be an undirected and connected graph. Then $\text{Aut}(G \square H) = \text{Aut}(G) \times \text{Aut}(H)$.*

Proof. Similarly as above, we get that $a(s(G \square \{h\})) = s(G_1) \square s(H_1)$. We do not assume that the digraph G has at least as many vertices as H , so we need to exclude the case $|V(G_1)| = 1$ differently. Here this would imply that G is a subgraph of H , but this is not possible as G has a directed edge while H does not. The conclusion follows as above. $\square$

The following proposition is modelled on an observation made in the proof of Theorem 6 of Watkins [28]:

Proposition 1.6. *Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be digraphs where G_2 is weakly connected. Suppose that every automorphism a of the digraph $G = G_1 * G_2$ maps rows onto rows. Then $\text{Aut}(G) = \text{Aut}(G_1) \times \text{Aut}(G_2)$.*

Proof. Let w_1 and w_2 be weak neighbors in G_2 and let $v \in V_1$ be arbitrarily chosen. Write $a(v, w_i) = (a_1(v, w_i), a_2(v, w_i))$. Since rows are mapped onto rows, a_2 does not depend on v . Hence, $a_2 \in \text{Aut}(G_2)$.

By the definition of the $*$ -product, (v, w_2) is the only vertex in $G_1^{(v, w_2)}$ that is not weakly adjacent to (v, w_1) . Hence $a(v, w_2) = (a_1(v, w_2), a_2(w_2))$ is the only vertex in $G_1^{a(v, w_2)}$ that is not weakly adjacent to $(a_1(v, w_1), a_2(w_1))$, so $a_1(v, w_1)$ must be equal to $a_1(v, w_2)$. By the weak connectivity of G_2 this means that a_1 only depends on v . It is easily seen that it is an automorphism of G_1 . Thus, for any $(v, w) \in V(G)$ we conclude that $a(v, w) = (a_1(v), a_2(w))$, where a_1, a_2 are automorphisms of G_1 , resp. G_2 . $\square$

2 Main result

The following theorem settles the problem when the direct product of automorphism groups of digraphs is an automorphism group of a digraph.

Theorem 2.1. *Let $A, B \in \text{DGR}(2)$. Then $A \times B \in \text{DGR}(2)$, unless $A \times B$ is $D_4 \times S_2$, $D_4 \times D_4$, $S_4 \times S_2 \times S_2$, $C_3 \times C_3$, or one of the groups $S_n \times S_n$, $n \geq 2$.*

The proof is broken up into a series of lemmas. Let us note first that we are given permutation groups $A = (A, V_A)$, $B = (B, V_B)$ and graphs $G_A = (V_A, E_A)$, $G_B = (V_B, E_B)$, where $\text{Aut}(G_A) = A$ and $\text{Aut}(G_B) = B$. Since $\text{Aut}(G) = \text{Aut}(\overline{G})$ for any G we may assume without loss of generality that both G_A and G_B are weakly connected. Moreover, by Theorem 1.2 we may also assume that they are prime if they have at least one directed edge.

We begin by extending Theorem 2.10 of [8] by Grech for undirected graphs to directed graphs.

Lemma 2.2. *Let $A, B \in \text{GR}(2)$. Then $A \times B \in \text{DGR}(2)$ if and only if $A \times B \in \text{GR}(2)$.*

Proof. By Theorem 2.10 of [8], $A \times B \in \text{GR}(2)$, unless $A \times B$ is $D_4 \times S_2$, $D_4 \times D_4$, $S_4 \times S_2 \times S_2$ or $S_n \times S_n$, for $n \geq 2$. In the exceptional cases the pair (v_2, v_1) belongs to the orbit of the pair (v_1, v_2) in the natural action of the group $(A \times B, V)$ on pairs of elements of V . Thus, every digraph G such that $A \times B \subseteq \text{Aut}(G)$ has to be an undirected graph. Hence, in all the cases, $A \times B \in \text{DGR}(2)$ would imply $A \times B \in \text{GR}(2)$. Consequently, in the exceptional cases, $A \times B \notin \text{DGR}(2)$. $\square$

Notice that this takes care of all exceptional groups of Theorem 2.1 that are different from $C_3 \times C_3$. The proof also shows that in what follows it suffices to consider only the cases where either A or B admits a digraph representation with at least one directed edge. We can thus assume without loss of generality that $A \in \text{EDGR}$.

Lemma 2.3. *Assume that A, B are non-isomorphic groups, where $A \in \text{EDGR}$ and $B \in \text{DGR}(2)$. Then $\text{Aut}(G_A \square G_B) = \text{Aut}(G_A) \times \text{Aut}(G_B)$*

Proof. As noted above, G_A and G_B can be chosen to be weakly connected, the complement being taken if necessary, with G_A being prime. Then, if $B \in \text{EDGR}$ so that G_B can also be chosen to be prime, the proof follows from Theorem 1.4, and from Theorem 1.5 otherwise. $\square$

This means that we can assume that $B \cong A$. Moreover, if we are able to find two non-isomorphic weakly connected digraphs, at least one of which is prime, with the same automorphism group A , then Theorem 1.4 also gives us a positive answer.

It therefore remains to consider the case $A \times A$, where A is the automorphism group of a weakly connected prime digraph G_A with at least one directed edge. In other words, we can assume that $A \in \text{EDGR}$ and that G_A is prime.

Lemma 2.4. *Let $A \in \text{EDGR}$ with prime G_A . If A is intransitive, then $A \times A \in \text{DGR}(2)$.*

Proof. We consider two copies $G_r = (V_r, E_r)$ and $G_c = (V_c, E_c)$ of G_A and will define a digraph $G = (V_r \times V_c, E)$ such that $\text{Aut}(G) = A \times A$. We call G_r the *row copy* and G_c the *column copy* of G_A .

Since A is intransitive, $G_A \neq K_{|V_A|}$. Let $W \subset V_c$ be one of the orbits of A in its action on G_c . The edge set E of the digraph $G = (V_r \times V_c, E)$ is then defined as the set of all pairs $((v_r, v_c), (w_r, w_c))$ satisfying one of the following conditions:

- (a) $(v_c, w_c) \in E_c$ and $v_r = w_r$;
- (b) $v_c = w_c$ and

- either $v_c \in W$

- or $v_c \notin W$, and $(v_r, w_r) \in E_r$.

Notice that there are no slant edges and that the subgraphs induced by the columns $\{v_r\} \times V_c$ are isomorphic to G_A , whereas the subgraphs induced by the rows $V_r \times \{v_c\}$ are isomorphic to $K_{|V_A|}$ if $v_c \in W$, otherwise they are isomorphic to G_A .

In other words, $V_r \times W$ induces the Cartesian product $K_{|V_A|} \square \langle W \rangle$, where $\langle W \rangle$ denotes the subgraph of G_A induced by W , and $V_r \times \{V_c \setminus W\}$ induces $G_A \square \langle V_c \setminus W \rangle$.

It is easy to see that $A \times A \subseteq \text{Aut}(G)$. We have to prove the converse. To that end it suffices to show that $\text{Aut}(G)$ maps rows onto rows and columns onto columns.

Consider a row $V_r \times \{v_c\}$, where $v_c \in W$. The row induces a complete subgraph. Because we have no slant edges, automorphisms can only map it into rows or columns. As all rows and columns have the same number of vertices and since $G_A \neq K_{|V_A|}$, it can only be mapped onto a $V_r \times \{w_c\}$, where $w_c \in W$.

We will now prove that automorphisms of G map columns onto columns. Pick a $v_c \in W$ to single out one of the rows of W , and let (w_r, w_c) be any vertex of G . As there are no slant edges in G , the paths realizing the weak distance of (w_r, w_c) to points (v_r, v_c) in the chosen row will be built of column edges and row edges. By analogy to the reasoning behind the distance formula for the cartesian product, the column edges of any such path projected onto the column graph G_c will form a weak path from w_c to v_c in G_c , just as in a cartesian product, but the row edges can go through regular rows or through $K_{|V_A|}$ rows. When v_r equals w_r , row edges are eliminated. That means that given a vertex (w_r, w_c) there is a unique vertex in the chosen row $V_r \times \{v_c\}$, to which weak distance ρ in G is minimal, this unique vertex (w_r, v_c) is in the same column as (w_r, w_c) and is unique in the above sense for all vertices (w_r, w_c) of that column.

Consider now an automorphism $a \in \text{Aut}(G)$. We already know that it will map the row $V_r \times \{v_c\}$ onto some other row $V_r \times \{x_c\}$. If the vertices $(x_r, x_c) = a(w_r, v_c)$ and $(y_r, y_c) = a(w_r, w_c)$ were in different columns, that is if $x_r \neq y_r$, there would be a vertex (y_r, x_c) in row x_c closest to (y_r, y_c) and different than (x_r, x_c) :

$$\rho((x_r, x_c), (y_r, y_c)) > \rho((y_r, x_c), (y_r, y_c)),$$

while after having applied a^{-1} on both sides we would get

$$\rho((w_r, v_c), (w_r, w_c)) > \rho((w'_r, v_c), (w_r, w_c)),$$

with $w'_r \neq w_r$ because of $x_r \neq y_r$, but that cannot be true. Hence, any automorphism maps columns onto columns, as vertices of G follow their closest vertices in the chosen row.

Since column edges are mapped by automorphisms onto column edges, row edges are mapped only to row edges, thus, the only way the image of a row can preserve its weak connectedness is for automorphisms to map entire rows onto entire rows. $\square$

Lemma 2.5. *Let $A \in \text{EDGR}$ with prime G_A . If A is transitive and $|V_A| \leq 4$, then $A \times A \in \text{EDGR}$ unless $A = C_3$.*

Proof. The group A is one of C_3 and C_4 . By a result of Babai [1], $C_3 \times C_3 \notin \text{EDGR}$. $C_4 \times C_4 \in \text{EDGR}$ by Theorem 1.4 for G_{C_4} and $\overline{G_{C_4}}$. $\square$

Observe that this takes care of the last exceptional case of Theorem 2.1.

Lemma 2.6. *Let $A \in \text{EDGR}$ with prime G_A , where A is transitive and $|V_A| > 4$. If $\overline{G_A}$ is weakly connected, then $A \times A \in \text{EDGR}$.*

Proof. Denote $n = |V_A|$. Since the graph $\overline{G_A}$ is weakly connected, we only need to consider the case $G_A \cong \overline{G_A}$ (otherwise the conclusion follows from Theorem 1.4). This implies that $d^0(G_A) = d^1(G_A)$. Because G_A is not undirected, we infer that $2d^f(G_A) > 1$. Then $d^0(G_A) + d^1(G_A) + 2d^f(G_A) = n - 1$ implies $2d^1(G_A) = 2d^0(G_A) < n - 2$.

We shall now prove that the graph $G = G_r * G_c$, where $G_r = (V_r, E_r)$ and $G_c = (V_c, E_c)$ are copies of G_A , has the property $\text{Aut}(G) = A \times A$. To this end, we will show that every undirected edge that is contained in a row is mapped, under the action of $\text{Aut}(G)$, onto an undirected edge which is contained in a row, and that the same is true for directed edges.

Let us compare the numbers of the common 1-neighbors of the ends of an undirected edge which is contained in a row, with the same number for the ends of an undirected slant edge. Denote the ends of the edge e by (v_r, v_c) and (w_r, w_c) . If e is contained in a row ($v_c = w_c$), then the common 1-neighbors of (v_r, v_c) and (w_r, w_c) are those contained in that row, together with all but two vertices in rows corresponding to 1-neighbors of $v_c = w_c$ in G_c , hence their number is equal to

$$N_{G_r}^1(v_r, w_r) + (n - 2)d^1(G_c), \quad (2.1)$$

where $N_{G_r}^1(v_r, w_r)$ is the number of common 1-neighbors of the vertices v_r and w_r (in G_r). If e is a slant edge, the common 1-neighbors of (v_r, v_c) and (w_r, w_c) are the 1-neighbors contained in both rows (excluding the vertex directly in front of the other end if it also is such a 1-neighbor), together with all but two vertices in rows corresponding to common 1-neighbors of both v_c and w_c in G_c . Thus, their number is

$$(n - 2)N_{G_c}^1(v_c, w_c) + 2d^1(G_r) - 2\delta, \quad (2.2)$$

where $N_{G_c}^1(v_c, w_c)$ is the number of common 1-neighbors of the vertices v_c and w_c (in G_c), and $\delta \in \{0, 1\}$.

The assumption that the numbers (2.1) and (2.2) are equal, implies

$$(n - 2)(d^1(G_c) - N_{G_c}^1(v_c, w_c)) + N_{G_r}^1(v_r, w_r) - 2d^1(G_r) + 2\delta = 0.$$

Since $d^1(G_c) > N_{G_c}^1(v_c, w_c)$ and $2d^1(G_r) < n - 2$, it cannot be true. Hence, an undirected edge which is contained in a row cannot be mapped onto a slant undirected edge. Since there are no undirected edges in columns of a $*$ -product, the set of the undirected edges that are contained in the rows is preserved by automorphisms.

We continue with a similar calculation for directed edges. Let e be a directed edge with ends as above. If e is contained in a row, then by similar reasoning as in the undirected case, the number of common forward-neighbors of (v_r, v_c) and (w_r, w_c) equals

$$N_{G_r}^f(v_r, w_r) + (n - 2)d^f(G_c), \quad (2.3)$$

where $N_{G_r}^f(v_r, w_r)$ is the number of common forward-neighbors of the vertices v_r and w_r (in G_r). If e is a slant edge, then this number is

$$(n - 2)N_{G_c}^f(v_c, w_c) + d^f(G_r) - \delta, \quad (2.4)$$

where $N_{G_c}^f(v_c, w_c)$ is the number of common forward-neighbors of the vertices v_c and w_c (in G_c), and $\delta \in \{0, 1\}$.

If it were possible for an automorphism from $\text{Aut}(G)$ to map a directed slant edge onto a directed row edge, the numbers (2.3) and (2.4) would need to be the same, which would imply

$$(n-2)(d^f(G_c) - N_{G_c}^f(v_c, w_c)) + N_{G_r}^f(v_r, w_r) - d^f(G_r) + \delta = 0. \quad (2.5)$$

If $e = ((v_r, v_c), (w_r, w_c))$ is a directed slant edge, then (v_c, w_c) is a directed edge of G_c . The equality $d^f(G_c) = N_{G_c}^f(v_c, w_c)$ would then imply that the set of forward-neighbors of each of the vertices v_c and w_c be identical, but this cannot be true, since w_c is a forward-neighbor of v_c but not of itself. Hence, $d^f(G_c) > N_{G_c}^f(v_c, w_c)$. Note that since A is transitive, every vertex has as many backward neighbours as forward neighbours. Therefore since $n > 4$, we infer $d^f(G_r) < n - 2$. Thus, equation (2.5) cannot be true and the set of directed edges that are contained in rows is preserved by automorphisms also in this case. Because G_A is weakly connected, it follows by Proposition 1.6 that $\text{Aut}(G) = A \times A$. $\square$

Lemma 2.7. *Let $A \in \text{EDGR}$, with prime G_A , be transitive. If $\overline{G_A}$ is disconnected, then $A \times A \in \text{EDGR}$.*

Proof. We first consider the structure of $\overline{G_A}$. Because A is transitive, the subgraphs of $\overline{G_A}$ induced by the vertices belonging to common weakly connected components of $\overline{G_A}$ are isomorphic, so $V_A = W' \times W$, where the weakly connected components of $\overline{G_A}$ are grouped as columns, with column size $s = |W| = n/t$, where $t = |W'| \geq 2$ is the number of weakly connected components of $\overline{G_A}$. Thus, the group A acts on the set of columns as S_t , and on every column independently as some A_1 , hence $A = S_t \text{ wr } A_1$. Since there are no edges between columns of $\overline{G_A}$ we infer that $(v, w) \in E(G_A)$ if v and w belong to different columns. Because A is transitive, and G_A is not undirected, we conclude that either $s \geq 4$ or $A_1 = C_3$.

In the latter case, we define $G = G_r * G_c$, where both G_r and G_c are isomorphic to G_A . Then, it is easy to see that the ends of the undirected edges in the rows have common forward-neighbors, and the ends of the undirected slant edges do not. Since the undirected edges in rows form spanning connected subgraphs of the rows, $\text{Aut}(G)$ maps rows onto rows. By Proposition 1.6 we conclude that $\text{Aut}(G) = A \times B$.

In the case $s \geq 4$, we define a graph $G = (V_r \times V_c, E)$ such that $((v_r, v_c), (w_r, w_c))$ is in E if either $v_c = w_c$ and $(v_r, w_r) \in E(G_r)$ or $(v_c, w_c) \in E(G_c)$, $v_r \neq w_r$, and the vertices v_r and w_r belong to the same weakly connected component in $\overline{G_r}$.

If a connected graph H has a disconnected complement, then the subgraphs of H that are induced by the vertices of the weakly connected components of $\overline{H}$ are sometimes called *Zykov components* of H . Our graph G thus consists of t copies of the $R * G_c$, where R is a Zykov-component of G_r , and the row-edges that are not in a copy of $R * G_c$. We say these row-edges are of type Q .

We wish to show that $\text{Aut}(G) = A \times A$. It is easy to check that $A \times A \subseteq \text{Aut}(G)$. We have to prove that the converse also holds. To this end, we count the common weak neighbors of the ends of the edges that are contained in a row. These edges have the form $\{(v_r, v_c), (w_r, w_c)\}$, where $v_c = w_c$. If v_r and w_r do not belong to the same Zykov component in G_r , then these edges are of type Q . The number of common weak neighbors of the endpoints of edges of type Q is

$$x = (t-2)s + 2d_W, \quad (2.6)$$

where d_W is the number of those weak neighbors of a vertex in G_r that belong to the same Zykov component of G_r . (The notation d_W is chosen, because all Zykov components are isomorphic to W as defined in the beginning of the proof.) For row-edges that are not of type Q the number of common weak neighbors of their endpoints is

$$y = (t-1)s + N_W(v_r, w_r) + (s-2)((t-1)s + d_W), \quad (2.7)$$

where $N_W(v_r, w_r)$ is the number of the common weak neighbors of the vertices v_r and w_r in their Zykov component. Since $x < (t-1)s + d_W$, it is obvious that $x < y$.

Moreover, the number of common weak neighbors of the ends of the slant edges of G is

$$2(d_W - \epsilon) + (s-2)N_W(v_c, w_c) + (s-2)(t-1)s$$

for some $\epsilon \in \{0, 1\}$ if the endpoints of $\{v_c, w_c\} \in E(G_c)$ belong to the same Zykov component of G_c , and

$$2(d_W - \epsilon) + 2(s-2)d_W + (s-2)(t-2)s$$

for some $\epsilon \in \{0, 1\}$ if the endpoints of $\{v_c, w_c\} \in E(G_c)$ belong to different Zykov components of G_c . It is easy to see that under our assumptions both numbers are strictly greater than x . Observe that the graph G has no edges that are contained in its columns.

This calculation implies that $\text{Aut}(G)$ preserves the set of edges of type Q . Since these edges form spanning subgraphs for all graphs induced by the rows of G , every $a \in \text{Aut}(G)$ maps rows of G onto rows. Moreover, a maps any copy of a Zykov component of G_r that is contained in a row in G onto a copy of a Zykov component of G_r that is contained in the image of that row.

To complete the proof, we have to show that each column of G is mapped onto a column. If we remove edges of type Q we are left with t identical subgraphs $R_i * G_c$ where $i = 1, \dots, t$. As any automorphism a of G maps rows into rows, it also maps subrows of the form $R_i \times \{v_c\}$ into subrows of the same form $R_j \times \{a(v_c)\}$.

Note that by assumption R has $s \geq 4$ vertices. Thus, every $R_i * G_c$ is weakly connected. From this we infer that automorphisms of G map entire subgraphs $R_i * G_c$ onto entire subgraphs $R_j * G_c$, as in G there are no slant edges between vertices belonging to different Zykov components.

Call subrows $R_i \times \{v_c\}$ and $R_j \times \{w_c\}$ of G adjacent if there is an edge or directed edge between some vertices of them, that is, when there is an edge or a directed edge between v_c and w_c in G_c and $i = j$.

If $R_i \times \{v_c\}$ and $R_i \times \{w_c\}$ are adjacent, so are $R_j \times \{a(v_c)\}$ and $R_j \times \{a(w_c)\}$ and the non-edges between vertices of the subrows are mapped to non-edges between vertices of the images of the subrows. But the non-edges of adjacent subrows span subgraphs that are isomorphic to copies of G_c and whose vertex sets are the columns. Therefore, columns of G are mapped onto columns. $\square$

This also completes the proof of the main theorem.

References

- [1] L. Babai, Finite digraphs with given regular automorphism groups, *Period. Math. Hung.* **11** (1980), 257–270, doi:10.1007/bf02107568.

- [2] L. Babai, Automorphism groups, isomorphism, reconstruction, in: R. L. Graham, M. Grötschel and L. Lovász (eds.), *Handbook of Combinatorics, Volume 2*, Elsevier, Amsterdam, pp. 1447–1540, 1995.
- [3] A. Bonisoli and D. Labbate, One-factorizations of complete graphs with vertex-regular automorphism groups, *J. Combin. Des.* **10** (2002), 1–16, doi:10.1002/jcd.1025.
- [4] J. Feigenbaum, Directed Cartesian-product graphs have unique factorizations that can be computed in polynomial time, *Discrete Appl. Math.* **15** (1986), 105–110, doi:10.1016/0166-218x(86)90023-5.
- [5] R. Frucht, Herstellung von Graphen mit vorgegebener abstrakter Gruppe, *Compositio Math.* **6** (1939), 239–250, http://www.numdam.org/item?id=CM_1939__6__239_0.
- [6] M. Giudici, C. H. Li, P. Potočník and C. E. Praeger, Homogeneous factorisations of graphs and digraphs, *European J. Combin.* **27** (2006), 11–37, doi:10.1016/j.ejc.2004.08.003.
- [7] C. D. Godsil, GRRs for nonsolvable groups, in: L. Lovász and V. T. Sós (eds.), *Algebraic Methods in Graph Theory, Volume I*, North-Holland, Amsterdam, volume 25 of *Colloquia Mathematica Societatis János Bolyai*, pp. 221–239, 1981, papers from the conference held in Szeged, August 24 – 31, 1978.
- [8] M. Grech, Direct products of automorphism groups of graphs, *J. Graph Theory* **62** (2009), 26–36, doi:10.1002/jgt.20385.
- [9] M. Grech, The graphical complexity of direct products of permutation groups, *J. Graph Theory* **66** (2011), 303–318, doi:10.1002/jgt.20504.
- [10] M. Grech, A. Jež and A. Kisielwicz, Graphical complexity of products of permutation groups, *Discrete Math.* **308** (2008), 1142–1152, doi:10.1016/j.disc.2007.04.004.
- [11] M. Grech and A. Kisielwicz, Direct product of automorphism groups of colored graphs, *Discrete Math.* **283** (2004), 81–86, doi:10.1016/j.disc.2003.11.008.
- [12] M. Grech and A. Kisielwicz, Totally symmetric colored graphs, *J. Graph Theory* **62** (2009), 329–345, doi:10.1002/jgt.20397.
- [13] R. Hammack, W. Imrich and S. Klavžar, *Handbook of Product Graphs*, Discrete Mathematics and its Applications, CRC Press, Boca Raton, Florida, 2nd edition, 2011, doi:10.1201/b10959.
- [14] W. Imrich, On products of graphs and regular groups, *Isr. J. Math.* **11** (1972), 258–264, doi:10.1007/bf02789317.
- [15] W. Imrich, On graphical regular representations of groups, in: A. Hajnal, R. Rado and V. T. Sós (eds.), *Infinite and Finite Sets, Volume II*, North-Holland, Amsterdam, volume 10 of *Colloquia Mathematica Societatis János Bolyai*, pp. 905–925, 1975, dedicated to Paul Erdős on his 60th birthday, proceedings of a colloquium held at Keszthely, June 25 – July 1, 1973.
- [16] W. Imrich, Graphical regular representations of groups of odd order, in: A. Hajnal and V. T. Sós (eds.), *Combinatorics, Volume II*, North-Holland, Amsterdam, volume 18 of *Colloquia Mathematica Societatis János Bolyai*, pp. 611–621, 1978, proceedings of the 5th Hungarian Colloquium held at Keszthely, June 28 – July 3, 1976.
- [17] W. Imrich and I. Peterin, Cartesian products of directed graphs with loops, *Discrete Math.* **341** (2018), 1336–1343, doi:10.1016/j.disc.2018.01.021.
- [18] W. Imrich and M. E. Watkins, On graphical regular representations of cyclic extensions of groups, *Pac. J. Math.* **55** (1974), 461–477, <http://projecteuclid.org/euclid.pjm/1102910980>.
- [19] A. Kisielwicz, Supergraphs and graphical complexity of permutation groups, *Ars Combin.* **101** (2011), 193–207.

- [20] D. König, *Theory of Finite and Infinite Graphs*, Birkhäuser, Boston, Massachusetts, 1990, doi:10.1007/978-1-4684-8971-2.
- [21] C. H. Li, T. K. Lim and C. E. Praeger, Homogeneous factorisations of complete graphs with edge-transitive factors, *J. Algebr. Combin.* **29** (2009), 107–132, doi:10.1007/s10801-008-0127-2.
- [22] C. H. Li and C. E. Praeger, On partitioning the orbitals of a transitive permutation group, *Trans. Am. Math. Soc.* **355** (2003), 637–653, doi:10.1090/s0002-9947-02-03110-0.
- [23] L. A. Nowitz, On the non-existence of graphs with transitive generalized dicyclic groups, *J. Comb. Theory* **4** (1968), 49–51, doi:10.1016/s0021-9800(68)80086-9.
- [24] L. A. Nowitz and M. E. Watkins, Graphical regular representations of non-abelian groups, I, *Canad. J. Math.* **24** (1972), 993–1008, doi:10.4153/cjm-1972-101-5.
- [25] L. A. Nowitz and M. E. Watkins, Graphical regular representations of non-abelian groups, II, *Canad. J. Math.* **24** (1972), 1009–1018, doi:10.4153/cjm-1972-102-3.
- [26] G. Sabidussi, Graph multiplication, *Math. Z.* **72** (1959/60), 446–457, doi:10.1007/bf01162967.
- [27] V. G. Vizing, The cartesian product of graphs (in Russian), *Vychisl. Sistemy* **9** (1963), 30–43.
- [28] M. E. Watkins, On the action of non-Abelian groups on graphs, *J. Comb. Theory Ser. B* **11** (1971), 95–104, doi:10.1016/0095-8956(71)90019-0.
- [29] M. E. Watkins, On graphical regular representations of $C_n \times Q$, in: Y. Alavi, D. R. Lick and A. T. White (eds.), *Graph Theory and Applications*, Springer, Berlin, volume 303 of *Lecture Notes in Mathematics*, pp. 305–311, 1972, doi:10.1007/bfb0067383, dedicated to the memory of J. W. T. Youngs, proceedings of the Conference at Western Michigan University, Kalamazoo, Michigan, May 10 – 13, 1972.
- [30] M. E. Watkins, Graphical regular representations of alternating, symmetric, and miscellaneous small groups, *Aequat. Math.* **11** (1974), 40–50, doi:10.1007/bf01837731.
- [31] M. E. Watkins and L. A. Nowitz, On graphical regular representations of direct products of groups, *Monatsh. Math.* **76** (1972), 168–171, doi:10.1007/bf01298284.
- [32] H. Wielandt, Permutation groups through invariant relations and invariant functions, in: B. Huppert and H. Schneider (eds.), *Group Theory*, Walter de Gruyter, Berlin, volume 1 of *Mathematische Werke/Mathematical Works*, pp. 237–297, 1994.
- [33] H. P. Yap, *Some Topics in Graph Theory*, volume 108 of *London Mathematical Society Lecture Note Series*, Cambridge University Press, Cambridge, 1986, doi:10.1017/cbo9780511662065.
- [34] A. Z. Zelikovskiy, The König problem for Abelian permutation groups, *Proc. Natl. Acad. Sci. Belarus, Ser. Phys.-Math. Sci.* **5** (1989), 34–39.