

Top-heavy phenomena for transformations*

Yaokun Wu [†] , Yinfeng Zhu 

*School of Mathematical Sciences and MOE-LSC, Shanghai Jiao Tong University,
Shanghai 200240, China*

Received 19 July 2018, accepted 5 February 2022, published online 19 August 2022

Abstract

Let S be a transformation semigroup acting on a set Ω . The action of S on Ω can be naturally extended to be an action on all subsets of Ω . We say that S is ℓ -homogeneous provided it can send A to B for any two (not necessarily distinct) ℓ -subsets A and B of Ω . On the condition that $k \leq \ell < k + \ell \leq |\Omega|$, we show that every ℓ -homogeneous transformation semigroup acting on Ω must be k -homogeneous. We report other variants of this result for Boolean semirings and affine/projective geometries. In general, any semigroup action on a poset gives rise to an automaton and we associate some sequences of integers with the phase space of this automaton. When this poset is a geometric lattice, we propose to investigate various possible regularity properties of these sequences, especially the so-called top-heavy property. In the course of this study, we are led to a conjecture about the injectivity of the incidence operator of a geometric lattice, generalizing a conjecture of Kung.

Keywords: Incidence operator, kernel space, rank, strong shape, valuated poset, weak shape.

Math. Subj. Class. (2020): 05B35, 20M20, 47D03, 52C35

1 Introduction

1.1 Transformation and phase space

Let Γ be a *digraph*, namely a pair consisting of its vertex set $V(\Gamma)$ and arc set $E(\Gamma) \subseteq V(\Gamma) \times V(\Gamma)$. We call Γ *symmetric* if $(u, v) \in E(\Gamma)$ holds if and only if so does $(v, u) \in E(\Gamma)$. For any $A \subseteq V(\Gamma)$, we adopt the notation $\Gamma[A]$ for the subdigraph of Γ induced by

*We thank Peter Cameron, Alexander Ivanov, Peter Šemrl, Johannes Siemons and Qing Xiang for useful discussions. Especially, Peter Šemrl reminded us the result of Taussky and Zassenhaus while Johannes Siemons suggested that the isomorphism of two permutation modules can be read from Figure 6. This work has been supported by NSFC (11971305, 11671258) and STCSM (17690740800).

[†]Corresponding author.

E-mail addresses: ykwu@sjtu.edu.cn (Yaokun Wu), fengzi@sjtu.edu.cn (Yinfeng Zhu)

A which has vertex set A and arc set $E(\Gamma) \cap (A \times A)$. The number of weakly connected components and the number of strongly connected components of Γ will be dubbed $wcc(\Gamma)$ and $scc(\Gamma)$, respectively.

For a set Ω , all maps from Ω to itself form the set Ω^Ω . For each $g \in \Omega^\Omega$ and $\alpha \in \Omega$, we write αg for the image of α under the map g . The composition of maps provides an associative product on the set Ω^Ω and thus turns it into a monoid, namely a semigroup with a multiplicative unit. We call this monoid the *full transformation monoid* on Ω and denote it by $T(\Omega)$. A subset of $T(\Omega)$ which is closed under map composition, whether or not it contains the identity map on Ω , is called a *transformation semigroup* acting on Ω . Let S be a transformation semigroup on Ω . We say that S is *transitive on a set* $A \subseteq \Omega$ if for every $\alpha, \beta \in A$ we can find $g \in S$ such that $\alpha g = \beta$; we call S *transitive* if S is transitive on Ω . If the transformation semigroup S is generated by a set $G \subseteq \Omega^\Omega$, namely S consists of products of elements of G of positive length, we call (S, G) a *deterministic automaton* on Ω [67, Section 1]. The *phase space* of an automaton (S, G) on Ω , denoted by $\Gamma(S, G)$, is the digraph with vertex set Ω and arc set $\{(\alpha, \alpha g) : \alpha \in \Omega, g \in G\}$. When Ω has at least two elements, the claim that S is transitive is equivalent to the claim that $\Gamma(S, G)$ is strongly connected for any generator set G of S . We write $\Gamma(S, S)$ simply as $\Gamma(S)$ and note that each strongly/weakly connected component of $\Gamma(S)$ coincides with a strongly/weakly connected component of $\Gamma(S, G)$ for any generator set G of S . For all work in this paper, we can simply focus on $\Gamma(S)$ instead of considering $\Gamma(S, G)$ for any specific generator set G . We emphasize $\Gamma(S, G)$ from the phase space viewpoint here to highlight the connection between semigroup theory and automata theory, and to indicate the role played by the choice of G in some problems related to various distance functions on the phase space, say the Černý conjecture. For any set Ω , a subset of $T(\Omega)$ forms a *permutation group* on Ω whenever it is a transformation semigroup and each element has an inverse in it, namely it is a set of bijective transformations of Ω and is closed under compositions and taking inverses. Permutation groups correspond to reversible deterministic automata.

Let Ω be a set. We follow the common practice to use 2^Ω for the power set of Ω . For each $g \in T(\Omega)$, let \bar{g} be the element in $T(2^\Omega)$ that sends each $A \in 2^\Omega$ to $A\bar{g} \doteq \{ag : a \in A\}$. More generally, for each $G \subseteq T(\Omega)$, \bar{G} refers to the set $\{\bar{g} : g \in G\}$. For any transformation semigroup S on Ω and any generator set G of S , \bar{S} , as a semigroup derived from S , is known to be the *powerset transformation semigroup of S* acting on 2^Ω and (\bar{S}, \bar{G}) is known to be the *powerset automaton of (S, G)* . It may be interesting to iterate the powerset automaton construction and examine the evolution of the phase spaces of the resulting automata.

When discussing transformation semigroups, we may often be more interested in those which preserve some structures, say simplicial maps for simplicial complexes, continuous maps for topological spaces, ordering preserving maps for posets, or adjacency-preserving maps in matrix geometry [52, 66]. Unlike the work on group actions on posets [3, 59] and matroids [19], very little has been done on semigroup actions on these structures [62]. Moving from group actions to semigroup actions is just to consider general deterministic automata instead of reversible ones.

1.2 Valuated poset and its shape

For any two sets Ω and Ψ , if they are different or if we do not emphasize that they may be equal, the image of $\omega \in \Omega$ under a map $g \in \Psi^\Omega$ is denoted $g(\omega)$; note that we often write it as ωg when $\Omega = \Psi$.

A poset P consists of a set Ω and a binary relation $<_P$ on it which is transitive and acyclic, namely we require that $\alpha <_P \alpha$ never happens, and that $\alpha <_P \beta$ and $\beta <_P \gamma$ implies $\alpha <_P \gamma$ for all $\alpha, \beta, \gamma \in \Omega$. We often just write P for its ground set Ω and we say the poset P is *finite* if $|P|$ is finite. For each $\alpha \in P$, the *principal ideal* generated by α is the set $\{\beta : \beta <_P \alpha\} \cup \{\alpha\} \subseteq P$, which we denote by $P_\downarrow(\alpha)$; the *principal filter* generated by α is the set $\{\beta : \alpha <_P \beta\} \cup \{\alpha\} \subseteq P$, which we denote by $P_\uparrow(\alpha)$. An *ideal (filter)* is a union of principal ideals (filters). A map g from a poset P to a poset Q is *order-preserving* if $g(\beta) \in Q_\downarrow(g(\alpha))$ holds whenever $\beta \in P_\downarrow(\alpha)$. We use $\text{End}(P)$ to denote the set of all order-preserving maps from P to itself.

Let $\mathbb{Z}_{\geq 0}$ be the set of nonnegative integers which carries a natural poset structure such that $a < b$ in $\mathbb{Z}_{\geq 0}$ if and only if $b - a$ is a positive integer. A *valuation* on a poset P is an order-preserving map r_P from P to the poset $\mathbb{Z}_{\geq 0}$; we call $r_P(x)$ the rank of x in the valuated poset. When we say P is a *valuated poset*, we are considering the poset P together with a valuation r_P , though the valuation may be only implicitly indicated. The *rank* of a valuated poset P , denoted by $r(P)$, is the maximum value of $r_P(\alpha)$ for $\alpha \in P$ if it exists and is ∞ otherwise. For a poset P , the symbols like $<_P$ and r_P will often be abbreviated to $<$ and r when no confusion can arise. Let P be a valuated poset. For any $k \in \mathbb{Z}_{\geq 0}$, we write P_k for the set $\{\alpha \in P : r(\alpha) = k\}$. We call the sequence $|P_0|, |P_1|, \dots$ the *shape* of the valuated poset and refer to it by $S(P)$. If $r(P) < \infty$, $S(P)$ is a sequence of $r(P) + 1$ nonnegative integers.

Let P be a valuated poset and let S be a subsemigroup of $\text{End}(P)$. The *weak shape* of P under the action of S is the sequence

$$\text{wcc}(\Gamma(S)[P_0]), \text{wcc}(\Gamma(S)[P_1]), \dots$$

which we denote by $\text{WS}(S, P)$; while the *strong shape* of P under the action of S is the sequence

$$\text{scc}(\Gamma(S)[P_0]), \text{scc}(\Gamma(S)[P_1]), \dots$$

which we denote by $\text{SS}(S, P)$. Note that

$$S(P) = \text{WS}(S, P) = \text{SS}(S, P)$$

when the semigroup S consists of the identity transformation from $\text{End}(P)$.

The main purpose of this note is to propose a study of the possible regularity in the strong/weak shape of a semigroup acting on a valuated poset.

1.3 Geometric lattice and top-heavy property

A matroid M consists of a ground set \mathcal{E}_M and a rank function r_M from $2^{\mathcal{E}_M}$ to the set of nonnegative integers plus infinity such that the rank axioms are satisfied [13, Section 1.5]. The flats of a matroid M , ordered by inclusion, form a very pretty structure, called the *matroid lattice* of M and denoted by $F(M)$. For each nonnegative integer t , let $F_t(M)$ be the set of all rank- t flats of the matroid M . A *geometric lattice* is an atomic and semimodular lattice which does not have any infinite chain [64, page 305]. We mention that a geometric lattice is cryptomorphic to a natural object called combinatorial geometry [64, Theorem 23.1] and that finite geometric lattice is nothing but finite matroid lattice [36, page 163, Birkhoff's Theorem]. A geometric/matroid lattice has a natural valuated poset structure, where the valuation is given by its rank function. For example, for a matroid M , all elements in $F_t(M)$ have rank t . In a geometric lattice, the elements of rank 1,

2 and 3 are viewed as points, lines and planes, respectively, thus giving geometric intuitions to many results about geometric lattices.

For each linear space V and each nonnegative integer k , we use $\text{Gr}(k, V)$ for the set of all k -dimensional linear subspaces of V and we call $\bigcup_{k=0}^{\infty} \text{Gr}(k, V)$ the *Grassmannian* of V , which is denoted by $\text{Gr}(V)$. If V is finite dimensional, $\text{Gr}(V)$ is surely a geometric lattice with elements from $\text{Gr}(k, V)$ having rank k .

Example 1.1. Let n and k be two positive integers such that $k < n$. Fix a non-degenerate inner product on \mathbb{Q}^n , say $\langle \cdot, \cdot \rangle$. For each $g \in \text{GL}_n(\mathbb{Q})$, let g^\top stand for the adjoint of g , namely the element such that $\langle ug, v \rangle = \langle u, vg^\top \rangle$ for all $u, v \in \mathbb{Q}^n$, and we write $g_\#$ for $(g^{-1})^\top$. Let $S \leq \text{GL}_n(\mathbb{Q})$ be a matrix group acting on \mathbb{Q}^n . If \bar{S} is transitive on the set of all dimension- k subspaces and if $g_\# \in S$ for all $g \in S$, then \bar{S} is transitive on the set of dimension- $(n - k)$ subspaces. To see this, fix a pair of subspaces (U, U') which are orthogonal complements to each other with respect to $\langle \cdot, \cdot \rangle$ and $(\dim U, \dim U') = (k, n - k)$. For each $g \in S$, we can see that $U\bar{g}$ and $U'\bar{g}_\#$ are orthogonal complements to each other with respect to the given inner product $\langle \cdot, \cdot \rangle$. Considering the set of pairs $\{(U\bar{g}, U'\bar{g}_\#) : g \in S\}$, we see that the transitivity on $\text{Gr}(k, \mathbb{Q}^n)$ implies transitivity on $\text{Gr}(n - k, \mathbb{Q}^n)$.

Motivated by Example 1.1, here is a very simple question on the very simple geometric lattice $\text{Gr}(\mathbb{Q}^3)$. Surprisingly, we even could not find any discussion of it in the literature.

Question 1.2. If S is a general matrix group acting on \mathbb{Q}^3 , can we draw the conclusion that \bar{S} is transitive on $\text{Gr}(1, \mathbb{Q}^3)$ from the assumption of its transitivity on $\text{Gr}(2, \mathbb{Q}^3)$? What about only assuming that S is a matrix semigroup?

Some seemingly weird properties of sequences turn out to be ubiquitous when we are examining some interesting structures or processes [6, 10, 11, 29, 58, 61]. We review some of them below. Let c_0, c_1, \dots , be a sequence of $n + 1$ real numbers, where n can be finite or infinite. We call it *t-top-heavy* if $c_k \leq t$ whenever there exists an integer ℓ such that $k \leq \ell \leq k + \ell \leq n$ and $c_\ell \leq t$; we call it *top-heavy* if it is *t-top-heavy* for all $t \in \mathbb{R}$, namely $c_k \leq c_\ell$ holds for all k, ℓ such that $k \leq \ell \leq k + \ell \leq n$; We call it *unimodal* if you cannot find three distinct integers i, j, k such that $0 \leq i < j < k \leq n$ and $c_i - c_j > 0 > c_j - c_k$; we call it *log-concave* if $c_i^2 \geq c_{i-1}c_{i+1}$ for all $i = 1, \dots, n - 1$. When n is finite, we call the sequence *real-rooted* provided the polynomial $c_0 + c_1x + \dots + c_nx^n$ in the unknown x only has real roots and we call it *ultra-log-concave* provided $\frac{c_0}{\binom{n}{0}}, \dots, \frac{c_n}{\binom{n}{n}}$ forms a log-concave sequence. Note that Question 1.2 is about the possible 1-top-heavy property of the strong shape of $\text{Gr}(\mathbb{Q}^3)$ under a matrix semigroup action.

In the 1970s, two log-concavity conjectures [61, Conjecture 3] appeared in combinatorics community which claim that the sequences of Whitney numbers of both the first kind and the second kind of a finite matroid are log-concave. The first conjecture was verified by Adiprasito, Huh and Katz [1]. Mason [40] had made variants and stronger versions of the second conjecture; but even the original conjecture is still open. Dowling and Wilson [23] conjectured that the sequence of Whitney numbers of the second kind of a finite matroid is top-heavy. When restricted to finite realizable matroids, this top-heavy conjecture was proved by Huh and Wang [28]. The second log-concavity conjecture as described above, which is about the Whitney numbers of the second kind [50], simply says that the shape of every geometric lattice is log-concave. The above-mentioned Dowling-Wilson top-heavy conjecture says that the shape of every finite geometric lattice is top-heavy. On

the condition that these two conjectures are both true, we know that the shape of a finite geometric lattice is both log-concave (and hence unimodal) and top-heavy. Can we draw this conclusion for the strong/weak shape of some semigroup actions on some geometric lattices?

Boolean lattices, partition lattices and projective/affine geometries are some most well-known geometric lattices. It is easy to see that their shapes are all ultra-log-concave (and hence real-rooted) and top-heavy [37]. The main result of this paper, Theorems 2.1 and 2.12, declare the top-heavy property for the strong/weak shape of some semigroups acting on Boolean lattices and projective/affine geometries. The semigroups considered by us are those derived from “simple” transformations. We also report our attempt at tackling the same problem for partition lattices and the Vámos matroid.

In Section 2, we will present our main results as well as pertinent problems, examples, and remarks. The first three subsections are devoted to Boolean lattices, partition lattices and projective/affine geometries. The last subsection is a simple discussion in the context of matroids. Before digging into the proofs of the main results, we develop some technical tools in Section 3. In the sequel, we provide in Sections 4 to 7 all the proofs missing from Sections 2.1 to 2.4. We conclude the paper in Section 8 with a brief discussion of the present work and some possible further research.

2 A top-heavy promenade

2.1 Boolean semiring and homogeneity

For any set Ω , the set $B_\Omega \doteq \bigcup_{k=0}^{\infty} \binom{\Omega}{k}$ forms a poset under the inclusion relationship, which is often known as the *Boolean semiring over Ω* , and the set 2^Ω gives rise to the *Boolean algebra over Ω* . When we view B_Ω as a valuated poset, unless stated otherwise, the valuation will be $r(A) = |A|$ for all $A \in B_\Omega$. If Ω is a finite set, B_Ω coincides with 2^Ω and is referred to as a *Boolean lattice*.

Let A and Ω be two sets with $A \subseteq \Omega$. For any $g \in \Omega^\Omega$, write $g|_A$ for the restriction of g on A . Let S be a transformation semigroup on Ω . For any positive integer $k \leq |\Omega|$, we name S *k-homogeneous* if the transformation semigroup \bar{S} is transitive on $\binom{\Omega}{k}$, that is, $\text{scc}(\Gamma(\bar{S})[\binom{\Omega}{k}]) = 1$. The *stabiliser permutation group* of (S, A) is the permutation group $S_A \doteq \{g|_A : g \in S, A\bar{g} = A\}$ acting on A . The *relative transformation semigroup* of (S, A) is the transformation semigroup $\tilde{S}_A \doteq \{g|_A : g \in S, A\bar{g} \subseteq A\}$ acting on A . Note that the action of \tilde{S}_A on A may not be transitive even if S acts on A transitively.

Theorem 2.1. *Let Ω be a set of size n . Let S be a transformation semigroup on Ω and let Γ be the phase space of \bar{S} .*

- (1) $\text{SS}(\bar{S}, B_\Omega)$ is 1-top-heavy.
- (2) Both $\text{WS}(\bar{S}, B_\Omega)$ and $\text{SS}(\bar{S}, B_\Omega)$ are top-heavy.
- (3) Let k and ℓ be two integers such that $0 \leq k \leq \ell \leq k + \ell \leq n + 1$. Let $A \in \binom{\Omega}{k}$ and $B \in \binom{\Omega}{\ell}$. If $n < \infty$ and S is ℓ -homogeneous, then $\text{scc}(\Gamma(S_A)) = \text{wcc}(\Gamma(S_A)) \leq \text{wcc}(\Gamma(S_B)) = \text{scc}(\Gamma(S_B))$.

Question 2.2. Take a finite set Ω and two integers k and ℓ such that $k \leq \ell < k + \ell \leq |\Omega| + 1$. Let S be an ℓ -homogeneous transformation semigroup acting on Ω . For any $A \in \binom{\Omega}{k}$ and $B \in \binom{\Omega}{\ell}$, does it always hold that $\text{wcc}(\Gamma(\tilde{S}_A)) \leq \text{wcc}(\Gamma(\tilde{S}_B))$?

When restricting to permutation groups, the results in Theorem 2.1 are all known more than 40 years ago: Claim (1) for an infinite set Ω was discovered by Brown [12, Corollary 1]; Claim (2) for a finite set Ω was derived by Livingstone and Wagner [38, Theorem 1]; Claim (3), as well as a positive answer to Question 2.2 for permutation groups, was proved by Cameron [15, Proposition 2.3] under the mild restriction of $k + \ell \leq |\Omega|$. Let G be a group acting on a finite set Ω . By Theorem 2.1(2), or more precisely Livingstone-Wagner Theorem [38, Theorem 1], we know that the strong/weak shape of 2^Ω under the action of \overline{G} is a symmetric unimodal distribution. This means that, for any two integers k and ℓ such that $k \leq \ell < k + \ell \leq |\Omega|$, the number of \overline{G} -orbits on $\binom{\Omega}{\ell}$ is equal to the sum of a nonnegative integer c plus the number of \overline{G} -orbits on $\binom{\Omega}{k}$. As an improvement of this fact, Siemons [56, Corollary 4.3] found a natural linear space whose dimension equals this integer c and he [56, Theorem 4.2] even obtained an algorithm to reconstruct the \overline{G} -orbits on $\binom{\Omega}{k}$ from the information on the \overline{G} -orbits on $\binom{\Omega}{\ell}$ without reference to the group G .

Question 2.3. Let Ω be a finite set, and let k and ℓ be two integers such that $k \leq \ell < k + \ell \leq |\Omega|$. Let S be a transformation semigroup on Ω and let Γ be the phase space of \overline{S} .

- (1) Is there a counterpart of [56, Corollary 4.3] which explains the nonnegativity constraint on the integer $\text{wcc}(\Gamma[\binom{\Omega}{\ell}]) - \text{wcc}(\Gamma[\binom{\Omega}{k}])$?
- (2) If S is $(\ell + 1)$ -homogeneous, is there a counterpart of [56, Corollary 4.3] which explains the nonnegativeness of the integer $\text{scc}(\Gamma(S_B)) - \text{scc}(\Gamma(S_A))$ for any $A \in \binom{\Omega}{k}$ and $B \in \binom{\Omega}{\ell+1}$?
- (3) Is there any algorithm to determine the weakly connected components of $\Gamma[\binom{\Omega}{k}]$ from the weakly connected components of $\Gamma[\binom{\Omega}{\ell}]$ without reference to the transformation semigroup S ?

Example 2.4. Let Ω be a set carrying a linear order \prec . A map $g \in \Omega^\Omega$ is order-preserving with respect to \prec provided αg is not bigger than βg in \prec whenever α is not bigger than β in \prec . Let S be the monoid consisting of all order-preserving maps on Ω with respect to the given linear order \prec . It is easy to see that S is ℓ -homogeneous for all $\ell \leq |\Omega|$ but it is even not 2-transitive; by contrast, this phenomenon never happens for permutation groups due to a result of Livingstone and Wagner [38, Theorem 2(b)]. Note that the only permutation contained in S is the identity map in case that Ω is a finite set. This suggests that you may not be able to read Theorem 2.1 or answer Question 2.3 directly from those known facts on permutation groups.

Example 2.5. Let $\Omega = \{1, \dots, 6\}$. Let r and b be two maps in $\text{T}(\Omega)$ such that

$$\begin{aligned} r(1) = r(2) = 3, \quad r(3) = r(4) = 5, \quad r(5) = r(6) = 1; \\ b(6) = b(1) = 2, \quad b(2) = b(3) = 4, \quad b(4) = b(5) = 6. \end{aligned}$$

Let $S = \langle r, b \rangle$. On the left of Figure 1, we depict the phase space $\Gamma(S, \{r, b\})$; on the right of Figure 1, we display both the strong shape and the weak shape of 2^Ω under the action of \overline{S} . Both weak shape and strong shape are unimodal and top-heavy. But neither of them is log-concave. Note that the peak of the weak shape does not happen at the middle rank 3.




Figure 1: $\Gamma(S, \{r, b\})$ and $\Gamma(\bar{S}, \{\bar{r}, \bar{b}\})[\binom{\Omega}{k}]$, $k \in \{0, 1, \dots, 6\}$. See Example 2.5.

Example 2.6. Let Ω be a set of size $n \geq 3$ and let S be a transformation semigroup acting on Ω . If $\text{SS}(\bar{S}, 2^\Omega)$ is not a sequence of all ones and has at least two ones at the beginning of it, then it cannot be log-concave. This happens when S is the alternating group of order $n \geq 4$ and when S is 2-homogeneous but not 3-homogeneous.

Example 2.7. Let n and k be two integers such that $1 \leq k \leq n$. Let Ω be a set of size n and take $X \in \binom{\Omega}{k}$. Let S be the set $\{f \in \text{T}(\Omega) : f|_X = \text{Id}|_X, \Omega \bar{f} = X\}$. Note that S is a transformation semigroup on 2^Ω satisfying

$$\text{wcc}(\Gamma(\bar{S})[\binom{\Omega}{i}]) = \begin{cases} 1, & \text{if } 0 \leq i \leq k; \\ \binom{n}{i}, & \text{if } k+1 \leq i \leq n. \end{cases}$$

This shows that the sequence $\text{WS}(\bar{S}, 2^\Omega)$ is unimodal and top-heavy and that it is not log-concave when $n \geq 2$. Note that $\text{SS}(\bar{S}, 2^\Omega)$ is a sequence of all ones.

Question 2.8. Let S be a transformation semigroup acting on an n -element set Ω . When can we conclude that the strong/weak shape of 2^Ω under the action of \bar{S} is unimodal?

Neumann [44] asked whether every λ -homogeneous permutation group is θ -homogeneous for all cardinals $\lambda > \theta \geq \aleph_0$. Assuming Martin's Axiom, Shelah and Thomas [55] gave a negative answer to it. Hajnal [26] supplied an example to show that 2^θ -homogeneity does not imply θ -homogeneity. An observation in the same vein by Penttila and Siciliano [47, Remark 4.6] was based upon the Generalized Continuum Hypothesis. For each statement in Theorem 2.1(3), Question 2.2 and 2.3, it is interesting to see whether or not it holds in the case that Ω is an infinite set. We are also wondering if the rich theory on oligomorphic permutation groups [16] should have a counterpart for transformation semigroups.

2.2 Partition lattice

Let Ω be a set. For any map $s \in \Omega^\Omega$, we define its *kernel map*, denoted by s^{-1} , to be the map from 2^Ω to 2^Ω that sends $X \in 2^\Omega$ to $Xs^{-1} = \{y \in \Omega : ys \in X\} \in 2^\Omega$. To illustrate the definition, we depict the phase space of a map s on the left of Figure 2 and part of the phase space of s^{-1} in the middle of Figure 2. A *partition* of Ω is a set of nonempty disjoint subsets of Ω whose union is Ω . We call these elements of a partition its *blocks*. The *rank* of a partition π is $\sum_{B \in \pi} (|B| - 1)$. Write $P(\Omega)$ for the set of all partitions of Ω of finite ranks. When $|\Omega| < \infty$, the set $P(\Omega)$ together with the refinement relation forms a geometric lattice, which we call the *partition lattice* of Ω . Note that the rank of a partition in this geometric lattice is $|\Omega|$ minus the number of its blocks. Let $P_k(\Omega)$ be the set of rank- k partitions of Ω , namely, those partitions of Ω of size $|\Omega| - k$. Each transformation $s \in \Omega^\Omega$ induces a transformation s^* of 2^Ω such that $\Pi s^* = \{\pi s^{-1} : \pi \in \Pi\} \setminus \{\emptyset\}$ for all $\Pi \in P(\Omega)$. We demonstrate part of the phase space of s^* on the right of Figure 2 for the map s as shown on the left there. Let S be a transformation semigroup on Ω . We have a derived transformation semigroup $S^* := \{s^* : s \in S\}$ on $P(\Omega)$, which we call the *kernel space* of S . We say that S is *k-kernel homogeneous* if for all $\Pi, \Pi' \in P_k(\Omega)$ there exists $s \in S$ such that $\Pi s^* = \Pi'$, which surely implies $\text{scc}(\Gamma(S^*)[P_k(\Omega)]) \leq 1$.




Figure 2: A map, its inverse and the derived action on partitions.

Example 2.9. On the left of Figure 3, we depict the so-called Černý automaton $\mathcal{C}_4 = \Gamma(S, G)$, where $G = \{a, b\}$ consists of two transformations on a four-element set Ω . On the right of Figure 3, we depict the automaton $\Gamma(S^*, G^*)$ where S^* is acting on $P(\Omega)$. Observe that $\text{WS}(S^*, P(\Omega)) = (1, 1, 1, 1)$ and $\text{SS}(S^*, P(\Omega)) = (1, 2, 2, 1)$ are both unimodal and top-heavy.

For any finite set Ω , $A \in 2^\Omega$, $\pi \in P(\Omega)$ and $s \in \Omega^\Omega$, it holds

$$r(A) \geq r(A\bar{s}) \quad \text{and} \quad r(\pi) \leq r(\pi s^*).$$

This difference between Boolean lattice and partition lattice somehow hints at our difficulty of turning the following conjecture into a result like Theorem 2.1.

Conjecture 2.10. Let Ω be a finite set and let S be a semigroup acting on Ω . Then both $\text{WS}(S^*, P(\Omega))$ and $\text{SS}(S^*, P(\Omega))$ are top-heavy.

For each set Ω and each positive integer $k \leq |\Omega|$, we use $P(\Omega, k)$ for the set of partitions of Ω into k blocks.




Figure 3: Černý automaton \mathcal{C}_4 and its kernel space. See Example 2.9.

- Question 2.11.** (1) Take two positive integers k and ℓ with $k < \ell$. Let Ω be an infinite set and let S be a semigroup S acting on Ω . If S^* is transitive on $P(\Omega, \ell)$, is it true that S^* is transitive on $P(\Omega, k)$?
- (2) The shapes of all Dowling lattices, which include all partition lattices, are real-rooted [8]. What about the top-heavy property of the (strong/weak) shapes of Dowling lattices under a semigroup action?

There has been an active study of those permutation groups which are transitive on the set of all ordered or unordered partitions of a set of a given shape [2, 21, 39, 43]. But even when confining our attention to permutation groups, we are not aware of any work related to Conjecture 2.10 and Question 2.11.

2.3 Subspace lattice

Let Ω be a possibly infinite set of size n , let k be a nonnegative integer with $k \leq n$ and let F be a finite field. We mention that $\text{Gr}(k, F^\Omega)$ is a q -analogue of $\binom{\Omega}{k}$ and their relationship is like the one between Johnson graphs and Grassmann graphs [45]. For each prime power q , we write \mathbb{F}_q for the q -element finite field. Write $\text{Mat}_n(\mathbb{F}_q)$ for the multiplicative semigroup of all Ω by Ω matrices over \mathbb{F}_q each row/column of which have finitely many nonzeros; and write $\text{Aff}_n(\mathbb{F}_q)$ for the semigroup of all affine linear transformation on \mathbb{F}_q^n equipped with the associated product of composition. We regard the empty set as the dimension- (-1) affine/linear subspace. The set of all nonempty finite linear subspaces of \mathbb{F}_q^n is denoted by $\mathcal{P}_{q,n} \doteq \text{Gr}(\mathbb{F}_q^n)$ and the set of all dimension- k linear subspaces of \mathbb{F}_q^n is denoted by $\mathcal{P}_{q,n}^k \doteq \text{Gr}(k, \mathbb{F}_q^n)$. By a finite affine subspace of a linear space V , we mean a translate of a finite linear subspace of V . The set of all finite affine subspaces of \mathbb{F}_q^n is denoted by $\mathcal{A}_{q,n}$ and the set of all dimension- k affine subspaces of \mathbb{F}_q^n is denoted by $\mathcal{A}_{q,n}^{k+1}$. Note that $\mathcal{P}_{q,n}$ and $\mathcal{A}_{q,n}$ are known as *projective geometry* and *affine geometry* over the field \mathbb{F}_q , respectively. For each nonnegative integer k , we put the rank of each element in $\mathcal{P}_{q,n}^k$ and the rank of each element in $\mathcal{A}_{q,n}^k$ to be k , thus getting two valuated posets $\mathcal{P}_{q,n}$ and $\mathcal{A}_{q,n}$,

which are geometric lattices when $n < \infty$.

We are ready to display Theorem 2.12, a q -analogue of Theorem 2.1. Kantor [31, Theorem 1] deduced a q -analogue of the aforementioned result of Livingstone and Wagner [38, Theorem 1]. If the semigroup $S \leq \text{Mat}_n(\mathbb{F}_q)$ is a subgroup of the general linear group $\text{GL}_n(\mathbb{F}_q)$, Stanley [59, Corollary 9.9] found that $\text{SS}(S^{\mathcal{P}}, \mathcal{P}_{q,n})$ and $\text{WS}(S^{\mathcal{P}}, \mathcal{P}_{q,n})$ are both symmetric and unimodal for finite n . Penttila and Siciliano [47, Theorem 4.4(ii), (iii)] generalized this result of Stanley for groups to the case that n is infinite.

Theorem 2.12. *Let n be the size of a nonempty set, and let q be a prime power.*

- (1) *Let $S \leq \text{Mat}_n(\mathbb{F}_q)$ be a linear transformation semigroup acting on \mathbb{F}_q^n . For each $g \in S$, write $g^{\mathcal{P}}$ for $\bar{g}|_{\mathcal{P}_{q,n}}$. Let $S^{\mathcal{P}}$ be the transformation semigroup $\{g^{\mathcal{P}} : g \in S\}$ acting on $\mathcal{P}_{q,n}$. Then $\text{SS}(S^{\mathcal{P}}, \mathcal{P}_{q,n})$ and $\text{WS}(S^{\mathcal{P}}, \mathcal{P}_{q,n})$ are both top-heavy.*
- (2) *Let $T \leq \text{Aff}_{n-1}(\mathbb{F}_q)$ be an affine linear transformation semigroup acting on \mathbb{F}_q^{n-1} . For each $g \in T$, write $g^{\mathcal{A}}$ for $\bar{g}|_{\mathcal{A}_{q,n-1}}$. Let $T^{\mathcal{A}}$ be the transformation semigroup $\{g^{\mathcal{A}} : g \in T\}$ acting on $\mathcal{A}_{q,n-1}$. Then $\text{SS}(T^{\mathcal{A}}, \mathcal{A}_{q,n-1})$ and $\text{WS}(T^{\mathcal{A}}, \mathcal{A}_{q,n-1})$ are both top-heavy.*

Remark 2.13. When n is infinite, Theorems 2.1 and 2.12 in the original version of this paper, submitted on 19 July 2018, contains weaker results. Following the proof presented by Bercov and Hobby for [9, Corollary 1] and also the proof of Roy for [51, Theorem], we used the existence of Ramsey number [49, Theorem A] to derive Theorem 2.1(1) for infinite n . A similar argument based on Ramsey number shows that both $\text{SS}(S^{\mathcal{P}}, \mathcal{P}_{q,n})$ and $\text{SS}(T^{\mathcal{A}}, \mathcal{A}_{q,n-1})$ are 1-top-heavy for infinite n in the setting of Theorem 2.12. After the acceptance of this paper in 2022, we notice the work of Penttila and Siciliano [47, Lemma 3.1], which was submitted on 30 April 2019 and published in 2021, and thus arrive at the corresponding strengthening in Theorem 2.1(2) and Theorem 2.12 via an application of their idea. See Lemma 3.6.

Remark 2.14. Kantor [32, Theorem 2] determined all the ordered-basis-transitive finite geometric lattices of rank at least three: Roughly speaking, they are Boolean lattices, projective (affine) geometries, and four sporadic designs. Kantor's classification theorem along with Theorems 2.1 and 2.12 may be a basis for getting homogeneity results about ordered-basis-transitive matroids.

Question 2.15. A general projective geometry is defined to be a modular combinatorial geometry that is connected in the sense that the point set cannot be expressed as the union of two proper flats [64, page 313]. Can we establish a counterpart of Theorem 2.12 for general projective geometries?

In mathematics we encounter quite some nice duality phenomena, say Chow's Theorem [45, Corollary 3.1] and many duality concepts for matroids [13]. For projectie geometry, we have the following duality result of Stanley [59, Corollary 9.9].

Theorem 2.16 (Stanley). *Let F be a finite field and let k and n be two positive integers with $k < n$. For any subgroup G of $\text{GL}(n, F)$, the number of orbits of the action of G on $\text{Gr}(k, F^n)$ must be the same with the number of orbits of G acting on $\text{Gr}(n - k, F^n)$.*

Question 2.17. If n is the size of an infinite set, does Theorem 2.16 still hold? Here, we should first of all choose a good definition for infinite Grassmannians [46].

2.4 A glimpse of matroid

In previous subsections, we discuss those poset endomorphisms which are derived from either set transformations or linear transformations. Since finite geometric lattices just encode information of finite matroids, it is natural to ask why not directly consider matroids and morphisms among matroids, namely those transformations which preserve “independence structure”.

Let M_1 and M_2 be two matroids and let f be a map from \mathcal{E}_{M_1} to \mathcal{E}_{M_2} . We call f a *weak map from M_1 to M_2* provided

$$r_{M_1}(A) \geq r_{M_2}(A\bar{f})$$

holds for all $A \subseteq \mathcal{E}_{M_1}$, and we call f a *strong map from M_1 to M_2* provided the preimage of any flat in M_2 is a flat of M_1 [33, 35, 57]. It is known that all strong maps must be weak maps.

Let M be a matroid on the ground set $\mathcal{E}_M = \Omega$. Let $T_M(\Omega)$ ($T_M^*(\Omega)$) be the monoid consisting of all elements of $T(\Omega)$ which are weak (strong) maps from M to itself. If we know that S is a subsemigroup of $T_M(\Omega)$ ($T_M^*(\Omega)$) acting on Ω , we can define a digraph $\Gamma_{M,t}(S)$ on $F_t(M)$ as follows: for any $X, Y \in F_t(M)$, there is an arc from X to Y if and only if there is $g \in S$ such that the minimum flat containing $X\bar{g}$ in M is Y . What is the relationship between the connectivity of $\Gamma_{M,t}(S)$ and $\Gamma_{M,r}(S)$ for different t and r ? We can ask the same question by imposing the extra condition that every element $f \in S$ is a bijection on Ω . If the matroid is a very special uniform matroid, namely a matroid in which all sets are independent, one can see that what is discussed in Section 1.3 becomes a very special case of this general setting.

Vámos matroid, also known as Vámos cube, is a famous non-algebraic matroid [5, 22, 42, 54]; see [24, Example 6.30] for a description of this rank-4 matroid over a ground set of size eight.


Example 2.18. Let M be the Vámos matroid and let S be a subsemigroup of $T_M^*(\mathcal{E}_M)$. It holds $\text{wcc}(\Gamma_{M,1}(S)) \leq \text{wcc}(\Gamma_{M,2}(S)) \leq \text{wcc}(\Gamma_{M,3}(S))$ and $\text{scc}(\Gamma_{M,1}(S)) \leq \text{scc}(\Gamma_{M,2}(S)) \leq \text{scc}(\Gamma_{M,3}(S))$.

Remark 2.19. Compared with the Fundamental Theorem of Projective (Affine) Geometry [17, 48], we think that weak/strong maps and bijective weak/strong maps for matroids are natural extensions of linear transformations and invertible linear transformations for linear spaces. We also mention the well-adopted viewpoint that the full permutation group and the full transformation semigroup can be interpreted as the general linear group and the linear transformation semigroup over the field with one element.

3 Valuated poset and incidence operator

3.1 Hereditary endomorphism and injective incidence operator

To prepare for a proof of our main results listed in Section 2, we will introduce a key property and then present a key lemma for our work. The key property is the so-called hereditary endomorphisms. The key lemma is Lemma 3.2, which gives us some information of the strong/weak shapes of a poset under some semigroup action, provided the semigroup consists of hereditary endomorphisms and that some linear map associated with the poset is injective.

Figure 4: An (ℓ, k) -hereditary endomorphism.

Let P be a valuated poset. For any nonnegative integers $k \leq \ell$, we call the poset P (k, ℓ) -finite provided $P_k \neq \emptyset$, $P_\ell \neq \emptyset$ and the set $P_\ell \cap P_\uparrow(\alpha)$ is finite for every $\alpha \in P_k$; we call P (ℓ, k) -finite provided $P_k \neq \emptyset$, $P_\ell \neq \emptyset$ and the set $P_\downarrow(\beta) \cap P_k$ is finite for every $\beta \in P_\ell$; we call $g \in \text{End}(P)$ a (k, ℓ) -hereditary endomorphism if for all $\alpha \in P_k$ which satisfies $r_P(g(\alpha)) = r_P(\alpha) = k$ it happens that g induces a bijection from the set $P_\ell \cap P_\uparrow(\alpha)$ to $P_\ell \cap P_\uparrow(\alpha g)$; we call $g \in \text{End}(P)$ an (ℓ, k) -hereditary endomorphism if for each $\beta \in P_\ell$, $r_P(\beta g) = r_P(\beta) = \ell$ ensures that g induces a bijection from the set $P_k \cap P_\downarrow(\beta)$ to $P_k \cap P_\downarrow(\beta g)$. See Figure 4 for an illustration. For any $k, \ell \in \mathbb{Z}_{\geq 0}$, we designate by $\text{hEnd}_{k, \ell}(P)$ the set of all (k, ℓ) -hereditary endomorphisms of the valuated poset P .

Let S be a transformation semigroup on a valuated poset P and let G be a generating set of S . For any two nonnegative integers k and ℓ with $k \leq \ell \leq r(P)$, we set $\Pi_{S, G}(k, \ell)$ to be the digraph with vertex set P_k and arc set

$$\{(\alpha, \alpha') \in P_k \times P_k : \exists g \in G, \beta \in P_\ell \text{ s.t. } \beta g \in P_\ell, \alpha' = \alpha g, \alpha \in P_\downarrow(\beta)\};$$

we set $\Pi_{S, G}(\ell, k)$ to be the digraph with vertex set P_ℓ and arc set

$$\{(\alpha, \alpha') \in P_\ell \times P_\ell : \exists g \in G, \beta \in P_k \text{ s.t. } \beta g \in P_k, \alpha' = \alpha g, \alpha \in P_\uparrow(\beta)\}.$$

We use the shorthand $\Pi_S(k, \ell)$ for $\Pi_{S, S}(k, \ell)$.

Lemma 3.1. *Let P be a valuated poset. Take two nonnegative integers k and ℓ such that $k, \ell \leq r(P)$ and that P is (ℓ, k) -finite. Let S be a sub-semigroup of $\text{hEnd}_{\ell, k}(P)$, let G be a generator set of S , and let $\Gamma \doteq \Gamma(S, G)$. Let $\beta \in P_\ell$ and let $\alpha \in P_k$ be an element comparable with β . Assume that g and h are two elements of S such that $\beta g \in P_\ell$ and $\beta gh = \beta$. Then there exists $f \in S$ such that $\beta gf \in P_\ell$ and $\alpha gf = \alpha$. Especially, if every weakly connected component of $\Gamma[P_\ell]$ is strongly connected, then so is $\Pi_{S, G}(k, \ell)$.*

Proof. The second claim is immediate from the first and so our task is just to prove the first one. Without loss of generality, we assume that $k < \ell$. Since $\beta(gh) = \beta$ and $gh \in S \leq \text{hEnd}_{\ell, k}(P)$, it follows that gh induces a permutation on $P_k \cap P_\downarrow(\beta)$. But from the assumption that P is (ℓ, k) -finite, we see that $P_k \cap P_\downarrow(\beta)$ is a finite set, which contains α . This means that there exists a positive integer r such that $\alpha(gh)^r = \alpha$. Accordingly, for $f = (hg)^{r-1}h \in S$ it holds $(\beta g)f = (\beta g)(hg)^{r-1}h = \beta(gh)^r = \beta \in P_\ell$ and $(\alpha g)f = (\alpha g)(hg)^{r-1}h = \alpha(gh)^r = \alpha$, finishing the proof. \square

For any set Ω , \mathbb{Q}^Ω refers to the linear space of all rational functions on Ω . If P is an (ℓ, k) -finite valuated poset, the incidence operator $\zeta_P^{k, \ell}: \mathbb{Q}^{P_k} \rightarrow \mathbb{Q}^{P_\ell}$ is the linear operator

such that for all $f \in \mathbb{Q}^{P_k}$ and $\beta \in P_\ell$, we have

$$(\zeta_P^{k,\ell}(f))(\beta) = \begin{cases} \sum_{\alpha \in P_k \cap P_\downarrow(\beta)} f(\alpha), & \text{if } k \leq \ell; \\ \sum_{\alpha \in P_k \cap P_\uparrow(\beta)} f(\alpha), & \text{if } k > \ell. \end{cases} \quad (3.1)$$

Lemma 3.2. *Let P be a valuated poset. Take two nonnegative integers k and ℓ not exceeding $\mathbf{r}(P)$ such that P is (ℓ, k) -finite, and hence $\zeta_P^{k,\ell}$ is well-defined. Let S be a subsemigroup of $\text{hEnd}_{\ell,k}(P)$ and let Γ stand for $\Gamma(S)$. Assume that $\zeta_P^{k,\ell}$ is an injective linear map from \mathbb{Q}^{P_k} to \mathbb{Q}^{P_ℓ} .*

$$(1) \quad \text{wcc}(\Gamma[P_k]) \leq \text{wcc}(\Pi_S(k, \ell)) \leq \text{wcc}(\Gamma[P_\ell]).$$

$$(2) \quad \text{scc}(\Gamma[P_k]) \leq \text{scc}(\Pi_S(k, \ell)) \leq \text{scc}(\Gamma[P_\ell]).$$

Proof. (1) The first inequality is a consequence of the fact that $E(\Pi_S(k, \ell)) \subseteq E(\Gamma[P_k])$.

Let $W \subseteq \mathbb{Q}^{P_\ell}$ be the subspace of all functions which are constant on each weakly connected component of $\Gamma[P_\ell]$; let $V \subseteq \mathbb{Q}^{P_k}$ be the subspace of all functions which are constant on each weakly connected component of $\Pi_S(k, \ell)$. Note that $\dim(V) = \text{wcc}(\Pi_S(k, \ell))$ and $\dim(W) = \text{wcc}(\Gamma[P_\ell])$ and so it suffices to demonstrate $\dim(V) \leq \dim(W)$.

By symmetry, we only deal with the case of $k \leq \ell$. For every $f \in V$ and every arc $(\beta, \beta g)$ of $\Gamma[P_\ell]$, we have

$$\begin{aligned} (\zeta_P^{k,\ell}(f))(\beta g) &= \sum_{\alpha' \in P_k \cap P_\downarrow(\beta g)} f(\alpha') \\ &= \sum_{\alpha \in P_k \cap P_\downarrow(\beta)} f(\alpha g) && (g \in \text{hEnd}_{\ell,k}(P)) \\ &= \sum_{\alpha \in P_k \cap P_\downarrow(\beta)} f(\alpha) && (f \in V) \\ &= (\zeta_P^{k,\ell}(f))(\beta). \end{aligned}$$

This says that $\zeta_P^{k,\ell}(f) \in W$ for all $f \in V$. Hence, by the injectivity of $\zeta_P^{k,\ell}$, $\dim(V) \leq \dim(W)$, as wanted.

(2) The first inequality is a consequence of the fact that $E(\Pi_S(k, \ell)) \subseteq E(\Gamma[P_k])$.

Let $W' \subseteq \mathbb{Q}^{P_\ell}$ be the subspace of all functions which are constant on each strongly connected component of $\Gamma[P_\ell]$; let $V' \subseteq \mathbb{Q}^{P_k}$ be the subspace of all functions which are constant on each strongly connected component of $\Pi_S(k, \ell)$. Note that $\dim(V') = \text{scc}(\Pi_S(k, \ell))$ and $\dim(W') = \text{scc}(\Gamma[P_\ell])$ and so it suffices to demonstrate $\dim(V') \leq \dim(W')$. Take $f \in V'$. As $\zeta_P^{k,\ell}$ is injective, we aim to show that $\zeta_P^{k,\ell}(f) \in W'$.

By symmetry, we only deal with the case of $k \leq \ell$. Assume that β and βg are from the same strongly connected component of $\Gamma[P_\ell]$, where $g \in S$. By the first claim of Lemma 3.1, for every $\alpha \in P_k \cap P_\downarrow(\beta)$, α and αg fall into the same strongly connected component of $\Gamma[P_k]$ and so, as $f \in V'$,

$$f(\alpha) = f(\alpha g). \quad (3.2)$$

This allows us to write

$$\begin{aligned}
 (\zeta_P^{k,\ell}(f))(\beta g) &= \sum_{\alpha' \in P_k \cap P_\downarrow(\beta g)} f(\alpha') \\
 &= \sum_{\alpha \in P_k \cap P_\downarrow(\beta)} f(\alpha g) && (g \in \text{hEnd}_{\ell,k}(P)) \\
 &= \sum_{\alpha \in P_k \cap P_\downarrow(\beta)} f(\alpha) && (\text{Equation (3.2)}) \\
 &= (\zeta_P^{k,\ell}(f))(\beta),
 \end{aligned}$$

proving that $\zeta_P^{k,\ell}(V') \subseteq W'$, as desired. \square

3.2 Injectivity

In order to apply Lemma 3.2, we may need to have some results to guarantee the injectivity of an incidence operator. In this regard, a good understanding of the incidence algebra of a poset may be valuable [36, 68]. We mention that Guiduli [4, Theorem 9.4] established an injectivity result for the so-called rank-regular semi-lattices. It may also be quite useful if the following conjecture [34, Conjecture 1.1] can be verified.

Conjecture 3.3 (Kung). *Let P be a finite geometric lattice. Let k and ℓ be two positive integers such that $k \leq \ell \leq \frac{r(P)}{2}$. Then $\ker(\zeta_P^{k,\ell}) = \{0\}$.*

We suggest a slight strengthening of Kung's Conjecture (Conjecture 3.3) as follows.

Conjecture 3.4. *Let P be a geometric lattice. Let k and ℓ be two nonnegative integers such that $k \leq \ell \leq k + \ell \leq r(P)$. If P is (ℓ, k) -finite, then $\zeta_P^{k,\ell}$ is an injective map.*

Remark 3.5. Let M be a matroid of rank r . Let S be a subsemigroup of $T_M^*(\mathcal{E}_M)$. For every $f \in S$, let $f': F(M) \rightarrow F(M)$ be the map sending a flat $X \in F(M)$ to the minimum flat containing $X\bar{f}$ in M . Assume that $f' \in \text{hEnd}_{\ell,k}(F(M))$ for every $f \in S$. In light of Lemma 3.2, if Conjecture 3.4 is valid for the lattice $F(M)$, we will be able to conclude that both the sequence $(\text{wcc}(\Gamma_{M,0}(S)), \dots, \text{wcc}(\Gamma_{M,r}(S)))$ and the sequence $(\text{scc}(\Gamma_{M,0}(S)), \dots, \text{scc}(\Gamma_{M,r}(S)))$ are top-heavy.

Let P be a valuated poset which is (ℓ, k) -finite for all nonnegative integers $k \leq \ell$. We say that P has a *top-heavy injective incidence operator* provided $\zeta_P^{k,\ell}$ is an injective linear map from \mathbb{Q}^{P_k} to \mathbb{Q}^{P_ℓ} for all nonnegative integers k and ℓ satisfying $k \leq \ell \leq k + \ell \leq r(P)$.

Penttila and Siciliano [47, Lemma 3.1] pointed out a simple way to establish some injectivity result for linear operators between infinite-dimensional linear spaces whenever they fulfil certain finiteness characteristics. We reformulate their observation below for the convenience of our later usage.

Lemma 3.6. *Let P be a valuated poset. Let $k \leq \ell$ be two nonnegative integers such that P is (ℓ, k) -finite. Assume that for every $A \in P_k$, we can find a finite subset Y of $P_{k+\ell}$ such that the ideal generated by Y in P , denoted Y^\downarrow and with the restriction of r_P as its rank function, contains A and possesses a top-heavy injective incidence operator. Then $\zeta_P^{k,\ell}$ is an injective linear map from \mathbb{Q}^{P_k} to \mathbb{Q}^{P_ℓ} .*

Proof. Take $f \in \ker \zeta_P^{k,\ell}$. Assume, for sake of contradiction, that $f(A) \neq 0$ for some $A \in P_k$. Choose $Y \subseteq P_{k+\ell}$ such that $A \in Y^\downarrow \cap P_k$ and Y^\downarrow possesses a top-heavy injective incidence operator. Let Q represent the resulting valuated poset on Y^\downarrow . Let g be the restriction of f on Q_k and let h be the restriction of $\zeta_P^{k,\ell}(f) = 0$ on Y . We have $0 = h = \zeta_Q^{k,\ell}(g)$ but $g(A) = f(A) \neq 0$, violating the assumption that Y^\downarrow has a top-heavy injective incidence operator. \square

3.3 Incidence operator as an intertwiner

For $f \in \Psi^\Omega$, we sometimes need to talk about $f(\omega)$ for $\omega \notin \Omega$. Following the practice of those mathematics with natural multivalued operations [7, 14, 65], we create a universal “don’t care” symbol $\star \notin \Psi$ and will set $f(\omega) = \star$. We often regard \star as all possible values in Ψ and so, whenever we have some addition operation $+$ on Ψ , we extend it to $\Psi \cup \{\star\}$ by setting $\star + \psi = \star$ for all $\psi \in \Psi \cup \{\star\}$.

Let P be a valuated poset. Let k and ℓ be two nonnegative integers no greater than $r(P)$. Let $g \in P^P$. For $f \in \mathbb{Q}^{P_k}$, we write $fg^{\dagger,k}$ for the element in $(\{\star\} \cup \mathbb{Q})^{P_k}$, where \star stands for “don’t care” and can be thought of as the whole set \mathbb{Q} , such that the following holds for all $\beta \in P_k$:

$$fg^{\dagger,k}(\beta) = \begin{cases} f(\beta g), & \text{if } \beta g \in P_k; \\ \star, & \text{if } \beta g \notin P_k. \end{cases}$$

Denote by $\text{Fix } g^{\dagger,k}$ the set of $f \in \mathbb{Q}^{P_k}$ for which

$$fg^{\dagger,k}(\beta) \in \{f(\beta), \star\}$$

holds for all $\beta \in P_k$. If $g \in \text{hEnd}_{\ell,k}(P)$, we say that it is a *good (ℓ, k) -hereditary endomorphism of P* provided that for any $\beta \in P_\ell$ with $\beta g \notin P_\ell$ it holds $\alpha g \notin P_k$ for some $\alpha \in P_k$ which is comparable to β in P . Assuming that g is a good (ℓ, k) -hereditary endomorphism of P , for any $\beta \in P_\ell$ and $f \in \mathbb{Q}^{P_k}$ we will have

$$\begin{aligned} (\zeta_P^{k,\ell}(f)g^{\dagger,\ell})(\beta) &= (\zeta_P^{k,\ell}(f))(\beta g) \\ &= \sum_{\alpha' \in P_k \cap (P_\downarrow(\beta g) \cup P_\uparrow(\beta g))} f(\alpha') \\ &= \sum_{\alpha \in P_k \cap (P_\downarrow(\beta) \cup P_\uparrow(\beta))} f(\alpha g) \\ &= (\zeta_P^{k,\ell}(fg^{\dagger,k}))(\beta) \end{aligned}$$

whenever $\beta g \in P_\ell$, and that

$$\begin{aligned} (\zeta_P^{k,\ell}(f)g^{\dagger,\ell})(\beta) &= (\zeta_P^{k,\ell}(f))(\beta g) \\ &= \star \\ &= (\zeta_P^{k,\ell}(fg^{\dagger,k}))(\beta) \end{aligned}$$

whenever $\beta g \notin P_\ell$. This observation can be summarized by the commutative diagram in Figure 5, which implies that $\text{Fix } g^{\dagger,k}$ is mapped by $\zeta_P^{k,\ell}$ to $\text{Fix } g^{\dagger,\ell}$ for all good (ℓ, k) -hereditary endomorphisms g of P .

$$\begin{array}{ccc}
 f & \xrightarrow{\zeta_P^{k,\ell}} & \zeta_P^{k,\ell}(f) \\
 g^{\dagger,k} \downarrow & & \downarrow g^{\dagger,\ell} \\
 fg^{\dagger,k} & \xrightarrow{\zeta_P^{k,\ell}} & \zeta_P^{k,\ell}(fg^{\dagger,k})
 \end{array}$$

Figure 5: The incidence operator intertwines with every good hereditary endomorphism.

- Example 3.7.** (1) Let Ω be a set of size n . Assume that $2 \leq k < \ell \leq n$. Here is an easy observation used often in the study of synchronizing automata: For any $g \in \Omega^\Omega$ and any $A \in \binom{\Omega}{\ell}$, we have $|A\bar{g}| = \ell$ if and only if $|B\bar{g}| = k$ for all $B \in \binom{A}{k}$. This conclusion is surely not valid any more when $k \leq 1$. Note that \bar{g} is a good (ℓ, k) -hereditary endomorphism of the Boolean lattice 2^Ω for each $g \in \Omega^\Omega$.
- (2) Take integers n, k and ℓ such that $2 \leq k < \ell \leq n$ and let q be a prime power. Let $P = \mathcal{P}_{q,n}$ or $P = \mathcal{A}_{q,n-1}$. Similar to the above claim on Boolean lattice, \bar{M} is a good (ℓ, k) -hereditary endomorphism of P for each $M \in \text{Mat}_n(\mathbb{F}_q)$ or $M \in \text{Aff}_{n-1}(\mathbb{F}_q)$, respectively.

4 Boolean semiring

Let Ω be a set and let k and ℓ be two nonnegative integers such that $k < \ell \leq |\Omega|$. For the valuated poset $P = B_\Omega$, we write the incidence operator $\zeta_P^{k,\ell}$ defined in Equation 3.1 as $\zeta_\Omega^{k,\ell}$. That is,

$$(\zeta_\Omega^{k,\ell}(f))(B) = \sum_{A \in \binom{B}{k}} f(A)$$

for all $f \in \mathbb{Q}^{\binom{\Omega}{k}}$ and $B \in \binom{\Omega}{\ell}$.

Following a common approach in establishing homogeneity of permutation groups [15, 41] [20, pages 20–22], we will make use of the ensuing result on the rank of the subset inclusion matrix. The result has been discovered independently by many but the earliest appearance of it dates back to the work of Gottlieb [25, Corollary 2]. Among many different proofs of this classical result, we refer the reader to [18, Corollary] and [56, Theorem 2.4]. Note that it gives a positive answer to Conjecture 3.4 for Boolean lattices.

Lemma 4.1 (Gottlieb). *Let Ω be a nonempty finite set. Then $\ker \zeta_\Omega^{k,\ell} = \{0\}$ for any two integers k and ℓ satisfying $0 \leq k \leq \ell \leq k + \ell \leq |\Omega|$.*

Let Ω be a set and S be a transformation semigroup on Ω . Let $\Omega^\# \doteq \{(\omega, C) : \omega \in C \in 2^\Omega\}$ and, for each $g \in S$, let $g^\#$ be the transformation on $\Omega^\#$ which sends (ω, C) to $(\omega g, C\bar{g})$ for all $(\omega, C) \in \Omega^\#$. Let $S^\#$ stand for the transformation semigroup on $\Omega^\#$ consisting of all elements $g^\#$ for $g \in S$. For all positive integers ℓ , we use the following notation:

$$\Omega_\ell^\# \doteq \{(\omega, C) : \omega \in C \in \binom{\Omega}{\ell}\}$$

and

$$\Gamma_\ell^\sharp(S) \doteq \Gamma(S^\sharp)[\Omega_\ell^\sharp].$$

Here is a result analogous to Lemma 3.1.

Lemma 4.2. *Let m be a positive integer and let S be an m -homogeneous transformation semigroup acting on a set Ω . Then the digraph $\Gamma_m^\sharp(S)$ is symmetric. Especially, $\text{wcc}(\Gamma_m^\sharp(S)) = \text{scc}(\Gamma_m^\sharp(S))$.*

Proof. Take $(\omega, C) \in \Omega_m^\sharp$ and $g \in S$ such that $|C\bar{g}| = m$. Our task is to show the existence of $h \in S$ such that $(\omega g, C\bar{g})h^\sharp = (\omega, C)$. As S is m -homogeneous, we can find $f \in S$ such that $C\bar{g}f = (C\bar{g})\bar{f} = C$. Hence, the fact that $|C| = m < \infty$ allows us to obtain a positive integer r for which $(gf)^r|_C$ is the identity map on C . This means that we can choose h to be $f(gf)^{r-1}$. \square

Lemma 4.3. *Let Ω be a set, let m be an integer satisfying $|\Omega| \geq m > 1$, and let S be a transformation semigroup on Ω . For every $X \in \binom{\Omega}{m}$, it holds*

$$\text{scc}(\Gamma(S_X)) = \text{wcc}(\Gamma(S_X)) \leq \text{wcc}(\Gamma_m^\sharp(S)) \leq \text{scc}(\Gamma_m^\sharp(S)). \quad (4.1)$$

Moreover, if S is m -homogeneous, then

$$\text{scc}(\Gamma(S_X)) = \text{wcc}(\Gamma(S_X)) = \text{wcc}(\Gamma_m^\sharp(S)) = \text{scc}(\Gamma_m^\sharp(S)). \quad (4.2)$$

Proof. It is trivial to see that $\text{wcc}(\Gamma(S_X)) = \text{scc}(\Gamma(S_X))$ and $\text{wcc}(\Gamma_m^\sharp(S)) \leq \text{scc}(\Gamma_m^\sharp(S))$. Let us call each strongly/weakly connected component of $\Gamma(S_X)$ a component. To prove Equation (4.1), let us find an injective map ψ from the set of components of $\Gamma(S_X)$ to the set of weakly connected components of $\Gamma_m^\sharp(S)$.

For each $\gamma \in X$, let the weakly connected component of $\Gamma_m^\sharp(S)$ containing (γ, X) be $\psi'(\gamma)$. Take γ_1, γ_2 from the same component of $\Gamma(S_X)$. We may assume that $\gamma_1 g = \gamma_2$ and $X\bar{g} = X$ for some $g \in S$. As $(\gamma_1, X)g^\sharp = (\gamma_1 g, X\bar{g}) = (\gamma_2, X)$, we see that $\psi'(\gamma_1) = \psi'(\gamma_2)$. For each component C of $\Gamma(S_X)$, we can now choose any $\gamma \in C$ and get a well-defined map ψ by setting $\psi(C) = \psi'(\gamma)$. For every weakly connected component C^\sharp of $\Gamma_m^\sharp(S)$, let $\phi(C^\sharp)$ be the set $\{\gamma \in X : (\gamma, X) \in C^\sharp\}$. It is routine to check that $\phi\psi(C) = C$ for every component C of $\Gamma(S_X)$, proving that ψ is injective, as desired.

Assume now S is m -homogeneous. It follows from Lemma 4.2 that $\text{wcc}(\Gamma_m^\sharp(S)) = \text{scc}(\Gamma_m^\sharp(S))$. We thus call each strongly/weakly connected component of $\Gamma_m^\sharp(S)$ simply a component. Since S is m -homogeneous, for every component C^\sharp of $\Gamma_m^\sharp(S)$, we have $\phi(C^\sharp) \neq \emptyset$. This verifies that ϕ and ψ are inverses of each other. We thus get Equation (4.2) and so finish the proof. \square

Proof of Theorem 2.1. (1) This is a special case of (2).

(2) This is direct from Lemmas 3.2, 3.6 and 4.1.

(3) Since S is ℓ -homogeneous, it follows from Lemma 4.3 that

$$\text{wcc}(\Gamma(S_A)) = \text{scc}(\Gamma(S_A)) \leq \text{wcc}(\Gamma_k^\sharp(S))$$

and

$$\text{wcc}(\Gamma(S_B)) = \text{scc}(\Gamma(S_B)) = \text{wcc}(\Gamma_\ell^\#(S)).$$

It then remains to prove $\text{wcc}(\Gamma_\ell^\#(S)) \geq \text{wcc}(\Gamma_k^\#(S))$.

We regard $\Omega^\#$ as a valued poset by putting $r((\alpha, X)) = |X|$ and requiring $(\alpha, X) < (\beta, Y)$ if and only if $\alpha = \beta \in \Omega$ and $X \subsetneq Y \subseteq \Omega$. Note that $S^\# \subseteq \text{hEnd}_{\ell, k}(\Omega^\#)$. In view of Lemma 3.2(1), it is sufficient to show that $\zeta_{\Omega^\#}^{k, \ell}$ is injective.

For each nonnegative integer m and each $\alpha \in \Omega$, let $\Omega_{m, \alpha}^\# \doteq \{(\alpha, A) : (\alpha, A) \in \Omega_m^\#\}$. Corresponding to the partition $\Omega_k^\# = \bigcup_{\alpha \in \Omega} \Omega_{k, \alpha}^\#$ and $\Omega_\ell^\# = \bigcup_{\beta \in \Omega} \Omega_{\ell, \beta}^\#$, the $\Omega_k^\# \times \Omega_\ell^\#$ matrix $\zeta_{\Omega^\#}^{k, \ell}$ is viewed as a partitioned matrix with blocks $\zeta_{\alpha, \beta}$, which are the submatrices with row index set $\Omega_{k, \alpha}^\#$ and column index set $\Omega_{\ell, \beta}^\#$, where $\alpha, \beta \in \Omega$. Observe that

$$\zeta_{\alpha, \beta} = \begin{cases} \zeta_{\Omega \setminus \{\alpha\}}^{k-1, \ell-1}, & \text{if } \alpha = \beta; \\ 0, & \text{otherwise.} \end{cases}$$

Since $(k-1) + (\ell-1) \leq |\Omega| - 1$, it follows from Lemma 4.1 that $\zeta_{\alpha, \alpha} = \zeta_{\Omega \setminus \{\alpha\}}^{k-1, \ell-1}$ is of full row rank for all $\alpha \in \Omega$. This implies that $\zeta_{\Omega^\#}^{k, \ell}$ is an injective linear map, as desired. \square

Remark 4.4. Let Ω be a set, which is not necessarily finite. Let k and ℓ be two integers with $k \leq \ell \leq k + \ell \leq |\Omega|$. For all $f \in \mathbb{Q}^{\binom{\Omega}{\ell}}$ and $A \in \binom{\Omega}{k}$, we put

$$(\zeta_\Omega^{\ell, k}(f))(A) = \sum_{A \subseteq B} f(B).$$

Making use of Lemma 4.1, it is easy to see that the linear transformation $\zeta_\Omega^{\ell, k} : \mathbb{Q}_{\text{fin}}^{\binom{\Omega}{\ell}} \rightarrow \mathbb{Q}_{\text{fin}}^{\binom{\Omega}{k}}$ is always a surjective map. Unfortunately, we do not see if this observation is helpful for getting a possible counterpart of Theorem 2.1(3) for an infinite set Ω .

5 A graded Möbius algebra

Möbius algebra is a semigroup algebra which plays an important role in combinatorics [36, Section 3.6]. Huh and Wang [28] introduced a graded Möbius algebra for geometric lattices. Let L be a finite geometric lattice with rank function (valuation) r . Define a \mathbb{Q} -algebra $M(L, \mathbb{Q})$, called the *graded Möbius algebra* of L [28], to be the linear space with L as a \mathbb{Q} -basis together with a multiplication given by

$$xy = \begin{cases} x \vee y, & \text{if } r(x) + r(y) = r(x \vee y) \\ 0, & \text{if } r(x) + r(y) > r(x \vee y), \end{cases}$$

and extended by linearity and distributivity. For any nonnegative integers $k \leq \ell$, it is easy to see that the linear map $\xi_L^{k, \ell}$ as specified below is well-defined:

$$\begin{aligned} \xi_L^{k, \ell} : \quad \mathbb{Q}^{L_k} &\rightarrow \mathbb{Q}^{L_\ell} \\ \phi &\mapsto \left(\sum_{x \in L_1} x \right)^{\ell-k} \phi. \end{aligned}$$

We call a finite geometric lattice a *realizable lattice* if it is the matroid lattice of a finite realizable matroid. Here is the main result of Huh and Wang [28, Theorem 6] in their work

on solving the realizable case of the top-heavy conjecture of Dowling-Wilson. Huh and Wang [28, Conjecture 7] conjectured that Theorem 5.1 holds without the assumption of realizability.

Theorem 5.1 (Huh and Wang). *Let L be a finite realizable geometric lattice with rank r . For any integers k and ℓ such that $k \leq \ell \leq k + \ell \leq r$, the linear map $\xi_L^{k,\ell}$ is injective.*

Remark 5.2. (1) The partition lattice $P(\Omega)$ is isomorphic with the flat lattice of the graphic matroid of the complete graph on Ω . Note that a graphic matroid is regular, namely it is representable over every field. This means that finite partition lattices are realizable.

(2) Assume that L is either a Boolean lattice, or a subspace lattice or a partition lattice. It is easy to see that $\xi_L^{k,\ell} = C_{L,k,\ell} \zeta_L^{k,\ell}$ for some positive integer $C_{L,k,\ell}$ which is determined by L, k and ℓ . Especially, $\xi_L^{k,k+1} = \zeta_L^{k,k+1}$. An important message here is that, $\zeta_L^{k,\ell}$ and $\xi_L^{k,\ell}$, as two \mathbb{Q} -linear maps, are either both injective or both non-injective.

Kung [34, Theorem 1.3] verified Conjecture 3.3 for partition lattices of finite sets. We can improve his result a little bit now. When Ω is finite, Lemma 5.3 claims that Conjecture 3.4 holds for partition lattices.

Lemma 5.3. *Let Ω be a set. Let k and ℓ be two integers such that $k \leq \ell \leq k + \ell \leq |\Omega|$. Then $\ker(\zeta_{P(\Omega)}^{k,\ell}) = \{0\}$.*

Proof. By Lemma 3.6, Theorem 5.1, and Remark 5.2. □

Let Ω be a finite set and let k and ℓ be two integers such that $0 \leq k \leq \ell \leq k + \ell \leq |\Omega|$. By virtue of Lemma 5.3, $\ker(\zeta_{P(\Omega)}^{k,\ell}) = \{0\}$. So, to prove Conjecture 2.10 via Lemma 3.2, we want to have $s^* \in \text{hEnd}_{\ell,k}(P(\Omega))$ for all $s \in \Omega^\Omega$. It is a pity that what we can have instead is $s^* \in \text{hEnd}_{k,\ell}(P(\Omega))$ for all $s \in \Omega^\Omega$.

For any transformation g on a set Ω , we associate a partition $\ker_\Omega(g)$ of Ω in which two elements α and β fall into the same part provided $\alpha g = \beta g$, and we call $\ker_\Omega(g)$ the *kernel* of g . Note that $\ker_\Omega(g_1 g_2) = \ker_\Omega(g_2) g_1^*$ for all $g_1, g_2 \in T(\Omega)$. For any transformation semigroup S on Ω , let $P^S(\Omega)$ stand for the set $\{\ker_\Omega(s) : s \in S\} = \{\ker_\Omega(\text{Id}_\Omega) s^* : s \in S\}$, and call it the *kernel partition subposet induced by S* . It is clear that $P^S(\Omega)$ is invariant under the action of the kernel space S^* . Inheriting the rank function on P_Ω , $P^S(\Omega)$ is still a valuated poset.

For a permutation group, all its elements have the same kernel. For a transformation semigroup, the existence of different kernels may make some arguments for permutation groups invalid. It looks interesting to study the action of the kernel space S^* on the kernel partition subposet $P^S(\Omega)$.

Example 5.4. Consider the Černý automaton $\mathcal{C}_4 = \Gamma(S, G)$ as illustrated in Figure 3, where $G = \{a, b\}$. All partitions of $\{1, 2, 3, 4\}$, excepting $\{\{0, 2\}, \{1, 3\}\}$ which is displayed in red in Figure 3, belong to $P^S(\Omega)$. One can check that

$$WS(S^*|_{P^S(\Omega)}, P^S(\Omega)) = (1, 1, 1, 1) \text{ and } SS(S^*|_{P^S(\Omega)}, P^S(\Omega)) = (1, 2, 1, 1),$$

both of which being unimodal.

Example 5.5. Let $\Omega = \{1, \dots, 6\}$ and let $S = \langle r, b \rangle$ be the transformation semigroup acting on Ω as defined in Example 2.5. Simple calculations shows that $P^S(\Omega)$ is given by

$$\{\{\{1\}, \{2\}, \{3\}, \{4\}, \{5\}, \{6\}\}, \{\{1, 2\}, \{3, 4\}, \{5, 6\}\}, \{\{1, 6\}, \{2, 3\}, \{4, 5\}\}\}.$$

One can further check that $WS(S^*|_{P^S(\Omega)}, P^S(\Omega)) = SS(S^*|_{P^S(\Omega)}, P^S(\Omega)) = (1, 0, 0, 1)$. If you delete those 0-entries (equivalently, adjusting the rank function for $P^S(\Omega)$), the resulting sequence $(1, 1)$ is still unimodal.

6 Linear space

6.1 Top-heavy shape

Let n be the size of a nonempty set Ω . Let k and ℓ be two integers satisfying $0 \leq k \leq \ell \leq n$. Let q be a prime power. As q -analogues of the set incidence operator specified in Equation (3.1), we define two linear transformations $M_{q,n}^{k,\ell}: \mathbb{Q}^{\mathcal{P}_{q,n}^k} \rightarrow \mathbb{Q}^{\mathcal{P}_{q,n}^\ell}$ and $N_{q,n}^{k,\ell}: \mathbb{Q}^{\mathcal{A}_{q,n-1}^k} \rightarrow \mathbb{Q}^{\mathcal{A}_{q,n-1}^\ell}$ as follows:

$$(M_{q,n}^{k,\ell}(f))(Y) \doteq \sum_{X \leq Y, X \in \mathcal{P}_{q,n}^k} f(X),$$

and

$$(N_{q,n}^{k,\ell}(f'))(Y') \doteq \sum_{X' \leq Y', X' \in \mathcal{A}_{q,n-1}^k} f(X'),$$

for all $f \in \mathbb{Q}^{\mathcal{P}_{q,n}^k}$, $Y \in \mathcal{P}_{q,n}^\ell$ and $f' \in \mathbb{Q}^{\mathcal{A}_{q,n-1}^k}$, $Y' \in \mathcal{A}_{q,n-1}^\ell$.

Kantor [30, Theorem] obtained a q -analogue of Gottlieb's Theorem [25, Corollary 2], which implies that Conjecture 3.4 holds for affine/projective geometries.

Lemma 6.1 (Kantor). *Let n be a positive integer. Let k and ℓ be two nonnegative integers such that $k \leq \ell \leq k + \ell \leq n$ and let q be any prime power. Then both $M_{q,n}^{k,\ell}$ and $N_{q,n-1}^{k,\ell}$ are injective.*

Proof of Theorem 2.12. Let k and ℓ be two integers such that $0 \leq k \leq \ell \leq k + \ell \leq n$. Note that $S^{\mathcal{P}} \subseteq \text{hEnd}_{k,\ell}(\mathcal{P}_{q,n})$ and $T^{\mathcal{A}} \subseteq \text{hEnd}_{k,\ell}(\mathcal{A}_{q,n-1})$. Since both $\mathcal{P}_{q,n}$ and $\mathcal{A}_{q,n-1}$ are (ℓ, k) -finite, the result follows readily from Lemmas 3.2, 3.6 and 6.1. \square

6.2 Duality: A result of Stanley

First Proof of Theorem 2.16. Let F be a field and Ω be a set. For each linear subspace $U \leq F^\Omega$, let U^\perp be the subspace of F^Ω given by

$$U^\perp \doteq \{f \in F^\Omega : \sum_{\omega \in \Omega} f(\omega)g(\omega) = 0 \text{ for all } g \in U\}.$$

Take a matrix $A \in F^{\Omega \times \Omega}$ and record its transpose by A^\top . For any $f \in F^\Omega$, which can be thought of as a row vector indexed by Ω , the image of f under the action of A , written as fA , can be thought of as the product of the row vector f and the matrix A . The matrix A induces a transformation \bar{A} on $\text{Gr}(F^\Omega)$ such that $U \in \text{Gr}(F^\Omega)$ is sent to

$$\begin{array}{ccc}
 f & \xrightarrow{M_{q,n}^{k,\ell}} & M_{q,n}^{k,\ell}(f) \\
 g^{\dagger,k} \downarrow & & \downarrow g^{\dagger,\ell} \\
 fg^{\dagger,k} & \xrightarrow{M_{q,n}^{k,\ell}} & M_{q,n}^{k,\ell}(fg^{\dagger,k})
 \end{array}$$

Figure 6: The incidence operator intertwines with every linear isomorphism g .

$U\hat{A} \doteq \{fA : f \in U\}$. It is easy to see that for any $U, W \in \text{Gr}(V)$ we have the implication

$$U\hat{A} = W \implies W^\perp \hat{A}^\top \leq U^\perp; \quad (6.1)$$

especially, when $A \in \text{GL}_n(F)$ it holds

$$U\hat{A} = W \iff W^\perp \hat{A}^\top = U^\perp. \quad (6.2)$$

According to Taussky and Zassenhaus [63, Theorem 1], we can find $P \in \text{GL}_n(F)$ such that $P = P^\top$ and $A^\top = PAP^{-1}$. This means that Equations (6.1), (6.2) become

$$U\hat{A} = W \implies (W^\perp \hat{P})\hat{A} \leq U^\perp \hat{P}$$

and

$$U\hat{A} = W \iff (W^\perp \hat{P})\hat{A} = U^\perp \hat{P}, \quad (6.3)$$

respectively. It is well-known that q -binomial coefficients (Gaussian coefficients) occur in pairs, namely in any n -dimensional linear space over a finite field, the number of k -dimensional subspaces is equal to the number of $(n - k)$ -dimensional subspaces [24, Proposition 5.31] [60, Section 3]. In general, as a consequence of Equation 6.3, for any $A \in \text{GL}_n(F)$, the number of k -dimension subspaces of F^n fixed by \hat{A} equals to the number of $(n - k)$ -dimension subspaces of F^n fixed by \hat{A} . If F is a finite field and G is a subgroup of $\text{GL}_n(F)$, in view of the Orbit Counting Lemma (also known as Burnside's Lemma), the above discussion leads to a proof of Theorem 2.16. \square

Second Proof of Theorem 2.16. Let $G \leq \text{GL}_n(\mathbb{F}_q)$ and let k be a positive integer fulfilling $k \leq \frac{n}{2}$. The group G can be seen as a permutation group acting on both $\text{Gr}(n - k, \mathbb{F}_q^n) = \mathcal{P}_{q,n}^k$ and $\text{Gr}(n - k, \mathbb{F}_q^n) = \mathcal{P}_{q,n}^{n-k}$; we use W_k and W_{n-k} for the two permutation modules accordingly. From Lemma 6.1, we see that $M_{q,n}^{k,n-k}$ is an \mathbb{F}_q -linear isomorphism from $\mathcal{P}_{q,n}^k$ to $\mathcal{P}_{q,n}^{n-k}$. From Figure 5 and Example 3.7, we have the commutative diagram in Figure 6 for $2 \leq k \leq \frac{n}{2}$; assuming that g comes from the group G , clearly our deduction of Figure 5 shows that Figure 6 is also valid for $k = 1$. This then shows that W_k and W_{n-k} are isomorphic permutation modules for G . In particular, the number of orbits of G on $\mathcal{P}_{q,n}^k$ and the number of its orbits on $\mathcal{P}_{q,n}^{n-k}$ must be equal. \square

By examining the proofs of Theorem 2.16, we intend to understand the challenge of extending some results on group actions to that on semigroup actions. The above two

proofs apply to a set of invertible linear operators over finite linear spaces. If we have a single linear operator $A \in \text{Mat}_n(F)$, by considering its action on the linear space obtained by “collapsing” the eventual kernel of A to zero, we may somehow still say something similar to above. When we have a subsemigroup S of the full linear transformation monoid acting on a finite linear space, different elements of S may have different eventual kernels and that makes it nontrivial to glean global information about the semigroup action.

7 Vámos matroid

Proof of Example 2.18. A simple calculation shows that $\ker(\zeta_{F(M)}^{k,\ell}) = \{0\}$ for $(k, \ell) \in \{(1, 2), (2, 3)\}$. Let $f \in S$ and let $f': F(M) \rightarrow F(M)$ be the map sending each flat $X \in F(M)$ to the minimum flat containing $X\bar{f}$ in M . By Lemma 3.2, we will be done if we can show that $f' \in \text{hEnd}_{\ell,k}(F(M))$ for $(k, \ell) = (1, 2), (2, 3)$.

If we know that f is a bijection or that $|\mathcal{E}_M \bar{f}| \leq 2$, we can easily check that $f' \in \text{hEnd}_{\ell,k}(F(M))$, as wanted. We intend to find a contradiction under the hypothesis that neither of them holds.

By assumption, we can take three distinct elements x, y, z in $\mathcal{E}_M \bar{f}$ such that $|xf^{-1}| \geq 2$. Let A be the minimum flat containing $\{x, y, z\}$ and let $B = Af^{-1}$. Observe that $|A| \in \{3, 4\}$. Since f is a strong map, B is a flat containing at least four elements and so $|B| \in \{4, 8\}$.

CASE 1: $|B| = 8$.

Take any $X \in \binom{A}{2}$. Note that X must be a flat and thus so is Xf^{-1} . Since $|\mathcal{E}_M \bar{f}| \geq 3$, we deduce that the flat Xf^{-1} is not equal to \mathcal{E}_M and so $|Xf^{-1}| \leq 4$. Considering that $|A| \in \{3, 4\}$, we find that $|A| = 4$ and each element in A has two perimages under f . Note that every element in $\binom{A}{2}$ is a flat. It follows that $\{Xf^{-1} : X \in \binom{A}{2}\}$ is a set of six distinct flats and each of them contains four elements, which cannot happen for the Vámos matroid M .

CASE 2: $|B| = 4$.

Thanks to the assumption of $|B| = 4$, we see that $C = \{x, y\}$ is a flat in M satisfying $|Cf^{-1}| = 3$. Note that no three-elements subset of any four-elements flat in M can be a flat. This means that Cf^{-1} is not a flat, violating the assumption that f is a strong map. \square

8 Concluding remarks


We have discussed some top-heavy phenomena for transformation semigroups acting on Boolean semirings, affine/projective geometries, and flat lattice of Vámos matroid; see Theorems 2.1 and 2.12 and Example 2.18. But some problems remain, say Question 2.2, 2.3 and 2.8, Conjecture 2.10 and Question 2.11, and Question 2.15. Our work relies on various injectivity results, say Lemmas 4.1, 5.3 and 6.1, which can all be deduced from Theorem 5.1 and Remark 5.2. We may think of Conjecture 3.4 as a natural companion to [28, Conjecture 7]. Since our results on comparing the number of components inside P_k and that of P_ℓ for various valuated posets P come from the injectivity of the relevant incidence operators (Lemma 3.2), we indeed have an injective map from components of P_k to that of P_ℓ which respects the poset structure. It is noteworthy that we do not find any general results on the unimodality of the strong/weak shape of a semigroup action on a valuated poset. We are wondering if the theory on the Lefschetz properties of commutative graded

algebras [27] will be useful in this line of research.

Penttila and Siciliano [47, Lemma 3.1] suggested a machinery (Lemma 3.6) to remove certain finiteness assumption. But we do not see any way to solve Question 1.2 and 2.17 by this means. Since there are many other approaches to go from finite to infinite [53], it will not be a surprise if Question 1.2 has a positive solution as simple as that for Theorem 2.12. Here is another such question. By our definition, a valuated poset only has nonnegative integers as ranks of its elements. We may allow ranks to be any (not necessarily finite) cardinal number and then examine all the work in this paper again. A few results of this kinds from the literature have been addressed at the end of Section 2.1.

ORCID iDs

Yaokun Wu  <https://orcid.org/0000-0002-6811-7067>

Yinfeng Zhu  <https://orcid.org/0000-0003-1724-5250>

References

- [1] K. Adiprasito, J. Huh and E. Katz, Hodge theory for combinatorial geometries, *Ann. of Math.* (2) **188** (2018), 381–452, doi:10.4007/annals.2018.188.2.1.
- [2] J. André, J. a. Araújo and P. J. Cameron, The classification of partition homogeneous groups with applications to semigroup theory, *J. Algebra* **452** (2016), 288–310, doi:10.1016/j.jalgebra.2015.12.025.
- [3] C. A. Athanasiadis, The symmetric group action on rank-selected posets of injective words, *Order* **35** (2018), 47–56, doi:10.1007/s11083-016-9417-9.
- [4] L. Babai and P. Frankl, *Linear Algebra Methods in Combinatorics With Applications to Geometry and Computer Science*, Department of Computer Science, The University of Chicago, 1992.
- [5] A. Bachem and A. Wanka, Separation theorems for oriented matroids, *Discrete Math.* **70** (1988), 303–310, doi:10.1016/0012-365x(88)90006-4.
- [6] M. Baker, Hodge theory in combinatorics, *Bull. Am. Math. Soc. (N.S.)* **55** (2018), 57–80, doi:10.1090/bull/1599.
- [7] M. Baker and N. Bowler, Matroids over partial hyperstructures, *Adv. Math.* **343** (2019), 821–863, doi:10.1016/j.aim.2018.12.004.
- [8] M. Benoumhani, Log-concavity of Whitney numbers of Dowling lattices, *Adv. in Appl. Math.* **22** (1999), 186–189, doi:10.1006/aama.1998.0621.
- [9] R. D. Bercov and C. R. Hobby, Permutation groups on unordered sets, *Math. Z.* **115** (1970), 165–168, doi:10.1007/bf01109854.
- [10] A. Björner and T. Ekedahl, On the shape of Bruhat intervals, *Ann. of Math.* (2) **170** (2009), 799–817, doi:10.4007/annals.2009.170.799.
- [11] F. Brenti, Log-concave and unimodal sequences in algebra, combinatorics, and geometry: an update, in: *Jerusalem Combinatorics '93*, Amer. Math. Soc., Providence, RI, volume 178 of *Contemp. Math.*, pp. 71–89, 1994, doi:10.1090/conm/178/01893.
- [12] M. Brown, Weak n -homogeneity implies weak $(n - 1)$ -homogeneity, *Proc. Am. Math. Soc.* **10** (1959), 644–647, doi:10.2307/2033668.
- [13] H. Bruhn, R. Diestel, M. Kriesell, R. Pendavingh and P. Wollan, Axioms for infinite matroids, *Adv. Math.* **239** (2013), 18–46, doi:10.1016/j.aim.2013.01.011.

- [14] V. M. Bukhshtaber, A. M. Vershik, S. A. Evdokimov and I. N. Ponomarenko, Combinatorial algebras and multivalued involutive groups, *Funct. Anal. Its Appl.* **30** (1996), 158–162, doi:10.1007/bf02509502.
- [15] P. J. Cameron, Transitivity of permutation groups on unordered sets, *Math. Z.* **148** (1976), 127–139, doi:10.1007/bf01214702.
- [16] P. J. Cameron, Oligomorphic permutation groups, in: *Perspectives in mathematical sciences. II*, World Sci. Publ., Hackensack, NJ, volume 8 of *Stat. Sci. Interdiscip. Res.*, pp. 37–61, 2009, doi:10.1142/9789814273657_0003.
- [17] A. Chubarev and I. Pinelis, Fundamental theorem of geometry without the 1-to-1 assumption, *Proc. Am. Math. Soc.* **127** (1999), 2735–2744, doi:10.1090/s0002-9939-99-05280-6.
- [18] D. de Caen, A note on the ranks of set-inclusion matrices, *Electron. J. Comb.* **8** (2001), Note 5, 2, doi:10.37236/1590.
- [19] E. Delucchi and S. Riedel, Group actions on semimatroids, *Adv. in Appl. Math.* **95** (2018), 199–270, doi:10.1016/j.aam.2017.11.001.
- [20] P. Dembowski, *Finite Geometries: Reprint of the 1968 Edition*, Springer Berlin, Heidelberg, 2012, doi:10.1007/978-3-642-62012-6.
- [21] E. Dobson and A. Malnič, Groups that are transitive on all partitions of a given shape, *J. Algebraic Comb.* **42** (2015), 605–617, doi:10.1007/s10801-015-0593-2.
- [22] R. Dougherty, C. Freiling and K. Zeger, Networks, matroids, and non-Shannon information inequalities, *IEEE Trans. Inform. Theory* **53** (2007), 1949–1969, doi:10.1109/tit.2007.896862.
- [23] T. A. Dowling and R. M. Wilson, Whitney number inequalities for geometric lattices, *Proc. Am. Math. Soc.* **47** (1975), 504–512, doi:10.2307/2039773.
- [24] G. Gordon and J. McNulty, *Matroids: A Geometric Introduction*, Cambridge University Press, Cambridge, 2012, doi:10.1017/cbo9781139049443.
- [25] D. H. Gottlieb, A certain class of incidence matrices, *Proc. Am. Math. Soc.* **17** (1966), 1233–1237, doi:10.2307/2035716.
- [26] A. Hajnal, A remark on the homogeneity of infinite permutation groups, *Bull. London Math. Soc.* **22** (1990), 529–532, doi:10.1112/blms/22.6.529.
- [27] T. Harima, T. Maeno, H. Morita, Y. Numata, A. Wachi and J. Watanabe, *The Lefschetz properties*, volume 2080 of *Lecture Notes in Mathematics*, Springer, Heidelberg, 2013, doi:10.1007/978-3-642-38206-2.
- [28] J. Huh and B. Wang, Enumeration of points, lines, planes, etc, *Acta Math.* **218** (2017), 297–317, doi:10.4310/acta.2017.v218.n2.a2.
- [29] O. Johnson, I. Kontoyiannis and M. Madiman, Log-concavity, ultra-log-concavity, and a maximum entropy property of discrete compound Poisson measures, *Discrete Appl. Math.* **161** (2013), 1232–1250, doi:10.1016/j.dam.2011.08.025.
- [30] W. M. Kantor, On incidence matrices of finite projective and affine spaces, *Math. Z.* **124** (1972), 315–318, doi:10.1007/bf01113923.
- [31] W. M. Kantor, Line-transitive collineation groups of finite projective spaces, *Israel J. Math.* **14** (1973), 229–235, doi:10.1007/bf02764881.
- [32] W. M. Kantor, Homogeneous designs and geometric lattices, *J. Comb. Theory Ser. A* **38** (1985), 66–74, doi:10.1016/0097-3165(85)90022-6.
- [33] J. P. S. Kung, Strong maps, in: *Theory of matroids*, Cambridge Univ. Press, Cambridge, volume 26 of *Encyclopedia Math. Appl.*, pp. 224–253, 1986, doi:10.1017/cbo9780511629563.011.

- [34] J. P. S. Kung, The Radon transforms of a combinatorial geometry. II. Partition lattices, *Adv. Math.* **101** (1993), 114–132, doi:10.1006/aima.1993.1044.
- [35] J. P. S. Kung and H. Q. Nguyen, Weak maps, in: *Theory of matroids*, Cambridge Univ. Press, Cambridge, volume 26 of *Encyclopedia Math. Appl.*, pp. 254–271, 1986, doi:10.1017/cbo9780511629563.012.
- [36] J. P. S. Kung, G.-C. Rota and C. H. Yan, *Combinatorics: the Rota way*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2009, doi:10.1017/cbo9780511803895.
- [37] E. H. Lieb, Concavity properties and a generating function for Stirling numbers, *J. Comb. Theory.* **5** (1968), 203–206, doi:10.1016/s0021-9800(68)80057-2.
- [38] D. Livingstone and A. Wagner, Transitivity of finite permutation groups on unordered sets, *Math. Z.* **90** (1965), 393–403, doi:10.1007/bf01112361.
- [39] W. J. Martin and B. E. Sagan, A new notion of transitivity for groups and sets of permutations, *J. London Math. Soc. (2)* **73** (2006), 1–13, doi:10.1112/s0024610705022441.
- [40] J. H. Mason, Matroids: unimodal conjectures and Motzkin’s theorem, in: *Combinatorics (Proc. Conf. Combinatorial Math., Math. Inst., Oxford, 1972)*, 1972 pp. 207–220.
- [41] V. B. Mnukhin and I. J. Siemons, On the Livingstone-Wagner theorem, *Electron. J. Comb.* **11** (2004), Research Paper 29, 8, doi:10.37236/1782.
- [42] M. M. Mukhopadhyay and E. Vassiliev, On the Vámos matroid, homogeneous pregeometries and dense pairs, *Australas. J. Comb.* **75** (2019), 158–170, https://ajc.maths.uq.edu.au/?page=get_volumes&volume=75.
- [43] Y. Nakashima, A partial generalization of the Livingstone-Wagner theorem, *Ars Math. Contemp.* **2** (2009), 207–215, doi:10.26493/1855-3974.92.46f.
- [44] P. M. Neumann, Homogeneity of infinite permutation groups, *Bull. London Math. Soc.* **20** (1988), 305–312, doi:10.1112/blms/20.4.305.
- [45] M. Pankov, *Geometry of Semilinear Embeddings: Relations to Graphs and Codes*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2015, doi:10.1142/9465.
- [46] M. Pankov, Apartments preserving transformations of Grassmannians of infinite-dimensional vector spaces, *Linear Algebra Appl.* **531** (2017), 498–509, doi:10.1016/j.laa.2017.06.016.
- [47] T. Penttilä and A. Siciliano, On the incidence map of incidence structures, *Ars Math. Contemp.* **20** (2021), 51–68, doi:10.26493/1855-3974.1996.db7.
- [48] A. Putman, The fundamental theorem of projective geometry, 2010, <https://www3.nd.edu/~andyp/notes/>.
- [49] F. P. Ramsey, On a Problem of Formal Logic, *Proc. London Math. Soc. (2)* **30** (1929), 264–286, doi:10.1112/plms/s2-30.1.264.
- [50] G.-C. Rota, Combinatorial theory, old and new, in: *Actes du Congrès International des Mathématiciens (Nice, 1970)*, Tome 3, Gauthier-Villars, Paris, pp. 229–233, 1971.
- [51] P. Roy, Another proof that weak n -homogeneity implies weak $(n - 1)$ -homogeneity, *Proc. Am. Math. Soc.* **49** (1975), 515–516, doi:10.2307/2040675.
- [52] P. Šemrl, The Optimal Version of Hua’s Fundamental Theorem of Geometry of Rectangular Matrices, *Mem. Am. Math. Soc.* **232** (2014), vi+74, <https://bookstore.ams.org/memo-232-1089>.
- [53] J.-P. Serre, How to use finite fields for problems concerning infinite fields, in: *Arithmetic, Geometry, Cryptography and Coding Theory*, Amer. Math. Soc., Providence, RI, volume 487 of *Contemp. Math.*, pp. 183–193, 2009, doi:10.1090/conm/487/09532.

- [54] P. D. Seymour, On secret-sharing matroids, *J. Comb. Theory Ser. B* **56** (1992), 69–73, doi:10.1016/0095-8956(92)90007-k.
- [55] S. Shelah and S. Thomas, Homogeneity of infinite permutation groups, *Arch. Math. Logic* **28** (1989), 143–147, doi:10.1007/bf01633987.
- [56] J. Siemons, On partitions and permutation groups on unordered sets, *Arch. Math. (Basel)* **38** (1982), 391–403, doi:10.1007/bf01304806.
- [57] M. T. Stamps, Topological representations of matroid maps, *J. Algebraic Comb.* **37** (2013), 265–287, doi:10.1007/s10801-012-0366-0.
- [58] R. P. Stanley, Two combinatorial applications of the Aleksandrov-Fenchel inequalities, *J. Comb. Theory Ser. A* **31** (1981), 56–65, doi:10.1016/0097-3165(81)90053-4.
- [59] R. P. Stanley, Some aspects of groups acting on finite posets, *J. Comb. Theory Ser. A* **32** (1982), 132–161, doi:10.1016/0097-3165(82)90017-6.
- [60] R. P. Stanley, $GL(n, \mathbb{C})$ for combinatorialists, in: *Surveys in combinatorics (Southampton, 1983)*, Cambridge Univ. Press, Cambridge, volume 82 of *London Math. Soc. Lecture Note Ser.*, pp. 187–199, 1983, <https://math.mit.edu/~rstan/pubs/>.
- [61] R. P. Stanley, Log-concave and unimodal sequences in algebra, combinatorics, and geometry, in: *Graph theory and its applications: East and West (Jinan, 1986)*, New York Acad. Sci., New York, volume 576 of *Ann. New York Acad. Sci.*, pp. 500–535, 1989, doi:10.1111/j.1749-6632.1989.tb16434.x.
- [62] T. Stokes, Semigroup actions on posets and preimage quasi-orders, *Semigroup Forum* **85** (2012), 540–558, doi:10.1007/s00233-012-9436-9.
- [63] O. Taussky and H. Zassenhaus, On the similarity transformation between a matrix and its transpose, *Pacific J. Math.* **9** (1959), 893–896, <http://projecteuclid.org/euclid.pjm/1103039127>.
- [64] J. H. van Lint and R. M. Wilson, *A Course in Combinatorics*, Cambridge University Press, Cambridge, 2nd edition, 2001, doi:10.1017/cbo9780511987045.
- [65] O. Y. Viro, On basic concepts of tropical geometry, *Proc. Steklov Inst. Math.* **273** (2011), 252–282, doi:10.1134/s0081543811040134.
- [66] Z.-X. Wan, *Geometry of Matrices*, World Scientific Publishing Co., Inc., River Edge, NJ, 1996, doi:10.1142/9789812830234.
- [67] Y. Wu, Z. Xu and Y. Zhu, A five-element transformation monoid on labelled trees, *European J. Comb.* **80** (2019), 401–415, doi:10.1016/j.ejc.2018.07.014.
- [68] Y. Wu and S. Zhao, Incidence matrix and cover matrix of nested interval orders, *Electron. J. Linear Algebra* **23** (2012), 43–65, doi:10.13001/1081-3810.1504.