



ARS MATHEMATICA
CONTEMPORANEA

ISSN 1855-3966 (printed edn.), ISSN 1855-3974 (electronic edn.)

ARS MATHEMATICA CONTEMPORANEA 16 (2019) 119–139

<https://doi.org/10.26493/1855-3974.1490.eea>

(Also available at <http://amc-journal.eu>)

The Doyen-Wilson theorem for 3-sun systems*

Giovanni Lo Faro , Antoinette Tripodi

Dipartimento di Scienze Matematiche e Informatiche, Scienze Fisiche e Scienze della Terra, Università di Messina, Messina, Italia

Received 25 September 2017, accepted 18 April 2018, published online 20 September 2018

Abstract

A solution to the existence problem of G -designs with given subdesigns is known when G is a triangle with $p = 0, 1$, or 2 disjoint pendent edges: for $p = 0$, it is due to Doyen and Wilson, the first to pose such a problem for Steiner triple systems; for $p = 1$ and $p = 2$, the corresponding designs are kite systems and bull designs, respectively. Here, a complete solution to the problem is given in the remaining case where G is a 3-sun, i.e. a graph on six vertices consisting of a triangle with three pendent edges which form a 1-factor.

Keywords: 3-sun systems, embedding, difference set.

Math. Subj. Class.: 05B05, 05B30

1 Introduction

If G is a graph, then let $V(G)$ and $E(G)$ be the vertex-set and edge-set of G , respectively. The graph K_n denotes the complete graph on n vertices. The graph $K_m \setminus K_n$ has vertex-set $V(K_m)$ containing a distinguished subset H of size n ; the edge-set of $K_m \setminus K_n$ is $E(K_m)$ but with the $\binom{n}{2}$ edges between the n distinguished vertices of H removed. This graph is sometimes referred to as a complete graph of order m with a *hole* of size n .

Let G and Γ be finite graphs. A G -*design* of Γ is a pair (X, \mathcal{B}) where $X = V(\Gamma)$ and \mathcal{B} is a collection of isomorphic copies of G (*blocks*), whose edges partition $E(\Gamma)$. If $\Gamma = K_n$, then we refer to such a design as a G -*design of order n* .

A G -design (X_1, \mathcal{B}_1) of order n is said to be *embedded* in a G -design (X_2, \mathcal{B}_2) of order m provided $X_1 \subseteq X_2$ and $\mathcal{B}_1 \subseteq \mathcal{B}_2$ (we also say that (X_1, \mathcal{B}_1) is a *subdesign* (or *subsystem*) of (X_2, \mathcal{B}_2) or (X_2, \mathcal{B}_2) contains (X_1, \mathcal{B}_1) as subdesign). Let $N(G)$ denote the set of integers n such that there exists a G -design of order n . A natural question to ask is: given $n, m \in N(G)$, with $m > n$, and a G -design (X, \mathcal{B}) of order n , does exists a G -design of order m containing (X, \mathcal{B}) as subdesign? Doyen and Wilson were the first to

*Supported by I.N.D.A.M. (G.N.S.A.G.A.).

E-mail addresses: lofaro@unime.it (Giovanni Lo Faro), atripodi@unime.it (Antoinette Tripodi)

pose this problem for $G = K_3$ (Steiner triple systems) and in 1973 they showed that given $n, m \in N(K_3) = \{v : v \equiv 1, 3 \pmod{6}\}$, then *any Steiner triple system of order n can be embedded in a Steiner triple system of order m if and only if $m \geq 2n+1$ or $m = n$* (see [3]). Over the years, any such problem has come to be called a “Doyen-Wilson problem” and any solution a “Doyen-Wilson type theorem”. The work along these lines is extensive ([1, 4, 5, 6, 7, 8, 9, 10, 13]) and the interested reader is referred to [2] for a history of this problem.

In particular, taking into consideration the case where G is a triangle with $p = 0, 1, 2$, or 3 mutually disjoint pendent edges, a solution to the Doyen-Wilson problem is known when $p = 0$ (Steiner triple systems, [3]), $p = 1$ (kite systems, [9, 10]) and $p = 2$ (bull designs, [4]). Here, we deal with the remaining case ($p = 3$) where G is a 3-sun, i.e. a graph on six vertices consisting of a triangle with three pendent edges which form a 1-factor, by giving a complete solution to the Doyen-Wilson problem for G -designs where G is a 3-sun (3-sun systems).

2 Notation and basic lemmas

The 3-sun consisting of the triangle (a, b, c) and the three mutually disjoint pendent edges $\{a, d\}, \{b, e\}, \{c, f\}$ is denoted by $(a, b, c; d, e, f)$. A 3-sun system of order n (briefly, $3SS(n)$) exists if and only if $n \equiv 0, 1, 4, 9 \pmod{12}$ and if (X, \mathcal{S}) is a $3SS(n)$, then $|\mathcal{S}| = \frac{n(n-1)}{12}$ (see [14]).

Let $n, m \equiv 0, 1, 4, 9 \pmod{12}$, with $m = u + n$, $u \geq 0$. The Doyen-Wilson problem for 3-sun systems is equivalent to the existence problem of decompositions of $K_{u+n} \setminus K_n$ into 3-suns.

Let r and s be integers with $r < s$, define $[r, s] = \{r, r+1, \dots, s\}$ and $[s, r] = \emptyset$. Let $Z_u = [0, u-1]$ and $H = \{\infty_1, \infty_2, \dots, \infty_t\}$, $H \cap Z_u = \emptyset$. If $S = (a, b, c; d, e, f)$ is a 3-sun whose vertices belong to $Z_u \cup H$ and $i \in Z_u$, let $S+i = (a+i, b+i, c+i; d+i, e+i, f+i)$, where the sums are modulo u and $\infty+i = \infty$, for every $\infty \in H$. The set $(S) = \{S+i : i \in Z_u\}$ is called the *orbit of S under Z_u* and S is a *base block* of (S) .

To solve the Doyen-Wilson problem for 3-sun systems we use the *difference method* (see [11, 12]). For every pair of distinct elements $i, j \in Z_u$, define $|i-j|_u = \min\{|i-j|, u - |i-j|\}$ and set $D_u = \{|i-j|_u : i, j \in Z_u\} = \{1, 2, \dots, \lfloor \frac{u}{2} \rfloor\}$. The elements of D_u are called *differences* of Z_u . For any $d \in D_u$, $d \neq \frac{u}{2}$, we can form a single 2-factor $\{\{i, d+i\} : i \in Z_u\}$, while if u is even and $d = \frac{u}{2}$, then we can form a 1-factor $\{\{i, i+\frac{u}{2}\} : 0 \leq i \leq \frac{u}{2}-1\}$. It is also worth remarking that 2-factors obtained from distinct differences are disjoint from each other and from the 1-factor.

If $D \subseteq D_u$, denote by $\langle Z_u \cup H, D \rangle$ the graph with vertex-set $V = Z_u \cup H$ and the edge-set $E = \{(i, j) : |i-j|_u = d, d \in D\} \cup \{(\infty, i) : \infty \in H, i \in Z_u\}$. The graph $\langle Z_u \cup H, D_u \rangle$ is the complete graph $K_{u+t} \setminus K_t$ based on $Z_u \cup H$ and having H as a hole. The elements of H are called *infinity points*.

Let X be a set of size $n \equiv 0, 1, 4, 9 \pmod{12}$. The aim of the paper is to decompose the graph $\langle Z_u \cup X, D_u \rangle$ into 3-suns. To obtain our main result the $\langle Z_u \cup X, D_u \rangle$ will be regarded as a union of suitable edge-disjoint subgraphs of type $\langle Z_u \cup H, D \rangle$ (where $H \subseteq X$ may be empty, while $D \subseteq D_u$ is always non empty) and then each subgraph will be decomposed into 3-suns by using the lemmas given in this section. From here on suppose $u \equiv 0, 1, 3, 4, 5, 7, 8, 9, 11 \pmod{12}$.

Lemmas 2.1–2.4 give decompositions of subgraphs of type $\langle Z_u \cup H, D \rangle$ where D

contains particular differences, more precisely, $D = \{2\}$, $D = \{2, 4\}$ or $D = \{1, \frac{u}{3}\}$.

Lemma 2.1. *Let $u \equiv 0 \pmod{4}$, $u \geq 8$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2\}, \{2\} \rangle$ can be decomposed into 3-suns.*

Proof. Consider the 3-suns

$$\begin{aligned} & (\infty_1, 2 + 4i, 4i; 3 + 4i, 4 + 4i, \infty_2), \\ & (\infty_2, 3 + 4i, 1 + 4i; 2 + 4i, 5 + 4i, \infty_1), \end{aligned}$$

for $i = 0, 1, \dots, \frac{u}{4} - 1$. \square

Lemma 2.2. *Let $u \equiv 0 \pmod{12}$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{2\} \rangle$ can be decomposed into 3-suns.*

Proof. Consider the 3-suns

$$\begin{aligned} & (\infty_1, 12i, 2 + 12i; 7 + 12i, \infty_3, \infty_4), \\ & (\infty_1, 4 + 12i, 6 + 12i; 9 + 12i, \infty_3, \infty_4), \\ & (\infty_1, 8 + 12i, 10 + 12i; 11 + 12i, \infty_3, \infty_4), \\ & (\infty_2, 2 + 12i, 4 + 12i; 1 + 12i, \infty_3, \infty_4), \\ & (\infty_2, 6 + 12i, 8 + 12i; 7 + 12i, \infty_3, \infty_4), \\ & (\infty_2, 10 + 12i, 12 + 12i; 11 + 12i, \infty_3, \infty_4), \\ & (\infty_3, 1 + 12i, 3 + 12i; 9 + 12i, \infty_1, \infty_2), \\ & (\infty_3, 5 + 12i, 7 + 12i; 11 + 12i, \infty_1, 9 + 12i), \\ & (\infty_4, 3 + 12i, 5 + 12i; 1 + 12i, \infty_1, \infty_2), \\ & (\infty_4, 9 + 12i, 11 + 12i; 7 + 12i, \infty_2, 13 + 12i), \end{aligned}$$

for $i = 0, 1, \dots, \frac{u}{12} - 1$. \square

Lemma 2.3. *The graph $\langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{2, 4\} \rangle$, $u \geq 7$, $u \neq 8$, can be decomposed into 3-suns.*

Proof. Let $u = 4k + r$, with $r = 0, 1, 3$, and consider the 3-suns

$$\begin{aligned} & (\infty_1, 4 + 4i, 6 + 4i; 5 + 4i, 8 + 4i, \infty_4), \\ & (\infty_2, 5 + 4i, 7 + 4i; 6 + 4i, 9 + 4i, \infty_1), \\ & (\infty_3, 6 + 4i, 8 + 4i; 7 + 4i, 10 + 4i, \infty_2), \\ & (\infty_4, 7 + 4i, 9 + 4i; 8 + 4i, 11 + 4i, \infty_3), \end{aligned}$$

for $i = 0, 1, \dots, k - 3$, $k \geq 3$, plus the following blocks as the case may be.

If $r = 0$,

$$\begin{aligned} & (\infty_1, 0, 2; 1, 4, \infty_4), \\ & (\infty_2, 1, 3; 2, 5, \infty_1), \\ & (\infty_3, 2, 4; 3, 6, \infty_2), \\ & (\infty_4, 3, 5; 4, 7, \infty_3), \\ & (\infty_1, 4k - 4, 4k - 2; 4k - 3, 0, \infty_4), \end{aligned}$$

$$\begin{aligned} & (\infty_2, 4k-3, 4k-1; 4k-2, 1, \infty_1), \\ & (\infty_3, 4k-2, 0; 4k-1, 2, \infty_2), \\ & (\infty_4, 4k-1, 1; 0, 3, \infty_3). \end{aligned}$$

If $r = 1$,

$$\begin{aligned} & (\infty_1, 0, 2; 1, 4, \infty_2), \\ & (\infty_2, 1, 3; 0, 5, \infty_1), \\ & (\infty_3, 2, 4; 3, 6, \infty_2), \\ & (\infty_4, 3, 5; 4, 7, \infty_3), \\ & (\infty_1, 4k-4, 4k-2; 4k-3, 4k, \infty_2), \\ & (\infty_2, 4k-3, 4k-1; 4k, 0, \infty_1), \\ & (\infty_3, 4k-2, 4k; 4k-1, 1, \infty_1), \\ & (\infty_4, 4k-1, 0; 4k-2, 2, \infty_3), \\ & (\infty_4, 4k, 1; 2, 3, \infty_3). \end{aligned}$$

If $r = 3$,

$$\begin{aligned} & (\infty_1, 0, 2; 1, 4, \infty_4), \\ & (\infty_2, 1, 3; 2, 5, \infty_1), \\ & (\infty_3, 2, 4; 3, 6, \infty_2), \\ & (\infty_4, 3, 5; 4, 7, \infty_3), \\ & (\infty_1, 4k-4, 4k-2; 4k-3, 4k, \infty_4), \\ & (\infty_2, 4k-3, 4k-1; 4k-2, 4k+1, \infty_1), \\ & (\infty_3, 4k-2, 4k; 4k-1, 4k+2, \infty_2), \\ & (\infty_4, 4k-1, 4k+1; 4k, 0, \infty_3), \\ & (\infty_1, 4k, 4k+2; 4k+1, 1, \infty_4), \\ & (\infty_2, 4k+1, 0; 4k+2, 2, \infty_4), \\ & (\infty_3, 4k+2, 1; 0, 3, \infty_4). \end{aligned}$$

With regard to the difference 4 in Z_7 , note that $|4|_7 = 3$ and the seven distinct blocks obtained for $k = 1$ and $r = 3$ gives a decomposition of $\langle Z_7 \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{2, 3\} \rangle$ into 3-suns. \square

Lemma 2.4. *Let $u \equiv 0 \pmod{3}$, $u \geq 12$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_8\}, \{1, \frac{u}{3}\} \rangle$ can be decomposed into 3-suns.*

Proof. If $u \equiv 0 \pmod{6}$ consider the 3-suns:

$$\begin{aligned} & (\infty_1, 2i, \frac{u}{3} + 2i; 2\frac{u}{3} + 2i, \infty_5, \infty_6), \quad i = 0, 1, \dots, \frac{u}{6} - 1, \\ & (\infty_1, 1 + 2i, \frac{u}{3} + 1 + 2i; 2\frac{u}{3} + 1 + 2i, \infty_6, \infty_5), \quad i = 0, 1, \dots, \frac{u}{6} - 1, \\ & (\infty_2, 2\frac{u}{3} + 2i, \frac{u}{3} + 2i; 2 + 2i, 2i, \infty_5), \quad i = 0, 1, \dots, \frac{u}{6} - 2, \\ & (\infty_2, 2\frac{u}{3} + 1 + 2i, \frac{u}{3} + 1 + 2i; 3 + 2i, 1 + 2i, \infty_6), \quad i = 0, 1, \dots, \frac{u}{6} - 2, \\ & (\infty_2, u - 2, 2\frac{u}{3} - 2; 0, \frac{u}{3} - 2, \infty_5), \end{aligned}$$

$$\begin{aligned}
& (\infty_2, u - 1, 2\frac{u}{3} - 1; 1, \frac{u}{3} - 1, \infty_6), \\
& (\infty_3, 2i, 1 + 2i; 2\frac{u}{3} + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u}{6} - 1, \\
& (\infty_3, \frac{u}{3} + 2i, \frac{u}{3} + 1 + 2i; 2\frac{u}{3} + 1 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u}{6} - 1, \\
& (\infty_4, 1 + 2i, 2 + 2i; 2\frac{u}{3} + 2 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u}{6} - 1, \\
& (\infty_4, \frac{u}{3} + 1 + 2i, \frac{u}{3} + 2 + 2i; 2\frac{u}{3} + 1 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u}{6} - 1, \\
& (\infty_5, 2\frac{u}{3} + 2i, 2\frac{u}{3} + 1 + 2i; 1 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u}{6} - 1, \\
& (\infty_6, 2\frac{u}{3} + 3 + 2i, 2\frac{u}{3} + 4 + 2i; 2 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u}{6} - 2, \\
& (\infty_6, 2\frac{u}{3} + 1, 2\frac{u}{3} + 2; 2\frac{u}{3}, \infty_7, \infty_8).
\end{aligned}$$

If $u \equiv 3 \pmod{6}$ consider the 3-suns:

$$\begin{aligned}
& (\infty_1, 2i, \frac{u}{3} + 2i; 2\frac{u}{3} + 2i, \infty_5, \infty_6), i = 0, 1, \dots, \frac{u-3}{6}, \\
& (\infty_1, 1 + 2i, \frac{u}{3} + 1 + 2i; 2\frac{u}{3} + 1 + 2i, \infty_6, \infty_5), i = 0, 1, \dots, \frac{u-9}{6}, \\
& (\infty_2, 2\frac{u}{3} + 2i, \frac{u}{3} + 2i; 2 + 2i, 2i, \infty_5), i = 0, 1, \dots, \frac{u-9}{6}, \\
& (\infty_2, u - 1, 2\frac{u}{3} - 1; 0, \frac{u}{3} - 1, \infty_5), \\
& (\infty_2, 2\frac{u}{3} + 1 + 2i, \frac{u}{3} + 1 + 2i; 3 + 2i, 1 + 2i, \infty_6), i = 0, 1, \dots, \frac{u-15}{6}, \\
& (\infty_2, u - 2, 2\frac{u}{3} - 2; 1, \frac{u}{3} - 2, \infty_6), \\
& (\infty_3, 2i, 1 + 2i; 2\frac{u}{3} + 2i, \infty_7, \infty_8), i = 2, 3, \dots, \frac{u-3}{6}, \\
& (\infty_3, 0, 1; 2\frac{u}{3}, \infty_6, \infty_8), \\
& (\infty_3, 2, 3; 2\frac{u}{3} + 2, \infty_6, \infty_8), \\
& (\infty_3, \frac{u}{3} + 1 + 2i, \frac{u}{3} + 2 + 2i; 2\frac{u}{3} + 1 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u-9}{6}, \\
& (\infty_4, 1 + 2i, 2 + 2i; 2\frac{u}{3} + 2 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u-9}{6}, \\
& (\infty_4, \frac{u}{3} + 2i, \frac{u}{3} + 1 + 2i; 2\frac{u}{3} + 1 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u-3}{6}, \\
& (\infty_5, 2\frac{u}{3} + 2i, 2\frac{u}{3} + 1 + 2i; 1 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u-9}{6}, \\
& (\infty_6, 2\frac{u}{3} + 1 + 2i, 2\frac{u}{3} + 2 + 2i; 4 + 2i, \infty_7, \infty_8), i = 0, 1, \dots, \frac{u-15}{6}, \\
& (\infty_6, u - 2, u - 1; 2\frac{u}{3}, \infty_7, \infty_8), \\
& (\infty_7, u - 1, 0; 2, \infty_5, \infty_8). \quad \square
\end{aligned}$$

Lemmas 2.5 – 2.9 allow to decompose $\langle Z_u \cup H, D \rangle$ where u is even and D contains the difference $\frac{u}{2}$.

Lemma 2.5. Let u be even, $u \geq 8$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle$ can be decomposed into 3-suns.

Proof. Consider the 3-suns

$$\begin{aligned}
& (\infty_1, 2i, 1 + 2i; \frac{u}{2} + 2 + 2i, \frac{u}{2} + 2i, \infty_3), i = 0, 1, \dots, \frac{u}{4} - 2, \\
& (\infty_1, \frac{u}{2} - 2, \frac{u}{2} - 1; \frac{u}{2}, u - 2, \infty_3), \\
& (\infty_2, 1 + 2i, \frac{u}{2} + 1 + 2i; 2i, 2 + 2i, \infty_1), i = 0, 1, \dots, \frac{u}{4} - 1, \\
& (\infty_3, \frac{u}{2} + 1 + 2i, \frac{u}{2} + 2i; 2i, \frac{u}{2} + 2 + 2i, \infty_2), i = 0, 1, \dots, \frac{u}{4} - 1. \quad \square
\end{aligned}$$

Lemma 2.6. Let $u \equiv 0 \pmod{12}$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{1, \frac{u}{2}\} \rangle$ can be decomposed into 3-suns.

Proof. Consider the 3-suns

$$\begin{aligned}
 & (\infty_1, 6i, \frac{u}{2} + 6i; 4 + 6i, \infty_3, \infty_2), \\
 & (\infty_1, 1 + 6i, \frac{u}{2} + 1 + 6i; 5 + 6i, \infty_4, \infty_2), \\
 & (\infty_1, 2 + 6i, \frac{u}{2} + 2 + 6i; \frac{u}{2} + 3 + 6i, \infty_4, \infty_3), \\
 & (\infty_2, 1 + 6i, 6i; \frac{u}{2} + 3 + 6i, \infty_3, \infty_4), \\
 & (\infty_2, 2 + 6i, 3 + 6i; \frac{u}{2} + 4 + 6i, 1 + 6i, \infty_4), \\
 & (\infty_2, 5 + 6i, 4 + 6i; \frac{u}{2} + 5 + 6i, 6 + 6i, 3 + 6i), \\
 & (\infty_3, 3 + 6i, \frac{u}{2} + 3 + 6i; 2 + 6i, \infty_1, \frac{u}{2} + 2 + 6i), \\
 & (\infty_3, 4 + 6i, \frac{u}{2} + 4 + 6i; \frac{u}{2} + 6i, \infty_4, \infty_1), \\
 & (\infty_3, 5 + 6i, \frac{u}{2} + 5 + 6i; \frac{u}{2} + 1 + 6i, \infty_4, \infty_1), \\
 & (\infty_4, \frac{u}{2} + 1 + 6i, \frac{u}{2} + 2 + 6i; \frac{u}{2} + 3 + 6i, \frac{u}{2} + 6i, \infty_2), \\
 & (\infty_4, \frac{u}{2} + 4 + 6i, \frac{u}{2} + 5 + 6i; \frac{u}{2} + 6i, \frac{u}{2} + 3 + 6i, \frac{u}{2} + 6 + 6i),
 \end{aligned}$$

for $i = 0, 1, \dots, \frac{u}{12} - 1$. \square

Lemma 2.7. Let u be even, $u \geq 8$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, \frac{u}{2}\} \rangle$ can be decomposed into 3-suns.

Proof. Consider the 3-suns

$$\begin{aligned}
 & (\infty_1, 2i, 1 + 2i; \frac{u}{2} + 2 + 2i, \frac{u}{2} + 2i, \infty_3), \quad i = 0, 1, \dots, \frac{u}{4} - 2, \\
 & (\infty_1, \frac{u}{2} - 2, \frac{u}{2} - 1; \frac{u}{2}, u - 2, \infty_3), \\
 & (\infty_2, 1 + 2i, \frac{u}{2} + 1 + 2i; 2i, \infty_6, \infty_1), \quad i = 0, 1, \dots, \frac{u}{4} - 1, \\
 & (\infty_3, \frac{u}{2} + 1 + 2i, \frac{u}{2} + 2i; 2i, \infty_6, \infty_2), \quad i = 0, 1, \dots, \frac{u}{4} - 1, \\
 & (\infty_4, 1 + 2i, 2 + 2i; \frac{u}{2} + 2 + 2i, \infty_5, \infty_6), \quad i = 0, 1, \dots, \frac{u}{4} - 1, \\
 & (\infty_5, \frac{u}{2} + 1 + 2i, \frac{u}{2} + 2 + 2i; 2 + 2i, \infty_4, \infty_6), \quad i = 0, 1, \dots, \frac{u}{4} - 1. \quad \square
 \end{aligned}$$

Lemma 2.8. Let $u \equiv 0 \pmod{12}$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_7\}, \{1, \frac{u}{2}\} \rangle$ can be decomposed into 3-suns.

Proof. Consider the 3-suns

$$\begin{aligned}
 & (\infty_1, 6i, \frac{u}{2} + 6i; 4 + 6i, \infty_7, \infty_2), \\
 & (\infty_1, 1 + 6i, \frac{u}{2} + 1 + 6i; \frac{u}{2} + 3 + 6i, \infty_7, \infty_4), \\
 & (\infty_1, 2 + 6i, \frac{u}{2} + 2 + 6i; \frac{u}{2} + 5 + 6i, \infty_5, \infty_2), \\
 & (\infty_2, 3 + 6i, \frac{u}{2} + 3 + 6i; 6i, \infty_1, \infty_4), \\
 & (\infty_2, 4 + 6i, \frac{u}{2} + 4 + 6i; 2 + 6i, \infty_7, \infty_1), \\
 & (\infty_2, 5 + 6i, \frac{u}{2} + 5 + 6i; \frac{u}{2} + 1 + 6i, \infty_1, \infty_7), \\
 & (\infty_3, 6i, 1 + 6i; \frac{u}{2} + 6i, \infty_5, \infty_6), \\
 & (\infty_3, 2 + 6i, 3 + 6i; \frac{u}{2} + 2 + 6i, \infty_7, \infty_6), \\
 & (\infty_3, 4 + 6i, 5 + 6i; \frac{u}{2} + 5 + 6i, \infty_5, \infty_6), \\
 & (\infty_4, 1 + 6i, 2 + 6i; \frac{u}{2} + 6 + 6i, \infty_2, \infty_6), \\
 & (\infty_4, 3 + 6i, 4 + 6i; \frac{u}{2} + 4 + 6i, \infty_7, \infty_6),
 \end{aligned}$$

$$\begin{aligned}
& (\infty_4, 5 + 6i, 6 + 6i; \frac{u}{2} + 5 + 6i, \infty_7, \infty_6), \\
& (\infty_5, \frac{u}{2} + 6i, \frac{u}{2} + 1 + 6i; 1 + 6i, \infty_7, \infty_3), \\
& (\infty_5, \frac{u}{2} + 2 + 6i, \frac{u}{2} + 3 + 6i; 3 + 6i, \infty_7, \infty_3), \\
& (\infty_5, \frac{u}{2} + 4 + 6i, \frac{u}{2} + 5 + 6i; 5 + 6i, \infty_7, \frac{u}{2} + 6 + 6i), \\
& (\infty_6, \frac{u}{2} + 1 + 6i, \frac{u}{2} + 2 + 6i; \frac{u}{2} + 5 + 6i, \infty_7, \infty_4), \\
& (\infty_6, \frac{u}{2} + 3 + 6i, \frac{u}{2} + 4 + 6i; \frac{u}{2} + 6 + 6i, \infty_7, \infty_3),
\end{aligned}$$

for $i = 0, 1, \dots, \frac{u}{12} - 1$. \square

Lemma 2.9. Let $u \equiv 0 \pmod{12}$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 2, \frac{u}{2}\} \rangle$ can be decomposed into 3-suns.

Proof. Consider the 3-suns

$$\begin{aligned}
& (\infty_1, 6i, 1 + 6i; \frac{u}{2} + 1 + 6i, \frac{u}{2} + 6i, 3 + 6i), \\
& (\infty_1, 2 + 6i, 3 + 6i; \frac{u}{2} + 5 + 6i, \frac{u}{2} + 2 + 6i, 5 + 6i), \\
& (\infty_1, 4 + 6i, 5 + 6i; \frac{u}{2} + 2 + 6i, \frac{u}{2} + 4 + 6i, 7 + 6i), \\
& (\infty_1, \frac{u}{2} + 3 + 6i, \frac{u}{2} + 4 + 6i; \frac{u}{2} + 6i, \frac{u}{2} + 2 + 6i, \infty_2), \\
& (\infty_2, 1 + 6i, \frac{u}{2} + 1 + 6i; \frac{u}{2} + 3 + 6i, 2 + 6i, \frac{u}{2} + 2 + 6i), \\
& (\infty_2, 3 + 6i, 4 + 6i; 2 + 6i, \frac{u}{2} + 3 + 6i, 6 + 6i), \\
& (\infty_2, 5 + 6i, \frac{u}{2} + 5 + 6i; \frac{u}{2} + 2 + 6i, 6 + 6i, \frac{u}{2} + 6 + 6i), \\
& (\infty_3, 2 + 6i, 6i; 1 + 6i, 4 + 6i, \infty_2), \\
& (\infty_3, \frac{u}{2} + 2 + 6i, \frac{u}{2} + 6i; 4 + 6i, \frac{u}{2} + 4 + 6i, \infty_2), \\
& (\infty_3, \frac{u}{2} + 1 + 6i, \frac{u}{2} + 3 + 6i; 3 + 6i, \frac{u}{2} + 6i, \frac{u}{2} + 5 + 6i), \\
& (\infty_3, \frac{u}{2} + 5 + 6i, \frac{u}{2} + 4 + 6i; 5 + 6i, \frac{u}{2} + 7 + 6i, \frac{u}{2} + 6 + 6i),
\end{aligned}$$

for $i = 0, 1, \dots, \frac{u}{12} - 1$. \square

The following lemma “combines” one infinity point with one difference $d \neq \frac{u}{2}, \frac{u}{3}$ such that $\frac{u}{\gcd(u, d)} \equiv 0 \pmod{3}$ (therefore, $u \equiv 0 \pmod{3}$).

Lemma 2.10. Let $u \equiv 0 \pmod{3}$ and $d \in D_u \setminus \{\frac{u}{2}, \frac{u}{3}\}$ such that $p = \frac{u}{\gcd(u, d)} \equiv 0 \pmod{3}$. Then the graph $\langle Z_u \cup \{\infty\}, \{d\} \rangle$ can be decomposed into 3-suns.

Proof. The subgraph $\langle Z_u, \{d\} \rangle$ can be decomposed into $\frac{u}{p}$ cycles of length $p = 3q$, $q \geq 2$.

If $q > 2$, let $(x_1, x_2, \dots, x_{3q})$ be a such cycle and consider the 3-suns

$$(\infty, x_{2+3i}, x_{3+3i}; x_{7+3i}, x_{1+3i}, x_{4+3i}),$$

for $i = 0, 1, \dots, q - 1$ (where the sum is modulo $3q$).

If $q = 2$, let $(x_1^{(j)}, x_2^{(j)}, x_3^{(j)}, x_4^{(j)}, x_5^{(j)}, x_6^{(j)})$, $j = 0, 1, \dots, \frac{u}{6} - 1$, be the 6-cycles decomposing $\langle Z_u, \{d\} \rangle$ and consider the 3-suns

$$\begin{aligned}
& (\infty, x_2^{(j)}, x_3^{(j)}; x_1^{(j+1)}, x_1^{(j)}, x_4^{(j)}), \\
& (\infty, x_5^{(j)}, x_6^{(j)}; x_4^{(j+1)}, x_4^{(j)}, x_1^{(j)}),
\end{aligned}$$

for $j = 0, 1, \dots, \frac{u}{6} - 1$ (where the sums are modulo $\frac{u}{6}$). \square

Subsequent Lemmas 2.11–2.14 allow to decompose $\langle Z_u \cup H, D \rangle$, where $|H| = 1, 2, 3, 5$, $|D| = 6 - |H|$ and $\frac{u}{2} \notin D$; here, u and D are any with the unique condition that if D contains at least three differences d_1, d_2, d_3 , then $d_3 = d_2 - d_1$ or $d_1 + d_2 + d_3 = u$.

Lemma 2.11. *Let $d_1, d_2, d_3, d_4, d_5 \in D_u \setminus \{\frac{u}{2}\}$ such that $d_3 = d_2 - d_1$ or $d_1 + d_2 + d_3 = u$. Then the graph $\langle Z_u \cup \{\infty\}, \{d_1, d_2, d_3, d_4, d_5\} \rangle$ can be decomposed into 3-suns.*

Proof. If $d_3 = d_2 - d_1$, consider the orbit of

$$(d_1, d_2, 0; \infty, d_2 + d_5, d_4)$$

(or $(d_1, d_2, 0; \infty, d_2 + d_5, -d_4)$, if $d_2 + d_5 = d_4$) under Z_u . If $d_1 + d_2 + d_3 = u$, consider the orbit of

$$(-d_1, d_2, 0; \infty, d_2 + d_5, d_4)$$

(or $(-d_1, d_2, 0; \infty, d_2 + d_5, -d_4)$, if $d_2 + d_5 = d_4$) under Z_u . \square

Lemma 2.12. *Let $d_1, d_2, d_3, d_4 \in D_u \setminus \{\frac{u}{2}\}$ such that $d_3 = d_2 - d_1$ or $d_1 + d_2 + d_3 = u$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2\}, \{d_1, d_2, d_3, d_4\} \rangle$ can be decomposed into 3-suns.*

Proof. Consider the orbit of $(d_1, d_2, 0; \infty_1, \infty_2, d_4)$ or $(-d_1, d_2, 0; \infty_1, \infty_2, d_4)$ under Z_u when, respectively, $d_3 = d_2 - d_1$ or $d_1 + d_2 + d_3 = u$. \square

Lemma 2.13. *Let $d_1, d_2, d_3 \in D_u \setminus \{\frac{u}{2}\}$ such that $d_3 = d_2 - d_1$ or $d_1 + d_2 + d_3 = u$. Then the graph $\langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{d_1, d_2, d_3\} \rangle$ can be decomposed into 3-suns.*

Proof. Consider the orbit of $(d_1, d_2, 0; \infty_1, \infty_2, \infty_3)$ or $(-d_1, d_2, 0; \infty_1, \infty_2, \infty_3)$ under Z_u when, respectively, $d_3 = d_2 - d_1$ or $d_1 + d_2 + d_3 = u$. \square

Lemma 2.14. *Let $d \in D_u \setminus \{\frac{u}{2}\}$, the graph $\langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4, \infty_5\}, \{d\} \rangle$ can be decomposed into 3-suns.*

Proof. The subgraph $\langle Z_u, \{d\} \rangle$ is regular of degree 2 and so can be decomposed into l -cycles, $l \geq 3$. Let (x_1, x_2, \dots, x_l) be a such cycle. Put $l = 3q + r$, with $r = 0, 1, 2$, and consider the 3-suns with the sums modulo l

$$\begin{aligned} & (\infty_1, x_{1+3i}, x_{2+3i}; x_{3+3i}, \infty_4, \infty_5), \\ & (\infty_2, x_{2+3i}, x_{3+3i}; x_{4+3i}, \infty_4, \infty_5), \\ & (\infty_3, x_{3+3i}, x_{4+3i}; x_{5+3i}, \infty_4, \infty_5), \end{aligned}$$

for $i = 0, 1, \dots, q-2$, $q \geq 2$, plus the following blocks as the case may be.

If $r = 0$,

$$\begin{aligned} & (\infty_1, x_{3q-2}, x_{3q-1}; x_{3q}, \infty_4, \infty_5), \\ & (\infty_2, x_{3q-1}, x_{3q}; x_1, \infty_4, \infty_5), \\ & (\infty_3, x_{3q}, x_1; x_2, \infty_4, \infty_5). \end{aligned}$$

If $r = 1$,

$$\begin{aligned} & (\infty_1, x_{3q-2}, x_{3q-1}; x_{3q+1}, \infty_4, \infty_5), \\ & (\infty_2, x_{3q-1}, x_{3q}; x_1, \infty_4, \infty_1), \\ & (\infty_3, x_{3q}, x_{3q+1}; x_2, \infty_4, \infty_2), \\ & (\infty_5, x_{3q+1}, x_1; x_{3q}, \infty_4, \infty_3). \end{aligned}$$

If $r = 2$,

$$\begin{aligned} & (\infty_1, x_{3q-2}, x_{3q-1}; x_{3q+2}, \infty_4, \infty_5), \\ & (\infty_2, x_{3q-1}, x_{3q}; x_1, \infty_4, \infty_5), \\ & (\infty_3, x_{3q}, x_{3q+1}; x_2, \infty_1, \infty_2), \\ & (\infty_4, x_{3q+1}, x_{3q+2}; x_{3q}, \infty_1, \infty_3), \\ & (\infty_5, x_{3q+2}, x_1; x_{3q+1}, \infty_2, \infty_3). \end{aligned}$$

□

Finally, after settling the infinity points by using the above lemmas, if u is large we need to decompose the subgraph $\langle Z_u, L \rangle$, where L is the set of the differences unused (*difference leave*). Since by applying Lemmas 2.1 – 2.13 it could be necessary to use the differences 1, 2 or 4, while Lemma 2.14 does not impose any restriction, it is possible to combine infinity points and differences in such a way that the difference leave L is the set of the “small” differences, where 1, 2 or 4 could possibly be avoided.

Lemma 2.15. *Let $\alpha \in \{0, 4, 8\}$ and u, s be positive integers such that $u > 12s + \alpha$. Then there exists a decomposition of $\langle Z_u, L \rangle$ into 3-suns, where:*

- i) $\alpha = 0$ and $L = [1, 6s]$;
- ii) $\alpha = 4$ and $L = [3, 6s + 2]$;
- iii) $\alpha = 8$ and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$.

Proof.

- i) Consider the orbits (S_j) under Z_u , where $S_j = (5s + 1 + j, 5s - j, 0; 3s, s, u - 2 - 2j)$, $j = 0, 1, \dots, s - 1$.
- ii) Consider the orbits in i), where (S_0) is replaced with the orbit of $(6s + 1, 4s, 0; s, 9s, 6s + 2)$.
- iii) Consider the orbits in i), where the orbits (S_0) and (S_1) are replaced with the orbits of $(6s + 1, 4s, 0; s, 9s, 6s + 4)$ and $(5s + 2, 5s - 1, 0; 3s, s, 6s + 2)$. □

3 The main result

Let (X, \mathcal{S}) be a 3SS(n) and $m \equiv 0, 1, 4, 9 \pmod{12}$.

Lemma 3.1. *If (X, \mathcal{S}) is embedded in a 3-sun system of order $m > n$, then $m \geq \frac{7}{5}n + 1$.*

Proof. Suppose (X, \mathcal{S}) is embedded in (X', \mathcal{S}') , with $|X'| = m = n + u$ (u positive integer). Let c_i be the number of 3-suns of \mathcal{S}' each of which contains exactly i edges in $X' \setminus X$. Then $\sum_{i=1}^6 i \times c_i = \binom{u}{2}$ and $\sum_{i=1}^5 (6 - i)c_i = u \times n$, from which it follows $6c_2 + 12c_3 + 18c_4 + 24c_5 + 30c_6 = \frac{u(5u-2n-5)}{2}$ and so $u \geq \frac{2}{5}n + 1$ and $m \geq \frac{7}{5}n + 1$. □

By previous Lemma:

1. if $n = 60k + 5r$, $r = 0, 5, 8, 9$, then $m \geq 84k + 7r + 1$;
2. if $n = 60k + 5r + 1$, $r = 0, 3, 4, 7$, then $m \geq 84k + 7r + 3$;
3. if $n = 60k + 5r + 2$, $r = 2, 7, 10, 11$, then $m \geq 84k + 7r + 4$;
4. if $n = 60k + 5r + 3$, $r = 2, 5, 6, 9$, then $m \geq 84k + 7r + 6$;

5. if $n = 60k + 5r + 4$, $r = 0, 1, 4, 9$, then $m \geq 84k + 7r + 7$.

In order to prove that the necessary conditions for embedding a 3-sun system (X, \mathcal{S}) of order n in a 3-sun system of order $m = n + u$, $u > 0$ are also sufficient, the graph $\langle Z_u \cup X, D_u \rangle$ will be expressed as a union of edge-disjoint subgraphs $\langle Z_u \cup X, D_u \rangle = \langle Z_u \cup X, D \rangle \cup \langle Z_u, L \rangle$, where $L = D_u \setminus D$ is the difference leave, and $\langle Z_u \cup X, D \rangle$ (if necessary, expressed itself as a union of subgraphs) will be decomposed by using Lemmas 2.1–2.14, while if $L \neq \emptyset$, $\langle Z_u, L \rangle$ will be decomposed by Lemma 2.15. To obtain our main result we will distinguish the five cases 1.–5. listed before by giving a general proof for any $k \geq 0$ with the exception of a few cases for $k = 0$, which will be indicated by a star $*$ and solved in Appendix. Finally, note that:

- a) $u \equiv 0, 1, 4$, or $9 \pmod{12}$, if $n \equiv 0 \pmod{12}$;
- b) $u \equiv 0, 3, 8$, or $11 \pmod{12}$, if $n \equiv 1 \pmod{12}$;
- c) $u \equiv 0, 5, 8$, or $9 \pmod{12}$, $n \equiv 4 \pmod{12}$;
- d) $u \equiv 0, 3, 4$, or $7 \pmod{12}$, if $n \equiv 9 \pmod{12}$.

Proposition 3.2. *For any $n = 60k + 5r$, $r = 0, 5, 8, 9$, there exists a decomposition of $K_{n+u} \setminus K_n$ into 3-suns for every admissible $u \geq 24k + 2r + 1$.*

Proof. Let $X = \{\infty_1, \infty_2, \dots, \infty_{60k+5r}\}$, $r = 0, 5, 8, 9$, and $u = 24k + 2r + 1 + h$, with $h \geq 0$. Set $h = 12s + l$, $0 \leq l \leq 11$ (l depends on r), and distinguish the following cases.

Case 1: $r = 0, 5, 8, 9$ and $l = 0$ (odd u).

Write $\langle Z_u \cup X, D_u \rangle = \langle Z_u \cup X, D \rangle \cup \langle Z_u, L \rangle$, where $D = [6s + 1, 12k + r + 6s]$, $|D| = 12k + r$, and $L = [1, 6s]$, and apply Lemmas 2.14 and 2.15.

Case 2: $r = 0, 9$ and $l = 8$ (odd u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{2, 6s + 3, 6s + 5\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4\}, \{1\} \rangle \cup \langle Z_u \cup \{\infty_5\}, \{6s + 4\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4, \infty_5\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 6, 12k + r + 6s + 4]$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.13, 2.10, 2.14 and 2.15.

Case 3: $r = 5, 8$ and $l = 4$ (odd u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{2, 4\} \rangle \cup \langle Z_u \cup \{\infty_5\}, \{1\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4, \infty_5\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 3, 12k + r + 6s + 2] \setminus \{6s + 4\}$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.3, 2.10, 2.14 and 2.15.

Case 4: $r = 0, 8$ and $l = 3$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \langle Z_u \cup \{\infty_4, \infty_5\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4, \infty_5\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 3, 12k + r + 6s + 1]$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.5, 2.1, 2.14 and 2.15.

Case 5: $r = 0$ and $l = 11$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 2, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5\}, \{4, 6s + 3, 6s + 5, 6s + 7\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4, \infty_5\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 6, 12k + 6s + 5] \setminus \{6s + 7\}$, $|D'| = 12k - 1$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.9, 2.12, 2.14 and 2.15.

Case 6: $r = 5$ and $l = 1$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_7, \infty_8, \infty_9, \infty_{10}\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_{10}\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 3, 12k + 6s + 5]$, $|D'| = 12k + 3$, and $L = [3, 6s + 2]$, and apply Lemmas 2.7, 2.2, 2.14 and 2.15.

Case 7: $r = 5, 9$ and $l = 9$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5\}, \{2, 6s + 3, 6s + 4, 6s + 5\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4, \infty_5\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 6, 12k + r + 6s + 4]$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.5, 2.12, 2.14 and 2.15.

Case 8: $r = 8$ and $l = 7$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 2, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4\}, \{4\} \rangle \cup \langle Z_u \cup \{\infty_5\}, \{6s + 5\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4, \infty_5\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 3, 12k + 6s + 11] \setminus \{6s + 4, 6s + 5\}$, $|D'| = 12k + 7$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.9, 2.10, 2.14 and 2.15.

Case 9: $r = 9$ and $l = 5$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \langle Z_u \cup \{\infty_4\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_5\}, \{4\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4, \infty_5\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 3, 12k + 6s + 11] \setminus \{6s + 4\}$, $|D'| = 12k + 8$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.5, 2.10, 2.14 and 2.15. \square

Proposition 3.3. For any $n = 60k + 5r + 1$, $r = 0, 3, 4, 7$, there exists a decomposition of $K_{n+u} \setminus K_n$ into 3-suns for every admissible $u \geq 24k + 2r + 2$.

Proof. Let $X = \{\infty_1, \infty_2, \dots, \infty_{60k+5r+1}\}$, $r = 0, 3, 4, 7$, and $u = 24k + 2r + 2 + h$, with $h \geq 0$. Set $h = 12s + l$, $0 \leq l \leq 11$, and distinguish the following cases.

Case 1: $r = 0, 3$ and $l = 1$ (odd u).

Write $\langle Z_u \cup X, D_u \rangle = \langle Z_u \cup \{\infty\}, \{6s + 2\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty\}), D' \rangle \cup \langle Z_u, L \rangle$, where $D' = [6s + 1, 12k + r + 6s + 1] \setminus \{6s + 2\}$, $|D'| = 12k + r$, and $L = [1, 6s]$, and apply Lemmas 2.10, 2.14 and 2.15.

Case 2: $r = 0, 3, 4, 7$ and $l = 9$ (odd u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 6s + 3, 6s + 4\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5, \infty_6\}, \{2, 6s + 5, 6s + 7\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_6\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 6, 12k + r + 6s + 5] \setminus \{6s + 7\}$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.13, 2.14 and 2.15.

Case 3: $r = 4^*, 7$ and $l = 5$ (odd u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{2, 4\} \rangle \cup \langle Z_u \cup \{\infty_5\}, \{1\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_6\}, \{6s + 8\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_6\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 3, 12k + r + 6s + 3] \setminus \{6s + 4, 6s + 8\}$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.3, 2.10, 2.14 and 2.15.

Case 4: $r = 0, 4$ and $l = 6$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5, \infty_6\}, \{2, 6s + 3, 6s + 5\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_6\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 4, 12k + r + 6s + 3] \setminus \{6s + 5\}$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.5, 2.13, 2.14 and 2.15.

Case 5: $r = 0$ and $l = 10$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_7, \infty_8, \infty_9, \infty_{10}\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_{11}\}, \{4, 6s + 3, 6s + 5, 6s + 6, 6s + 7\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_{11}\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 8, 12k + 6s + 5]$, $|D'| = 12k - 2$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.7, 2.2, 2.11, 2.14 and 2.15.

Case 6: $r = 3, 7$ and $l = 0$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_6\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = \{2\} \cup [6s + 3, 12k + r + 6s]$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.7, 2.14 and 2.15.

Case 7: $r = 3$ and $l = 4$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{1, \frac{u}{2}\} \rangle \cup \langle Z_u \cup \{\infty_5\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_6\}, \{6s + 5\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_6\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s + 3, 12k + 6s + 5] \setminus \{6s + 5\}$, $|D'| = 12k + 2$, and $L = [3, 6s + 2]$, and apply Lemmas 2.6, 2.10, 2.14 and 2.15.

Case 8: $r = 4$ and $l = 2$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{1, \frac{u}{2}\} \rangle \cup \langle Z_u \cup \{\infty_5, \infty_6\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_6\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s + 3, 12k + 6s + 5]$, $|D'| = 12k + 3$, and $L = [3, 6s + 2]$, and apply Lemmas 2.6, 2.1, 2.14 and 2.15.

Case 9: $r = 7$ and $l = 8$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 2, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5, \infty_6\}, \{4, 6s + 3, 6s + 7\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_6\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s + 5, 12k + 6s + 11] \setminus \{6s + 7\}$, $|D'| = 12k + 6$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.9, 2.13, 2.14 and 2.15. \square

Proposition 3.4. *For any $n = 60k + 5r + 2$, $r = 2, 7, 10, 11$, there exists a decomposition of $K_{n+u} \setminus K_n$ into 3-suns for every admissible $u \geq 24k + 2r + 2$.*

Proof. Let $X = \{\infty_1, \infty_2, \dots, \infty_{60k+5r+2}\}$, $r = 2, 7, 10, 11$, and $u = 24k + 2r + 2 + h$, with $h \geq 0$. Set $h = 12s + l$, $0 \leq l \leq 11$, and distinguish the following cases.

Case 1: $r = 2, 11$ and $l = 3$ (odd u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1\}, \{6s + 2\} \rangle \cup \langle Z_u \cup \{\infty_2\}, \{6s + 4\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s + 1, 12k + r + 6s + 2] \setminus \{6s + 2, 6s + 4\}$, $|D'| = 12k + r$, and $L = [1, 6s]$, and apply Lemmas 2.10, 2.14 and 2.15.

Case 2: $r = 2, 7, 10, 11$ and $l = 7$ (odd u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2\}, \{1, 2, 6s + 3, 6s + 4\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 5, 12k + r + 6s + 4]$, $|D'| = 12k + r$, and $L = [3, 6s + 2]$, and apply Lemmas 2.12, 2.14 and 2.15.

Case 3: $r = 7, 10$ and $l = 11$ (odd u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 6s + 3, 6s + 4\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5, \infty_6\}, \{2, 6s + 5, 6s + 7\} \rangle \cup \langle Z_u \cup \{\infty_7\}, \{6s + 8\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_7\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 6, 12k + r + 6s + 6] \setminus \{6s + 7, 6s + 8\}$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.13, 2.10, 2.14 and 2.15.

Case 4: $r = 2$ and $l = 6$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_5, \infty_6, \infty_7\}, \{2, 6s + 3, 6s + 5\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_7\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 4, 12k + 6s + 5] \setminus \{6s + 5\}$, $|D'| = 12k + 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.6, 2.13, 2.14 and 2.15.

Case 5: $r = 2, 10$ and $l = 10$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_7\}, \{2, 6s + 3, 6s + 4, 6s + 5, 6s + 6\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_7\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 7, 12k + r + 6s + 5]$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.7, 2.11, 2.14 and 2.15.

Case 6: $r = 7, 11$ and $l = 4$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5, \infty_6, \infty_7\}, \{2, 4\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_7\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 3, 12k + r + 6s + 2] \setminus \{6s + 4\}$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.5, 2.3, 2.14 and 2.15.

Case 7: $r = 7$ and $l = 8$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5, \infty_6\}, \{2, 6s + 3, 6s + 5\} \rangle \cup \langle Z_u \cup \{\infty_7\}, \{6s + 7\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_7\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 4, 12k + 6s + 11] \setminus \{6s + 5, 6s + 7\}$, $|D'| = 12k + 6$, and $L = [3, 6s + 2]$, and apply Lemmas 2.5, 2.13, 2.10, 2.14 and 2.15.

Case 8: $r = 10$ and $l = 2$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, \frac{u}{2}\} \rangle \cup \langle Z_u \cup \{\infty_7\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_7\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 3, 12k + 6s + 11]$, $|D'| = 12k + 9$, and $L = [3, 6s + 2]$, and apply Lemmas 2.7, 2.10, 2.14 and 2.15.

Case 9: $r = 11$ and $l = 0$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_7\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_7\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = \{2\} \cup [6s + 3, 12k + 6s + 11]$, $|D'| = 12k + 10$, and $L = [3, 6s + 2]$, and apply Lemmas 2.8, 2.14 and 2.15. \square

Proposition 3.5. *For any $n = 60k + 5r + 3$, $r = 2, 5, 6, 9$, there exists a decomposition of $K_{n+u} \setminus K_n$ into 3-suns for every admissible $u \geq 24k + 2r + 3$.*

Proof. Let $X = \{\infty_1, \infty_2, \dots, \infty_{60k+5r+3}\}$, $r = 2, 5, 6, 9$, and $u = 24k + 2r + 3 + h$, with $h \geq 0$. Set $h = 12s + l$, $0 \leq l \leq 11$, and distinguish the following cases.

Case 1: $r = 2, 5, 6, 9$ and $l = 4$ (odd u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 6s + 3, 6s + 4\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = \{2\} \cup [6s + 5, 12k + r + 6s + 3]$, $|D'| = 12k + r$, and $L = [3, 6s + 2]$, and apply Lemmas 2.13, 2.14 and 2.15.

Case 2: $r = 2, 5$ and $l = 8$ (odd u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2\}, \{1, 6s + 3, 6s + 4, 6s + 5\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_3\}, \{2\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 6, 12k + r + 6s + 5]$, $|D'| = 12k + r$, and $L = [3, 6s + 2]$, and apply Lemmas 2.12, 2.10, 2.14 and 2.15.

Case 3: $r = 6, 9$ and $l = 0$ (odd u).

If $s = 0$, then write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_8\}, \{1, \frac{u}{3}\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_8\}), D' \rangle, \end{aligned}$$

where $D' = [2, 12k + r + 1] \setminus \{\frac{u}{3}\}$, $|D'| = 12k + r - 1$, and apply Lemmas 2.4 and 2.14.

If $s > 0$, then write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 5s, 5s + 1\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5, \infty_6\}, \{2, 6s + 1, 6s + 3\} \rangle \cup \langle Z_u \cup \{\infty_7\}, \{6s + 2\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_8\}, \{6s + 4\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_8\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = \{2s + 1, 4s\} \cup [6s + 5, 12k + r + 6s + 1]$, $|D'| = 12k + r - 1$, and $L = [3, 6s] \setminus \{2s + 1, 4s, 5s, 5s + 1\}$, and apply Lemmas 2.13, 2.10 and 2.14 to decompose the first five subgraphs, while to decompose the last one apply Lemma 2.15 i) and delete the orbit (S_0).

Case 4: $r = 2, 6$ and $l = 1$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = \{2\} \cup [6s + 3, 12k + r + 6s + 1]$, $|D'| = 12k + r$, and $L = [3, 6s + 2]$, and apply Lemmas 2.5, 2.14 and 2.15.

Case 5: $r = 2^*$ and $l = 5$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_7, \infty_8, \infty_9, \infty_{10}\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_{11}, \infty_{12}, \infty_{13}\}, \{4, 6s + 3, 6s + 7\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_{13}\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 5, 12k + 6s + 5] \setminus \{6s + 7\}$, $|D'| = 12k$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.7, 2.2, 2.13, 2.14 and 2.15.

Case 6: $r = 5, 9$ and $l = 7$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_7, \infty_8\}, \{2, 6s + 3, 6s + 4, 6s + 5\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_8\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 6, 12k + r + 6s + 4]$, $|D'| = 12k + r - 1$, and $L = [3, 6s + 2]$, and apply Lemmas 2.7, 2.12, 2.14 and 2.15.

Case 7: $r = 5$ and $l = 11$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_5, \infty_6\}, \{2, 6s + 3, 6s + 5, 6s + 6\} \rangle \cup \langle Z_u \cup \{\infty_7\}, \{4\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_8\}, \{6s + 7\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_8\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 8, 12k + 6s + 11]$, $|D'| = 12k + 4$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.6, 2.12, 2.10, 2.14 and 2.15.

Case 8: $r = 6$ and $l = 9$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4, \infty_5, \infty_6\}, \{2, 6s + 3, 6s + 5\} \rangle \cup \langle Z_u \cup \{\infty_7\}, \{4\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_8\}, \{6s + 7\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_8\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 6, 12k + 6s + 11] \setminus \{6s + 7\}$, $|D'| = 12k + 5$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.5, 2.13, 2.10, 2.14 and 2.15.

Case 9: $r = 9$ and $l = 3$ (even u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 2, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = [6s + 3, 12k + 6s + 11]$, $|D'| = 12k + 9$, and $L = [3, 6s + 2]$, and apply Lemmas 2.9, 2.14 and 2.15. \square

Proposition 3.6. *For any $n = 60k + 5r + 4$, $r = 0, 1, 4, 9$, there exists a decomposition of $K_{n+u} \setminus K_n$ into 3-suns for every admissible $u \geq 24k + 2r + 3$.*

Proof. Let $X = \{\infty_1, \infty_2, \dots, \infty_{60k+5r+4}\}$, $r = 0, 1, 4, 9$, and $u = 24k + 2r + 3 + h$, with $h \geq 0$. Set $h = 12s + l$, $0 \leq l \leq 11$, and distinguish the following cases.

Case 1: $r = 0, 1^*, 4, 9$ and $l = 2$ (odd u).

Write

$$\begin{aligned} \langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{2, 4\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4\}), D' \rangle \cup \langle Z_u, L \rangle, \end{aligned}$$

where $D' = \{1, 6s + 3\} \cup [6s + 5, 12k + r + 6s + 2]$, $|D'| = 12k + r$, and $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.3, 2.14 and 2.15.

Case 2: $r = 0, 9$ and $l = 6$ (odd u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, 6s+3, 6s+4\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4\}, \{2\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s+5, 12k+r+6s+4]$, $|D'| = 12k+r$, and $L = [3, 6s+2]$, and apply Lemmas 2.13, 2.10, 2.14 and 2.15.

Case 3: $r = 1, 4$ and $l = 10$ (odd u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2\}, \{1, 6s+3, 6s+5, 6s+6\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_3\}, \{2\} \rangle \cup \langle Z_u \cup \{\infty_4\}, \{6s+4\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s+7, 12k+r+6s+6]$, $|D'| = 12k+r$, and $L = [3, 6s+2]$, and apply Lemmas 2.12, 2.10, 2.14 and 2.15.

Case 4: $r = 0, 4$ and $l = 5$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_7, \infty_8, \infty_9\}, \{2, 6s+3, 6s+5\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_9\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s+4, 12k+r+6s+3] \setminus \{6s+5\}$, $|D'| = 12k+r-1$, and $L = [3, 6s+2]$, and apply Lemmas 2.7, 2.13, 2.14 and 2.15.

Case 5: $r = 0$ and $l = 9$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_5, \infty_6, \infty_7\}, \{2, 6s+3, 6s+5\} \rangle \cup \langle Z_u \cup \{\infty_8\}, \{4\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_9\}, \{6s+7\} \rangle \cup \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_9\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s+6, 12k+6s+5] \setminus \{6s+7\}$, $|D'| = 12k-1$, and $L = [3, 6s+4] \setminus \{4, 6s+3\}$, and apply Lemmas 2.6, 2.13, 2.10, 2.14 and 2.15.

Case 6: $r = 1$ and $l = 7$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_7\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_8, \infty_9\}, \{2, 4, 6s+3, 6s+5\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_9\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s+6, 12k+6s+5]$, $|D'| = 12k$, and $L = [3, 6s+4] \setminus \{4, 6s+3\}$, and apply Lemmas 2.8, 2.12, 2.14 and 2.15.

Case 7: $r = 1, 9$ and $l = 11$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_4\}, \{2, 4, 6s+3, 6s+5, 6s+6\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s+7, 12k+r+6s+6]$, $|D'| = 12k+r$, and $L = [3, 6s+4] \setminus \{4, 6s+3\}$, and apply Lemmas 2.5, 2.11, 2.14 and 2.15.

Case 8: $r = 4$ and $l = 1$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{1, \frac{u}{2}\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = \{2\} \cup [6s+3, 12k+6s+5]$, $|D'| = 12k+4$, and $L = [3, 6s+2]$, and apply Lemmas 2.6, 2.14 and 2.15.

Case 9: $r = 9$ and $l = 3$ (even u).

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3\}, \{1, \frac{u}{2}\} \rangle \cup \langle Z_u \cup \{\infty_4\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \infty_3, \infty_4\}), D' \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $D' = [6s+3, 12k+6s+11]$, $|D'| = 12k+9$, and $L = [3, 6s+2]$, and apply Lemmas 2.5, 2.10, 2.14 and 2.15. \square

Combining Lemma 3.1 and Propositions 3.2 – 3.6 gives our main theorem.

Theorem 3.7. Any 3SS(n) can be embedded in a 3SS(m) if and only if

$$m \geq \frac{7}{5}n + 1 \quad \text{or} \quad m = n.$$

References

- [1] D. E. Bryant and C. A. Rodger, The Doyen-Wilson theorem extended to 5-cycles, *J. Comb. Theory Ser. A* **68** (1994), 218–225, doi:10.1016/0097-3165(94)90101-5.
- [2] D. E. Bryant and C. A. Rodger, On the Doyen-Wilson theorem for m -cycle systems, *J. Combin. Des.* **2** (1994), 253–271, doi:10.1002/jcd.3180020405.
- [3] J. Doyen and R. M. Wilson, Embeddings of Steiner triple systems, *Discrete Math.* **5** (1973), 229–239, doi:10.1016/0012-365x(73)90139-8.
- [4] C.-M. Fu, Y.-L. Lin, S.-W. Lo, Y.-F. Hsu and W.-C. Huang, The Doyen-Wilson theorem for bull designs, *Discrete Math.* **313** (2013), 498–507, doi:10.1016/j.disc.2012.11.023.
- [5] H.-L. Fu and C. C. Lindner, The Doyen-Wilson theorem for maximum packings of K_n with 4-cycles, *Discrete Math.* **183** (1998), 103–117, doi:10.1016/s0012-365x(97)00080-0.
- [6] H.-L. Fu, C. C. Lindner and C. A. Rodger, Two Doyen-Wilson theorems for maximum packings with triples, *Discrete Math.* **178** (1998), 63–71, doi:10.1016/s0012-365x(97)81817-1.

- [7] H. Gao and J. Wang, Doyen-Wilson theorem for perfect hexagon triple systems, *Discrete Math.* **311** (2011), 1006–1014, doi:10.1016/j.disc.2011.03.004.
- [8] W.-C. Huang and W.-C. Yang, The Doyen-Wilson theorem for extended directed triple systems, *Ars Combin.* **84** (2007), 77–83.
- [9] G. Lo Faro and A. Tripodi, The Doyen-Wilson theorem for kite systems, *Discrete Math.* **306** (2006), 2695–2701, doi:10.1016/j.disc.2006.03.074.
- [10] G. Lo Faro and A. Tripodi, Embeddings of λ -fold kite systems, $\lambda \geq 2$, *Australas. J. Combin.* **36** (2006), 143–150, https://ajc.maths.uq.edu.au/pdf/36/ajc_v36_p143.pdf.
- [11] R. Peltesohn, Eine Lösung der beiden Heffterschen Differenzenprobleme, *Compositio Math.* **6** (1939), 251–257, http://www.numdam.org/item?id=CM_1939__6__251_0.
- [12] G. Stern and H. Lenz, Steiner triple systems with given subspaces; another proof of the Doyen-Wilson-theorem, *Boll. Un. Mat. Ital. A Serie 5* **17** (1980), 109–114.
- [13] J. Wang, Perfect hexagon triple systems with given subsystems, *Discrete Math.* **309** (2009), 2930–2933, doi:10.1016/j.disc.2008.07.004.
- [14] J.-X. Yin and B.-S. Gong, Existence of G -designs with $|V(G)| = 6$, in: W. D. Wallis, H. Shen, W. Wei and L. Zhu (eds.), *Combinatorial Designs and Applications*, Marcel Dekker, New York, volume 126 of *Lecture Notes in Pure and Applied Mathematics*, 1990 pp. 201–218.

Appendix

- $n = 21, u = 12s + 15$

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{2, 4\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_5\}, \{1\} \rangle \cup \langle Z_u \cup \{\infty_6\}, \{6s + 7\} \rangle \cup \\ &\quad \langle Z_u \cup (X \setminus \{\infty_1, \infty_2, \dots, \infty_6\}), \{6s + 3, 6s + 5, 6s + 6\} \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.3, 2.10, 2.14 and 2.15.

- $n = 13, u = 12s + 12$

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \dots, \infty_6\}, \{1, 6s + 6\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_7, \infty_8, \infty_9, \infty_{10}\}, \{2\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_{11}, \infty_{12}, \infty_{13}\}, \{4, 6s + 3, 6s + 5\} \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $L = [3, 6s + 4] \setminus \{4, 6s + 3\}$, and apply Lemmas 2.7, 2.2, 2.13 and 2.15.

- $n = 9, u = 12s + 7$

Write

$$\begin{aligned}\langle Z_u \cup X, D_u \rangle &= \langle Z_u \cup \{\infty_1, \infty_2, \infty_3, \infty_4\}, \{2, 4\} \rangle \cup \\ &\quad \langle Z_u \cup \{\infty_5, \infty_6, \infty_7, \infty_8, \infty_9\}, \{1\} \rangle \cup \langle Z_u, L \rangle,\end{aligned}$$

where $L = [3, 6s + 3] \setminus \{4\}$, and apply Lemmas 2.3, 2.14 and decompose $\langle Z_u, L \rangle$ as in Lemma 2.15 iii), taking in account that $|6s + 4|_{12s+7} = 6s + 3$.