An overview of monitoring for raptors in Ireland

Pregled monitoringa populacij ptic roparic na Irskem

Allan Mee

Irish Raptor Study Group, Glenanaar, Ardpatrick, Kilmallock, Co. Limerick, Ireland, e-mail: allanmee@goldeneagle.ie

Ireland holds a low diversity of breeding raptors as a result of its location on the western edge of Europe but also due to historical persecution leading to the loss of at least seven species. Recolonisation by Buzzards Buteo buteo and the recent reintroduction of three species, Golden Eagle Aquila chrysaetos, Whitetailed Eagle Haliaeetus albicilla and Red Kite Milvus milvus, has helped redress such losses. Monitoring for raptors is carried out by the statutory agencies, NGOs and two university research groups. Decadal and semi-decadal surveys are undertaken nationally in the Republic of Ireland and Northern Ireland for Peregrine Falcon Falco peregrinus and Hen Harriers Circus cyaneus, respectively. Long term monitoring projects have been established for some key species such as Barn Owl Tyto alba. However, some species receive little monitoring effort (e.g. Sparrowhawk Accipiter nisus, Merlin F. columbarius). A dedicated Raptor Monitoring Scheme to determine long-term population trends across a range of species is lacking and remains an urgent priority. Development of an Action Plan for raptors and/or single key species would further help identify priorities and raise awareness of the need of monitoring for raptors.

Key words: birds of prey, owls, monitoring, survey, Ireland **Ključne besede**: ujede, sove, monitoring, popis, Irska

1. Introduction

Ireland holds perhaps the lowest diversity of raptor species in Europe (Table 1) in part as a result of its location as an island on the western edge of Europe but also due to the loss of several species as a result of centuries of human persecution and habitat loss (D'ARCY 1999). However, in the last few decades four species previously extinct have now either been re-established through reintroduction programmes (O'TOOLE *et al.* 2002) or have recolonised naturally (NORRISS 1991), while two other species have bred intermittently and may re-establish themselves in the near future (HILLIS 2008).

Perhaps due to different conservation priorities and the low diversity of breeding raptors, monitoring of raptors in Ireland has received much less attention than other bird groups (e.g., seabirds, wildfowl) until recently (see Table 1, Figure 1).

2. Main players

Statutory responsibility for raptor monitoring and conservation rests with the National Parks and Wildlife Service (NPWS) in the Republic of Ireland (RoI) and the Northern Ireland Environment Agency (NIEA) in Northern Ireland. Such responsibilities include the designation of Special Protection Areas (SPA) for Birds Directive Annex I raptors within the Natura 2000 network, the management of six National Parks in the RoI, some of which hold important raptor populations, and the enforcement of the Wildlife Act (1976, 2000) to protect raptor populations (see www. npws.ie).

Two NGOs have a specific remit to monitor raptor populations. The Irish Raptor Study Group (IRSG) is a voluntary organisation solely dedicated to monitor and improve the conservation status of raptors in the RoI (IRSG 2006). The Golden Eagle Trust

Table 1: Summary of monitoring programmes for raptors in Ireland (including Northern Ireland)

Tabela 1: Povzetek programov monitoringa ptic roparic na Irskem (vključno s Severno Irsko)

Species / Vrsta	Population Estimate/ Populacijska ocena	Project name/ Naziv projekta	Project type/ Vrsta projekta	Duration/ Trajanje	Organisation/ Organizacija
Red Kite Milvus milvus	25–30	Irish Red Kite reintroduction ¹	Reintroduction	2007–2011	GET, NPWS, RSPB
White-tailed Eagle Haliaeetus albicilla	10	Irish White- tailed Eagle reintroduction ²	Reintroduction	2007–2011	GET, NPWS
Marsh Harrier Circus aeruginosus	0-2	No studies			IRSG
Hen Harrier Circus cyaneus	190–220	Planforbio ³	Research		University College Cork
Goshawk Accipiter gentilis	0-5	No studies			D. Scott (in litt.)
Sparrowhawk Accipiter nisus	11,000	No studies			IRSG
Buzzard Buteo buteo	3,500-4,000	Buzzard Ecology & Biology Project ⁴	PhD/ Monitoring	2010-	Queens University, IRSG
Golden Eagle Aquila chrysaetos	5–10	Irish Golden Eagle Reintroduction ⁵	Reintroduction	2001-	GET, NPWS
Kestrel Falco tinnunculus	10,000	Kestrel Monitoring Project ⁶	Monitoring/ Research		BWI
Merlin Falco columbarius	250+	Merlin pilot project ⁷	Research	2010-	BWI, NPWS
Peregrine Falcon Falco peregrinus	450-500	Wicklow/NW Ireland pop. Study ⁸	Survey/ Research	2011	D. Clarke/ NPWS/IRSG
Barn Owl <i>Tyto alba</i>	400-600	Barn Owl Research Project ⁹	Research	2006–	BWI
Snowy Owl Bubo scandiacus	0-1	No studies			IRSG
Long-eared Owl Asio otus	3,500+	LEO Monitoring Project ¹⁰	Survey/ Monitoring	2012—	BWI, NPWS
Short-eared Owl Asio flammeus	0-5	No studies			

¹Co. Wicklow, Co. Down; ²Co. Kerry; ³Cos. Kerry, Cork, Limerick, Clare; ⁴Co. Cork (IRSG), N. Ireland (Queens); ⁵Co. Donegal; ⁶SW Ireland; ⁷Wicklow, Donegal, Connemara; ⁸Co. Wicklow, NW Ireland; ⁹SW Ireland; ¹⁰Connemara, SW Ireland.

(GET) works to restore populations of threatened or previously extinct raptor species. The GET manages reintroduction programmes for Golden Eagles *Aquila chrysaetos*, White-tailed Eagles *Haliaeetus albicilla*, and Red Kites *Milvus milvus* in the RoI in collaboration with NPWS (O'TOOLE 2002, MEE 2009 & 2010). A third NGO, BirdWatch Ireland (BWI), the BirdLife

International partner in the RoI, is dedicated to the conservation of Irish birds and has recently established a Raptor Conservation Project. BWI established a long term monitoring programme for Barn Owls *Tyto alba* in 2006 and has more recently initiated research and monitoring of Long-eared Owl *Asio otus*, Kestrel *Falco tinnunculus* and Merlin *F. columbarius* to determine

 $GET-Golden\ Eagle\ Trust;\ NPWS-National\ Parks\ and\ Wildlife\ Service;\ IRSG-Irish\ Raptor\ Study\ Group;\ BWI-BirdWatch\ Ireland\ Study\ Group;\ BWI-BirdWatch\ Ireland\ Group;\ BWI-Bird$

densities, trends, ecological requirements and factors affecting the conservation status of these populations (Lusby 2009, 2012A & 2012B, Lusby et al. 2010A, 2010B, 2011, 2012A & 2012B, FERNÁNDEZ-BELLON & Lusby 2011A & 2011B, O'CLEARY et al. 2012). In Northern Ireland, the Royal Society for the Protection of Birds (RSPB) manages the reintroduction programme for Red Kites in County Down. The Northern Ireland Raptor Study Group (NIRSG) is the sister group of the IRSG, monitoring raptors in Northern Ireland, especially an intensive monitoring programme for Peregrine Falcons F. peregrinus (Wells 2007).

Two university research teams monitor raptor populations. The Planforbio research team at University College Cork has been working on the ecology of Hen Harrier *Circus cyaneus* nesting in commercial forestry plantations since 2008, work which has informed the State forestry body, Coillte, of forest management possibilities for harriers (IRWIN *et al.* 2008, WILSON *et al.* 2009 & 2012). The Quercus programme at Queens University, Belfast, Northern Ireland, has instigated studies of harriers and Peregrine Falcon (RUDDOCK 2007 & 2008), as well as raptor monitoring workshops (www.qub.ac.uk/sites/Quercus). A study of the population dynamics of Buzzards *Buteo buteo* is also ongoing (ROONEY 2013, ROONEY & MONTGOMERIE 2013).

Collaboration in monitoring for raptors is primarily with the UK including breeding and winter Atlas studies (all species including raptors), which are run simultaneously across Ireland and Britain (Sharrock 1976, Lack 1986, Gibbons et al. 1993). Single species studies of Peregrine Falcon (Crick & Ratcliffe 1995, Norriss 1995, Banks et al. 2003, Madden et al. 2009) and Hen Harrier (Sim et al. 2001, Norriss et al. 2002, Barton et al. 2006, Sim et al. 2007, Ruddock et al. 2010) have also been carried out in both regions in similar time periods. There is also close collaboration with Britain on ringing studies and activities as Irish ringers use rings supplied by the British Trust for Ornithology (BTO) and supply raptor information to BTO studies such as the Nest Record Scheme.

Raptor monitoring data submitted to statutory agencies are important in determining population status and trends and may form the basis for conservation action such as the SPA designation. Declining species may be added to the Red List of species of conservation concern and become priorities for conservation (Lynas et al. 2007). NGOs, such as the IRSG, NIRSG, GET and BWI, supply much of the data needed for setting conservation priorities. State agencies such as Coillte may use data on Hen Harrier to inform forest management.




Figure 1: Monitoring programmes for raptors in Ireland. Numbers refer to project names in Table 1.

Şlika 1: Programi monitoringa ptic roparic na Irskem. Številke ustrezajo nazivom projektov v Tabeli 1.

3. National coverage

Apart from Atlas studies (all species), coverage for single species raptor surveys has been coordinated by NPWS/NIEA, NGOs or private consultants working on behalf of the statutory agencies so that populations in all regions are assessed (e.g. MADDEN et al. 2009, RUDDOCK et al. 2010). NGOs have also coordinated their own national surveys to determine population status and trends (Lusby 2012B, MEE & CLARKE in prep.). No national scheme exists for long-term monitoring of raptor population trends, although this has been proposed by the IRSG-NIRSG to the statutory agencies. At present, little or no funding appears to be available for this programme. The IRSG-NIRSG provide a framework for coordinating raptor monitoring but, except for dedicated national surveys, coverage is patchy and tends to focus on species of interest to individuals or those that can be relatively easily accessed without specialist skills such as tree-climbing, while some of the more widespread and apparently common raptors (e.g. Sparrowhawk Accipiter nisus) are little monitored. Coverage also tends to be lowest in more remote, mountain areas.

4. Key species and key issues

The focus of monitoring for raptors to date has been national and regional studies of Annex I species, particularly Peregrine Falcon (Moore et al. 1992 & 1997, Wells 2007) and Hen Harrier (Scott 2000, O'DONOGHUE 2010) as well as local and regional studies of Merlin (CLARKE & SCOTT 1994, McElheron 2005, Norriss et al. 2010, Fernández-BELLON & LUSBY 2011A & 2011B). Other key species have been those recently listed on the Red (high) and Amber (medium) list of birds of conservation concern (Lynas et al. 2007) including the recently reintroduced Golden Eagle and Red Kite. Breeding populations of the reintroduced Golden Eagle (northwestern Ireland) and Red Kite (County Wicklow) are comprehensively monitored by the GET using radio and satellite telemetry as well as public sighting of wing-tagged individuals (see www.goldeneagle.ie). White-tailed Eagles are similarly monitored, with first breeding in the wild expected in 2012/2013 (MEE 2009 & 2010).

Although historically habitat loss, principally loss of native forest cover and wetland drainage, has had an undoubted effect of Irish raptor populations, direct human persecution has in the past caused most population extinctions including the loss of breeding Osprey, Golden Eagle *Pandion haliaetus*, White-tailed Eagle, Red Kite, Goshawk *A. gentilis* and Buzzard (D'ARCY 1999). All but Osprey have now either been reintroduced or recolonised to varying extents (NORRISS 1991, HILLIS 2008, MEE 2010).

However, poisoning remains the greatest threat to re-establishing or maintaining populations. Between 2007 and early 2013, of 27 White-tailed Eagles recovered dead, 12 were confirmed poisoned in RoI and another seven suspected (GET unpubl.). At least three Golden Eagles and 16 Red Kites have also been poisoned during this period. Most if not all such poisoning appears to be accidental, foxes and crows being the intended target. However, several Buzzards and even Peregrine Falcons have also been deliberately poisoned. Most recently (Sept-Dec 2011), eight Red Kites were recovered dead, at least six apparently due to rodenticide (brodifacoum) poisoning (GET unpubl.) to which Barn Owls in Ireland are highly susceptible (J. Lusby unpubl.). Following a formal complaint to the EU (GET 2009), the use of poisons for the control of foxes has been banned in the RoI (Oct 2010). However, illegal use and misuse of substances

remains a problem. A formal protocol for postmortem handling, toxicological analysis and reporting was established in 2011 by NPWS in collaboration with the Department of Agriculture and the State Laboratory. Rodenticides also kill Kestrel and Buzzard, but population level threats are unknown. Other threats include road (Barn Owl) and rail (Red Kite) collisions, e.g. 214 road mortalities in 2006–2012 (J. Lusby pers. comm.). The recent expansion of wind farm developments, including some within SPAs, is also likely to pose a threat through displacement and/or collision (Scott & McHaffie 2008, Pearce-Higgins et al. 2009). In 2011, two White-tailed Eagles were recovered dead due to turbine collisions in south-western Ireland (GET unpubl.).

No currently breeding Irish raptors are truly migratory, although there is evidence that some Hen Harriers winter as far as Western Europe (B. O'Donoghue *in litt.*). Sharing information on dispersal patterns might improve the conservation status of this species. Issues facing raptors in Ireland such as poisoning and windfarm collisions would benefit from international networking to establish best practice for monitoring and assessing threats to populations.

5. Strengths and weaknesses

Ireland is a relatively small country with a smaller suite of breeding raptors than most other European countries. Therefore it should be in a position to devise and implement a comprehensive monitoring programme. Decadal and semi-decadal national surveys for key species such as Peregrine Falcon and Hen Harrier have worked well because the populations have been small enough, nesting sites reasonably well known, or populations have been discrete (e.g. MADDEN et al. 2009, RUDDOCK et al. 2010). However, such surveys are labour intensive and as some populations recover and expand (e.g. Peregrine Falcon) random selection of study areas will be essential. Other key species (e.g. Merlin) have never been surveyed on a national scale, while baseline data do not exist for some very rare breeders (e.g. Short-eared Owl Asio flammeus). Barn Owls have been increasingly monitored in recent years (NAGLE 2007, O'CONNELL et al. 2007, LUSBY 2009, 2012A & 2012B, LUSBY et al. 2010A & 2010B), but little specific monitoring of breeding Long-eared or Short-eared Owl populations or their ecology has been undertaken (but see and SLEEMAN 1988, SMAL 1989, ANDREWS 1992, COOKE et al. 1995).

Lack of a Raptor Monitoring Scheme (RMS) to

determine long-term population trends across a range of species, including some widespread but possibly declining species such as Kestrel, remain a major weakness. International best practice and information sharing on appropriate monitoring schemes would be useful in designing a RMS.

6. Priorities, capacity-building

Establishing an island-wide RMS is a high priority. However, lack of funding and the small pool of existing raptor fieldworkers will inhibit development of this scheme without full time personnel to drive the RMS.

Semi-decadal surveys for Hen Harriers should be continued. Merlin needs to be monitored at an appropriate level on at least a semi-decadal frequency. High priority reintroduced raptors should continue to be intensively monitored at least until all populations are well established. Rare and little known species should also be targeted to establish baseline data for conservation. Ultimately funding will be needed to drive much of this effort. Where key skills are lacking, training workshops will be useful in upskilling fieldworkers and generating support for monitoring programmes.

Group Action Plans have been devised for suites of species including raptors based on ecosystem to identify specific conservation requirements, targets and the actions needed to achieve such targets (e.g. BIRDWATCH IRELAND 2010). Development of an Action Plan for raptors or single key species would further help identify priorities and raise awareness of the need of monitoring for raptors.

Further, the timely publication and dissemination of monitoring results is important not only for improving our knowledge of species and setting priorities but would also further capacity-building for raptors.

Acknowledgements: Thanks to the dedicated raptor fieldworkers who provided information on their monitoring and/or research projects including as yet unpublished data. Special thanks to Damian Clarke (NPWS), John Lusby (BWI), Tony Nagle (IRSG), and Eimear Rooney (Queens University). John Lusby made helpful comments on the manuscript.

7. Povzetek

Na Irskem je pestrost ptic roparic majhna, a ne zgolj zaradi geografske lege te države na zahodnem robu Evrope, marveč tudi zaradi zgodovinskega preganjanja

teh ptic na otoku in posledično izumrtja najmanj sedmih vrst. Te izgube se je Irski posrečilo do neke mere nadomestiti z nedavno ponovno naselitvijo treh vrst, in sicer planinskega orla Aquila chrysaetos, belorepca Haliaeetus albicilla in rjavega škarnika Milvus milvus, medtem ko je kanja Buteo buteo pričela ponovno gnezditi sama. Za monitoring ptic roparic skrbijo različne javne agencije, nevladne organizacije in dve univerzitetni raziskovalni skupini. V Republiki Irski poteka, na primer, nacionalni popis sokola selca Falco peregrinus vsakih deset let, na Severnem Irskem pa popis pepelastega lunja Circus cyaneus vsakih pet let. Za nekatere ključne vrste, kakršna je pegasta sova *Tyto alba*, so bili izdelani projekti za njihov dolgoročni monitoring. Nekatere druge vrste, npr. skobec Accipiter nisus in mali sokol F. columbarius, pa so po drugi strani deležne komaj omembe vredne pozornosti. Najnujnejša prioriteta ostaja program temeljitega monitoringa ptic roparic, s katerim bi ugotavljali dolgoročne trende v arealih posameznih vrst. Razvoj akcijskega načrta za to skupino ptic in/ ali posamezne ključne vrste pa bi nadalje pripomogel k ugotavljanju prioritet in spodbujal ozaveščenost o potrebi stalnega spremljanja ptic roparic.

8. References

Andrews, D.J. (1992): The diet of wintering Short-eared Owls on Strangford Lough, Co. Down. – Irish Birds 4: 549–554.

Banks, A.N., Coombes, R.H. & Crick, H.Q.P. (2003): The Peregrine Falcon breeding population in the UK and Isle of Man in 1992. BTO Research Report No. 333. – BTO, Thetford.

Barton, C., Pollock, C., Norriss, D.W., Nagle, T., Oliver, G.A. & Newton, S. (2006): The second national survey of breeding Hen Harriers *Circus cyaneus* in Ireland 2005. – Irish Birds 8: 1–20.

BIRDWATCH IRELAND (2010): Action Plan for Upland Birds in Ireland 2011–2020. BirdWatch Ireland's Group Action Plans for Irish Birds. — BirdWatch Ireland, Kilcoole, Co. Wicklow.

CLARKE, R. & SCOTT, D. (1994): Breeding season diet of the Merlin in County Antrim. – Irish Birds 5: 205–206.

Сооке, A., Nagle, A. & Fairley, J.S. (1995): The diet of Long-eared Owls within the range of the Bank Vole in Co. Cork. – Irish Birds 5: 305–307.

CRICK, H.Q.P. & RATCLIFFE, D.A. (1995): The Peregrine *Falco peregrinus* population of the United Kingdom in 1991. – Bird Study 42 (1): 1–19.

D'ARCY, G. (1999): Ireland's Lost Birds. – Four Courts Press, Dublin.

GIBBONS, D.W., REID, J.B. & CHAPMAN, R.A. (1993): The New Atlas of Breeding Birds in Britain and Ireland 1988–1991. – T & A D Poyser, London.

Fernández-Bellon, D. & Lusby, J. (2011A): The effectiveness of playback as a method for monitoring

- breeding Merlin *Falco columbarius* in Ireland. Irish Birds 9: 155–158.
- FERNÁNDEZ-BELLON, D. & LUSBY, J. (2011B): The feeding ecology of Merlin *Falco columbarius* during the breeding season in Ireland, and an assessment of current diet analysis methods. – Irish Birds 9: 159–164.
- GOLDEN EAGLE TRUST LTD. (2009): Ireland's failure to protect birds of prey. Unpublished report to the EU Commission Directorate General, November 2009.
- HILLIS, J.P. (2008): Rare Irish breeding birds, 2007. The Seventh Annual Report of the Irish Rare Breeding Birds Panel. – Irish Birds 8: 365–372.
- IRISH RAPTOR STUDY GROUP (2006): Birds of Prey and Owls in Ireland. Dublin.
- Irwin, S., Wilson, M., Kelly, T.C., O'Donoghue, B.,
 O'Mahony, B., Oliver, G., Cullen, C., O'Donoghue,
 T. & O'Halloran, J. (2008): Aspects of the breeding biology of Hen Harriers *Circus cyaneus* in Ireland. Irish Birds 8: 331–334.
- LACK, P. (1986): The Atlas of Wintering Birds in Britain and Ireland. – British Trust for Ornithology & Irish Wildbird Conservancy, Calton.
- LUSBY, J. (2009): The impact of two introduced mammals on the Barn Owl (*Tyto alba*) population in Ireland. Poster presentation. Raptor Research Foundation Annual Conference, 29 September—4 October 2009, Pitlochry, Scotland.
- LUSBY, J. (2012A): Exposure of Barn Owls *Tyro alba* in Ireland to second generation anticoagulant rodenticides (2006–2011). Confidential report to the National Parks and Wildlife Service and the Department of Agriculture, Food and the Marine.
- LUSBY, J. (2012A): Barn Owl Monitoring Programme. Raptor Conservation Project, Final Report. – BirdWatch Ireland, Kilcoole.
- LUSBY, J., WATSON, D. & COPLAND, A. (2010A): The ecology and conservation of the Barn Owl *Tyto alba* in County Cork; The Cork Barn Owl Research Project. Report prepared for Cork County Council.
- LUSBY, J., WATSON, D. & O'CLERY, M. (2010B): The ecology and conservation of the Barn Owl *Tyto alba* in County Kerry; The Kerry Barn Owl Research. Report to Kerry County Council.
- LUSBY, J., FERNÁNDEZ-BELLON, D., NORRISS, D. & LAUDER, A. (2011): Assessing the effectiveness of monitoring methods for Merlin Falco columbarius in Ireland: the Pilot Merlin Survey 2010. – Irish Birds 9: 143–154.
- Lusby, J., Cregg, P. & O'Clery, M. (2012A): The West Offaly Raptor Conservation Project. Final Report to Offaly Local Development Company.
- LUSBY, J., MCDONNELL, B. AND O'CLERY, M. (2012B): The ecology and conservation of the Barn Owl *Tyto alba* in Duhallow; Duhallow Barn Owl Project. Final Report prepared for IRD Duhallow.
- Lynas, P., Newton, S.F. & Robinson, J.A. (2007): The status of birds in Ireland: an analysis of conservation concern 2008–2013. Irish Birds 8: 149–166.
- MADDEN, B., HUNT, J. & NORRISS, D. (2009): The 2002 survey of the peregrine *Falco peregrines* breeding population in the republic of Ireland. Irish Birds 8: 543–548.

- McElheron, A. (2005): Merlins of the Wicklow Mountains. – Currach Press, Dublin.
- Mee, A. (2009): The Irish White-tailed Eagle reintroduction programme 2007–2008. Unpublished report to the Directorate of Nature Management, Trondheim, Norway.
- MEE, A. (2010): Determining habitat use in a reintroduced population of White-tailed Sea Eagle in SW Ireland. Project ROO527. Unpublished report to The Heritage Council, Kilkenny, Ireland.
- MEE, A. & CLARKE, D. (in prep.): Population and range expansion in Ireland: the 2011–2012 national soaring survey.
- Moore, N.P., Kelly, P.F. & Lang, F. (1992): Quarry-nesting by Peregrine Falcons in Ireland. – Irish Birds 4: 519–524.
- MOORE, N.P., KELLY, P.F. & LANG, F., Lynch, J.M. & Langton, S.D. (1997): The Peregrine *Falco peregrinus* in quarries: current status and factors influencing occupancy in the Republic of Ireland. Bird Study 44 (2): 176–181.
- Nagle, T. (2007): The loss of Barn Owl *Tyto alba* breeding and roost sites in Co. Cork: a contributory factor in the species decline. Irish Birds 8: 314–315.
- Norris, D.W. (1991): The status of the buzzard as a breeding species in the Republic of Ireland, 1977–1991.

 Irish Birds 4: 291–198.
- NORRISS, D.W. (1995): The 1991 survey and weather impacts on the Peregrine *Falco peregrinus* breeding population in the Republic of Ireland. – Bird Study 42 (1): 20–30.
- NORRISS, D.W., MARSH, J., McMahon, D. & OLIVER, G.A. (2002): A national survey of breeding Hen Harriers *Circus cyaneus* in Ireland 1998–2000. Irish Birds 7: 1–12.
- NORRISS, D.W., HARAN, B., HENNIGAN, A., McELHERON, D.J., McLaughlin, D.J., Swan, V. & Walsh, A. (2010): Breeding biology of Merlins *Falco columbarious* in Ireland, 1986–1992. Irish Birds 9: 23–30.
- O'CLERY, M., McDonnell, B. & Lusby, J. (2012): The Duhallow Raptor Conservation Project. Final Report to IRD Duhallow.
- O'Connell, P., Cogan, R. & Dunne, J. (2007): The diet of the Barn Owl *Tyto alba* at two sites in Co. Galway. – Irish Birds 8: 91–96.
- O'DONOGHUE, B. (2010): The hen harrier (*Circus cyaneus*) in Ireland; its ecology and conservation. PhD thesis. University College Cork, Ireland.
- O'Toole, L., Fielding, A.H. & Haworth, P.F. (2002): Reintroduction of the golden eagle into the Republic of Ireland. – Biological Conservation 103 (3): 303–312.
- Pearce-Higgins, J.W., Stephen, L., Langston, R.H.W., Bainbridge, I.P. & Bullman, R. (2009): The distribution of breeding birds around upland wind farms. – Journal of Applied Ecology 46 (6): 1323–1331.
- ROONEY, E. (2013): Ecology and breeding biology of the Common Buzzard *Buteo buteo*, in Ireland. Unpublished thesis. Queen's University, Belfast.
- ROONEY, E. & MONTGOMERIE, W.I. (2013): Diet diversity of the Common Buzzard (*Buteo buteo*) in a vole-less environment. Bird Study 60 (2): 147–155.
- Ruddock, M. (2007): The importance of Peregrine diet

- studies in resolving predator-prey conflicts. pp. 50 In: Sielicki, S. & Sielicki, J. (eds.): Abstracts of the Peregrine conference, 19–23 September 2007, Poland. TURUL Robert Zmuda, Warszawa.
- RUDDOCK, M. (2008): Housewives' planning: Population ecology, predation and prey selection in the peregrine falcon. Unpublished thesis. – Queen's University, Belfast.
- RUDDOCK, M., DUNLOP, B.J., O' TOOLE, L., MEE, A. & NAGLE, T. (2011): Republic of Ireland Hen Harrier survey 2010. Unpublished report by the Irish Raptor Study Group and the Golden Eagle Trust Ltd for the Department of Environment, Heritage and Local Government, Ireland.
- Scott, D. (2000): Marking a decade of tree-nesting by Hen Harriers *Circus cyaneus* in Northern Ireland, 1991–2000. – Irish Birds 6: 586–589.
- Scott, D. & McHaffie, P. (2008): Hen Harrier *Circus cyaneus* killed at windfarm site in County Antrim. Irish Birds 8: 436–437.
- SHARROCK, J.T.R. (1976): The Atlas of Breeding Birds in Britain and Ireland. T & A D Poyser, Berkhamsted, Herts.
- SIM, I.M.W., GIBBONS, D.W., BAINBRIDGE, I.P. & MATTINGLEY, W.A. (2001): Status of hen harrier *Circus cyaneus* in the UK and the Isle of Man in 1998. – Bird Study 48 (3): 341–353.
- SIM, I.M.W., DILLON, I.A., EATON, M.A., ETHERIDGE, B., LINDLEY, P., RILEY, H., SAUNDERS, R., SHARPE, C. & TICKNER, M. (2007): Status of the Hen Harrier *Circus cyaneus* in the UK and Isle of Man in 2004, and a comparison with the 1988/89 and 1998 surveys. Bird Study 54 (2): 256–267.
- SMAL, C.M. (1989): Notes on the diet of Short-eared Owls in Ireland. Irish Birds 4: 73–75.
- Walsh, P.M. & Sleeman, D.P. (1988): Avian prey of a wintering Short-eared Owl population in south-west Ireland. – Irish Birds 3: 589–591.
- Wells, J. (2007): The Northern Ireland Peregrine population 1977–2007. pp. 57 In: Sielicki, S. & Sielicki, J. (eds.): Abstracts of the Peregrine conference, 19–23 September 2007, Poland. – TURUL Robert Zmuda, Warszawa.
- WILSON, M.W., IRWIN, S., NORRISS, D.W., NEWTON, S.F., COLLINS, K., KELLY, T.C. & O'HALLORAN, J. (2009): The importance of pre-thicket conifer plantations for nesting Hen Harriers *Circus cyaneus* in Ireland. – Ibis 151 (2): 332–343.
- WILSON, M.W., O'DONOGHUE, B., O'MAHONY, B., CULLEN, C., O'DONOGHUE, T., OLIVER, G., RYAN, B., TROAKE, P., IRWIN, S., KELLY, T.C., ROTELLA, J.J. & O'HALLORAN, J. (2012): Mismatches between breeding success and habitat preferences in Hen Harriers Circus cyaneus breeding in forested landscapes. – Ibis 154 (3): 578–589.

Arrived / Prispelo: 27. 3. 2013 Accepted / Sprejeto: 1. 7. 2013