

NEW LOCALITY AND SITES OF *PULMONARIA AUSTRALIS* IN THE TRIGLAV MOUNTAINS (THE JULIAN ALPS, NW SLOVENIA)

NOVO NAHAJALIŠČE IN RASTIŠČA VRSTE *PULMONARIA AUSTRALIS* V TRIGLAVSKEM POGORJU (JULIJSKE ALPE, SLOVENIJA)

Igor DAKSKOBLER¹ & Branko ZUPAN²

ABSTRACT UDC 581.9(234.323.6:497.4):582.901
New locality and sites of *Pulmonaria australis* in the Triglav Mountains (the Julian Alps, NW Slovenia)

The paper describes a new locality and sites of *Pulmonaria australis* in the Triglav Mountains under the peak Spodnja Brda on the northern edge of the Pokljuka plateau. *Pulmonaria australis* is a new species in the flora of both the Triglav National Park and the eastern Julian Alps. This locality is very different from its other known localities in Slovenia in terms of its elevation belt (almost 1800 m a.s.l.) and its sites (stony grassland on an overgrown scree slope with dominating species of the class *Elyno-Seslerietea*); it is, however, similar to some *Pulmonaria australis* sites in other Alpine regions, such as South Tyrol in N Italy.

Key words: phytogeography, phytosociology, *Pulmonaria australis*, Julian Alps, Triglav National Park, north-western Slovenia

IZVLEČEK UDK 581.9(234.323.6:497.4):582.901
Novo nahajališče in rastišča vrste *Pulmonaria australis* v Triglavskem pogorju (Julijske Alpe, Slovenija)

Opisujemo novo nahajališče in rastišča vrste *Pulmonaria australis* v Triglavskem pogorju, pod vzpetino Spodnja Brda na severnem robu Pokljuke. To je novost za floro Triglavskega narodnega parka in vzhodnih Julijskih Alp. Po višinskem pasu uspevanja (skoraj 1800 m nm. v.) in rastiščih (kamnito travnišče na poraslem melišču s prevladujočimi vrstami razreda *Elyno-Seslerietea*) se to nahajališče precej razlikuje od njenih drugih znanih nahajališč v Sloveniji, podobno pa je nekaterim njenim nahajališčem v drugih alpskih pokrajinah, na primer na Južnem Tirolskem v severni Italiji.

Ključne besede: fitogeografija, fitocenologija, *Pulmonaria australis*, Julijske Alpe, Triglavski narodni park, severozahodna Slovenija

¹ Dr., the Jovan Hadži Institute of Biology of the Scientific Research Centre of the Slovenian Academy of Sciences and Arts, Regional research unit, Brunov drevored 13, SI-5220 Tolmin, igor.dakskobler@zrc-sazu.si

² Savica 6, SI-4264 Bohinjska Bistrica

1 INTRODUCTION

Pulmonaria australis is an Alpine-Illyrian species, a character species of thermophilous forest edges and tall herbs on nutrient-poor sites from the class *Trifolio-Gerenietea*. In the Alps, it occurs in the montane and subalpine belt, rarely also in the colline and even in the alpine belt on calcareous, mixed calcareous-silicate, rarely also on silicate bedrock, on neutral to alkaline dry soils (AESCHIMANN et al. 2004: 74). Its sites in Slovenia include dry meadows, shrubs and light-filled woods from lowland to the montane belt, with certainty mainly in the sub-Mediterranean and Dinaric phytogeographical regions (JOGAN 2007: 538). Its distribution in Slovenia has not yet been fully researched. In the past, there was no differentiation between the similar taxa *Pulmonaria angustifolia* and *P. australis* (the first description of the latter dates back to 1975 – SAUER 1975), with both determined as *P. angustifolia*. For *Pulmonaria angustifolia*, HORVATÍČ & TOMAŽIČ (1939: 82) list localities beyond the Javornik hills, in the Notranjska and Gorica regions (citing Paulin), scattered and rare localities in the Podravje region around Radgona, near Slovenj Gradec and Celje (Sv. Miklavž) – the source for these localities is Hayek. They report its find on the southern margins of Mt. Begunjščica by R. Justin (herbarium), but this specimen is currently not held in LJU (BAČIČ, in litt.). As a new locality they mention the Polhograjsko hribovje hills (to the south of Grmada, north and south from Goljak). JOGAN (ibid.) is of the opinion that *Pul-*

monaria angustifolia most likely occurs only in eastern Slovenia. JOGAN et al. (2001: 305) published only the distribution map for the species aggregate *Pulmonaria angustifolia* agg., but without any marked localities in the Alps (not even Justin's locality under Mt. Begunjščica). Without a thorough revision of herbarium material it is therefore impossible to give a detailed presentation of the distribution of *Pulmonaria australis* in Slovenia. According to our information it is generally distributed in the Central Soča Valley, but is frequent also in Karst (see also ZUPANČIČ & ŽAGAR 2006) and Istria (including the Italian part of the Gorica and Trieste Karst – POLDINI 2009: 436). In the foothills of the Julian Alps it was found on the Hlevnik ridge and under the neighbouring Kolovrat ridge (9848/1) – DAKSKOBLER (1998), as well as on the southern slopes of Mt. Matajur above Jevšček, at the elevation of 940 m (DAKSKOBLER 2005, Table 2, relevé No. 13). It is more frequent in the foothills of the Italian part of the Julian Alps (the ridge of Stol, Monte Musi/Muzci, Resia, the Raccolana Valley) – POLDINI 2002: 394, GOBBO & POLDINI 2005: 183). We find Branko Zupan's discovery in the Triglav Mts., at the elevation of almost 1800 m and in completely different communities than those known until then, surprising and worthy of detailed presentation. We therefore present the new locality and compare its site with other sites known so far in the southern foothills of the Julian Alps.

2 METHODS

Floristic records and phytosociological relevés were made according to the established Central European methods (EHRENDORFER & HAMANN 1965, BRAUN-BLANQUET 1964) and entered into the FloVegSi database (T. SELIŠKAR, VREŠ & A. SELIŠKAR 2003). Combined cover-abundance values were transformed into numerical values 1–9 (van der Maarel 1979). Three relevés were mutually compared by means of hierarchical classification using the software package SYN-TAX 2000 (PODANI 2001). We selected the unweighted pair group method with arithmetic mean (UPGMA) and used Wishart's similarity ratio. In the determination of *Pulmonaria australis* we used the determination keys from the Mala flora Slovenije (JOGAN 2007)

and from the Austrian Excursion Flora (FISCHER et al. 2008) as well as descriptions in MERXMÜLLER & SAUER (1973) and SAUER (1975). The nomenclature source for the names of vascular plants is the Mala flora Slovenije (MARTINČIČ et al. 2007). MARTINČIČ (2003, 2011) is the nomenclature source for the names of mosses. The nomenclature sources for the names of syntaxa are THEURILLAT (2004) and ŠILC & ČARNI (2012). The source for the names of soil types are URBANČIČ et al. (2005). The climatic data are cited from J. PRISTOV, N. PRISTOV & B. ZUPANČIČ (1998), B. ZUPANČIČ (1998) and CEGNAR (1998), and the data on geological bedrock from JURKOVŠEK (1987 a,b).

3 RESULTS AND DISCUSSION

3.1 Description of the new locality of *Pulmonaria australis* in the Julian Alps

9649/2 (UTM 33TVM13): Slovenia, the Julian Alps, Triglav Mts., northern edge of Pokljuka, Lipanca pasture above the path towards Klek pasture under Brda pasture or the peak Spodnja Brda, Triassic limestone and talus scree, stony grassland on an overgrown scree slope under a rock step, lithosol and rendzina, 1760 m – 1780 m a.s.l. Leg. B. Zupan & P. Strgar, 29. 5. 2012 and 22. 7. 2013; leg. & det. B. Zupan & I. Dakskobler, 1.8. 2013, herbarium LJS.

Pulmonaria australis was found in an area totaling about 10 acres (Figure 1). It grows on an overgrown scree slope under a rock step, in communities

whose species composition is presented in Table 1. Localities are still in the forest belt, in the belt of open larch-spruce forest classified into the syntaxa *Adenostylo glabrae-Piceetum laricetosum* and *Rhodothamno-Laricetum*. In places, on more extreme sites, dwarf pine stands with larch (*Rhododendro hirsuti-Pinetum mugo laricetosum* = *Rhodothamno-Pinetum mugo laricetosum*) occur. Rock walls, gravelly gullies and depressions where snow from avalanches accumulates each year are not covered with forests. Fragments of green alder (*Rhododendro hirsuti-Alnetum viridis*) and single shrubs of *Rhamnus fallax* were spotted in moist gullies. Slopes face the northeast, east and partly the southeast. The floristic composition of overgrown screes indicates a certain similarity with the stands of the associations *Astrantio carniolicae-Adenostyletum glabrae*, *Caricetum ferruginei* s. lat., *Rhodothamno-Rhododendretum hirsuti* and *Ranunculo hybridi-Caricetum sempervirentis*. Relevés were adapted to the occurrence of *Pulmonaria australis* and therefore do not represent stands typical for said communities, but indicate syndynamic transitions between them. In terms of proportion they are dominated by diagnostic species of subalpine-alpine grasslands from the class *Elyno-Seslerietea*, tall herbs and shrubs from the classes *Mulgedio-Aconitetea* and *Betulo-Alnetea*, scree species from the class *Thlaspietea rotundifolii* and pine and spruce forest species from the classes *Erico-Pinetea* and *Vaccinio-Piceetea*. The only diagnostic species of the class *Trifolio-Geranietea* is *Pulmonaria australis* (Table 2). Under the peak Spodnja Brda on the northern edge of the Pokljuka plateau it grows on shallow initial soil, in stony subalpine grasslands in the belt of larch-spruce forest that continues into dwarf pine on steeper slopes. The climate is montane, with mean annual temperature of 2 °C – 4 °C and mean annual precipitation between 2000 mm and 2500 mm.



Figure 1: Localities of *Pulmonaria australis* under the peak Spodnja Brda in the Triglav Mountains
Slika 1: Nahajališča vrste *Pulmonaria australis* pod vzpetino Spodnja Brda v Triglavskem pogorju

3.2 Comparison of sites of *Pulmonaria australis* in the Triglav Mountains with sites in the southern foothills of the Julian Alps

In the southern foothills of the Julian Alps, *Pulmonaria australis* was most frequently recorded on dry to mesic moist grasslands on mixed geological bedrock (limestone and marlstone, limestone, marlstone, chert, flysch) and on relatively deep soils (brown calcareous soils and Eutric brown soils). 26 relevés of dry and semi-dry grasslands (mainly of the order *Scorzonero villosae-Chrysopogonetalia grylli* and the alliance *Scor-*

zonarion villosae, some of which have already been published – DAKSKOBLER 2005) from the foothills of Matajur, from Kanalski Kolovrat, Korada and Banjšice were selected for the purpose of comparison of floristic composition. These relevés comprise very few, a total of 13 species growing in the stands in the Triglav Mountains. In addition to *Pulmonaria australis* these species include *Anemone nemorosa*, *Buphthalmum salicifolium*, *Carlina acaulis*, *Erica carnea*, *Hypericum maculatum*, *Knautia drymeia*, *Listera ovata*, *Lotus corniculatus* s. lat., *Salix appendiculata*, *Solidago virgaurea*, *Ranunculus nemorosus* agg. and *Betonica alopecu-*

rus. Floristic similarity of the three relevés from the Triglav Mts. with 26 relevés from the southern margins of the Julian Alps according to SØRENSEN (1948) is only 7%. On the northern edge of Pokljuka, *Pulmonaria australis* grows in very different site conditions and within very different communities than known in Slovenia so far. This, however, does not refer to its sites elsewhere in the Alps. In South Tyrol in northern Italy it is characteristic for dry grasslands in open larch forests and can grow as high as 2000 m a.s.l. (WILHALM, in litt.), in conditions that at least slightly resemble those on the newly described site.

4 CONCLUSIONS

Pulmonaria australis is an Alpine-Illyrian species from the aggregate species *Pulmonaria angustifolia* agg., which has not been fully researched with certainty as of yet (FISCHER et al. 2008). The same applies to its distribution in Slovenia (JOGAN 2007), where it is frequent in the Primorska region and partly also in the Notranjska region, in the pre-Alpine, sub-Mediterranean and Dinaric phytogeographical regions. Its occurrence in other Slovenian regions is still to be verified. The new locality described in the Triglav Mts. is new both to the flora of the Triglav National Park and to the eastern part of the Julian Alps. In terms of its elevation belt and sites it is very different from other localities described so far in Slovenia. The exception

are some of its sites in other Alpine regions (AESCHIMANN et al., ibid, WILHALM, in litt.). As it is very inconspicuous in the summer and often covered by other plants it is likely to grow on other sites on the northern edge of Pokljuka, under the ridge Mrežce–Brda–Debela Peč–Klečica. On the sites known to us in the Primorska region it most often grows in communities from the classes *Trifolio-Geranietea*, *Festuco-Brometea* and *Quercu-Fagetea*, which do not usually grow on elevations of 1700 m or higher. Its phytosociological amplitude obviously exceeds the elevation belt of communities from the class *Trifolio-Geranietea*, so it is only conditionally diagnostic for this class.

5 POVZETEK

5.1 Uvod

Pulmonaria australis je alpsko-ilirska vrsta, značilnica združb termofilnih gozdnih robov in visokih steblik na s hranili siromašnih rastiščih iz razreda *Trifolio-Geranietea*. V Alpah uspeva v montanskem in subalpinskem pasu, redkeje tudi v kolinskem in celo alpskem pasu na karbonatni, mešani karbonatno-silikatni in redko tudi na silikatni podlagi, na nevtralnih do bazičnih suhih tleh (AESCHIMANN et al. 2004: 74). V Sloveniji so njena rastišča suhi travniki, grmovna mesta in svetli gozdovi od nižine do montanskega pasu, zanesljivo predvsem v submediteranskem in dinarskem fitogeografskem območju (JOGAN 2007: 538). Njena celotna razširjenost v Sloveniji je še slabo raziskana. V preteklosti niso razlikovali med podobnima taksonoma *Pulmonaria angustifolia* in *P. australis*

(slednji je bil opisan šele leta 1975 – SAUER 1975) in so oba določali kot vrsto *P. angustifolia*. HORVATIĆ & TOMAŽIČ (1939: 82) tako za vrsto *Pulmonaria angustifolia* naštevata nahajališča onstran Javornikov na Notranjskem in Goriškem (tu povzemata Paulina), raztresena in redka nahajališča v Podravju okoli Radgone, pri Slovenj Gradcu in Celju (Sv. Miklavž) – tu povzemata Hayeka. Na južnih obronkih Begunjščice naj bi jo našel R. Justin (herbarij) – v LJU žal tega primerka zdaj nimajo (BAČIČ, in litt.), kot novo nahajališče pa navajata Polhograjsko hribovje (južno od Grmade, severno in južno od Goljaka). JOGAN (ibid.) meni, da vrsta *Pulmonaria angustifolia* najverjetneje uspeva le v vzhodni Sloveniji. JOGAN et al. (2001: 305) so objavili le arealno karto za skupno vrsto *Pulmonaria angustifolia* agg., v njej pa ni označeno nobeno nahajališče v Alpah (tudi ne Justinovo nahajališče

pod Begunjščico). Brez podrobne revizije herbarijskega gradiva torej za zdaj ne moremo prikazati natančne razširjenosti vrste *Pulmonaria australis* v Sloveniji. Po naših podatkih je splošno razširjena v Srednjem Posočju, pogosta tudi na Krasu (glej tudi ZUPANČIČ & ŽAGAR 2006) in v Istri (to velja tudi za italijanski del goriškega in tržaškega Krasa – POLDINI 2009: 436). V prigorju Julijskih Alp smo jo našli na grebenu Hlevnika in pod sosednjim grebenom Kolovrata (9848/1) – DAKSKOBLER (1998) in na južnih pobočjih Matajurja nad Jevščkom, na nadmorski višini 940 m (DAKSKOBLER 2005, tabela 2, popis št. 13). Pogostejša je v prigorju italijanskega dela Julijskih Alp (Stolov greben, Muzci, Rezija, Reklanska dolina) – POLDINI 2002: 394, GOBBO & POLDINI 2005: 183). Najdba starejšega od naju, Branka Zupana, v Triglavskem pogorju, na nadmorski višini skoraj 1800 m, in v povsem drugačnih združbah, kot smo jo poznali doslej, se nam zdi presenetljiva in vredna, da novo nahajališče podrobneje predstavimo in primerjamo njeno rastišče s drugimi do zdaj znanimi rastišči v južnem prigorju Julijskih Alp.

5.2 Metode

Floristične in fitocenološke popise smo naredili po ustaljenih srednjeevropskih metodah (EHRENDORFER & HAMANN 1965, BRAUN-BLANQUET 1964) in jih vnesli v bazo podatkov FloVegSi (T. SELIŠKAR, VREŠ & A. SELIŠKAR 2003). Kombinirane ocene zastiranja in obilnosti smo pretvorili v števila od 1 do 9 (van der MAAREL 1979). Tri fitocenološke popise smo med seboj primerjali s hierahično klasifikacijo in pri tem uporabili programski paket SYN-TAX 2000 (PODANI 2001). Izbrali smo metodo kopičenja na podlagi povezovanja (netehtanih) srednjih razdalj (UPGMA) in kot količnik različnosti uporabili Wishartov koeficient similarity ratio. Pri določanju vrste *Pulmonaria australis* smo uporabili določevalna ključa v Mali flori Slovenije (JOGAN 2007) in avstrijski ekskurzijski flori (FISCHER et al. 2008) kot tudi opisa v MERXMÜLLER & SAUER (1973) in SAUER (1975). Nomenklaturni vir za imena praprotnic in semenk je Mala flora Slovenije (MARTINČIČ et al. 2007). MARTINČIČ (2003, 2011) je nomenklaturni vir za imena mahov. Nomenklaturni vir za imena sintaksonov so THEURILLAT (2004) in ŠILC & ČARNI (2012). Vir za imenovanje talnih tipov so URBANČIČ et al. (2005). Podnebne podatke povzemamo po J. PRISTOV, N. PRISTOV & B. ZUPANČIČ (1998), B. ZUPANČIČ (1998) in CEGNAR (1998), podatke o geološki podlagi pa po JURKOVŠEK (1987 a, b).

5.3 Rezultati in razprava

5.3.1 Oznaka novega nahajališča vrste *Pulmonaria australis* v Julijskih Alpah

9649/2 (UTM 33TVM13): Slovenija, Julijske Alpe, Triglavsko pogorje, severni rob Pokljuke, pl. Lipanca, nad potjo proti pl. Klek, pod pl. Brda oz. vzpetino Spodnja Brda, triasni apnenec in pobočni grušč, kamnito travnišče na poraslem melišču pod skalno stopnjo, litosol in rendzina, 1760 m do 1780 m nm. v. Leg. B. Zupan & P. Strgar, 29. 5. 2012 in 22. 7. 2013; leg. & det. B. Zupan & I. Dakskobler, 1. 8. 2013, herbarij LJS.

Vrsto *Pulmonaria australis* smo opazili na skupni površini okoli 10 arov. Raste na poraslem melišču pod skalno stopnjo, v združbah, katerih vrstno sestavo prikazuje preglednica 1. Nahajališča so še v gozdnem pasu in sicer v pasu vrzelastega macesnovo-smrekovega gozda, ki ga uvrščamo v sintaksona *Adenostylo glabrae-Piceetum laricetosum* in *Rhododendro-Laricetum*. Na bolj skrajnih rastiščih ponekod uspeva ruševje z macesnom (*Rhododendro hirsuti-Pinetum mugo laricetosum* = *Rhododendro-Pinetum mugo laricetosum*). Z gozdom neporasli so stene ter gruščnati žlebovi in kotanje, kjer se vsako leto kopičijo snežni plazovi. V vlažnih žlebovih smo opazili tudi fragmente zelenega jelševja (*Rhododendro hirsuti-Alnetum viridis*) in posemne grme kranjske krhlike (*Rhamnus fallax*). Pobočja so obrnjena proti severovzhodu, vzhodu in deloma jugovzhodu. Floristična sestava poraslih melišč kaže določeno podobnost s sestoji asociacij *Astrantio carniolicae-Adenostyletum glabrae*, *Caricetum ferruginei* s. lat., *Rhododendro-Rhododendretum hirsuti* in *Ranunculo hybridi-Caricetum sempervirentis*. Fitocenološke popise smo prilagodili prisotnosti vrste *Pulmonaria australis* in zato ne predstavljajo za omenjene združbe tipičnih sestojev, temveč kažejo na sindinamske prehode med njimi. V njih po deležu prevladujejo vrste subalpinsko-alpinskih travnišč iz razreda *Elvino-Seslerietea*, vrste visokih steblik in grmišč iz razredov *Mulgedio-Aconitetea* in *Betulo-Alnetea*, vrste melišč iz razreda *Thlaspietea rotundifoliae* ter vrste borovih in smrekovih gozdov iz razredov *Erico-Pinetea* in *Vaccinio-Piceetea*. Edina diagnostična vrsta razreda *Trifolio-Geranietea* je *Pulmonaria australis* (preglednica 2). Pod vzpetino Spodnja Brda na severnem robu Pokljuke ta vrsta torej raste na plitvih, inicialnih tleh, v subalpinskih kamnitih travniščih v pasu macesnovo-smrekovega gozda, ki v strminah prehaja v ruševje. Podnebje je gorsko, s povprečno letno temperaturo 2 °C – 4 °C in povprečno letno množino padavin med 2000 mm in 2500 mm.

5.3.2 Primejava rastišč vrste *Pulmonaria australis* v Triglavskem pogorju z rastišči v južnem prigorju Julijskih Alp

V južnem prigorju Julijskih Alp smo vrsto *Pulmonaria australis* največkrat popisali na suhih do zmerno vlažnih traviških na mešani geološki podlagi (apnenec in laporovec, apnenec, laporovec, roženec, fliš) in na razmeroma globokih tleh (rjava pokarbonatna in evtrična rjava tla). Za primerjavo floristične sestave smo izbrali 26 popisov suhih in polsuhih travišč (večinoma priopadajo redu *Scorzenero villosae*-*Chrysopogonetalia grylli* in zvezi *Scorzenerion villosae*, nekateri so bili že objavljeni – DAKSKOBLER 2005) iz prigorja Matajurja, s Kanalskega Kolovrata, Korade in Banjšič. V teh popisih je zelo malo vrst, ki uspevajo tudi v sestojih v Triglavskem pogorju, skupno 13. Poleg vrste *Pulmonaria australis* so to še vrste *Anemone nemorosa*, *Buphthalmum salicifolium*, *Carlina acaulis*, *Erica carnea*, *Hypericum maculatum*, *Knautia drymeia*, *Listera ovata*, *Lotus corniculatus* s. lat., *Salix appendiculata*, *Solidago virgaurea*, *Ranunculus nemorosus* agg. in *Betonica alopecurus*. Floristična podobnost treh popisov iz Triglavskega pogorja s 26 popisi iz južnega obrobja Julijskih Alp po SØRENSEN (1948) je le 7 %. Vrsta *Pulmonaria australis* na severnem robu Pokljuke uspeva v precej drugačnih rastiščnih razmerah in v precej drugačnih združbah, kot smo jo poznali doslej v Sloveniji. To pa ne velja za njena rastišča drugod v Alpah. Na Južnem Tirolskem v Severni Italiji je ta vrsta tipična za suha rastišča v vrzelastih macesnovih gozdovih in raste tudi na nadmorski višini več kot 2000 m

(WILHALM, in litt.), torej vsaj v nekoliko podobnih razmerah kot na opisanem novem nahajališču.

5.4 Zaključki

Pulmonaria australis je alpsko-ilirska vrsta iz agregata *Pulmonaria angustifolia* agg., ki še ni povsem zanesljivo raziskana (FISCHER et al. 2008). Enako velja za njeno razširjenost v Sloveniji (JOGAN 2007), kjer je pogosta na Primorskem in deloma Notranjskem, v predalpskem, submediteranskem in dinarskem fitogeografskem območju. Njeno pojavljanje v drugih slovenskih pokrajinah je treba še preveriti. Novo opisano nahajališče v Triglavskem pogorju je novost za floro Triglavskega narodnega parka in vzhodnega dela Julijskih Alp. Po višinskem pasu uspevanja in rastiščih se zelo razlikuje od doslej znanih ostalih nahajališč v Sloveniji. Ta ugotovitev pa ne velja za nekatera njena nahajališča v drugih alpskih pokrajinah (AESCHIMANN et al., ibid, WILHALM, in litt.). Verjetno na severnem robu Pokljuke, pod grebenom Mrežce–Brda–Debela peč–Klečica uspeva še kje, saj je v poletnem času razmeroma slabo opazna in jo zakrijejo druge rastline. Na nam znanih nahajališčih na Primorskem največkrat raste v združbah iz razredov *Trifolio-Geranieta*, *Festuco-Brometea* in *Quercu-Fagetea*. Združbe iz teh treh razredov navadno ne uspevajo na nadmorski višini 1700 m in več. Očitno njena fitocenološka amplituda presega višinski pas uspevanja združb iz razreda *Trifolio-Geranieta* in je za ta razred diagnostična le pogojno.

ACKNOWLEDGEMENTS – ZAHVALA

The information on the sites of *Pulmonaria australis* in South Tyrol was kindly provided to us by Dr. Thomas Wilhalm (Bozen / Bolzano) and the information on the herbarium sheets for this species in LJU by Doc. Dr. Tinka Bačič. We are grateful to Academician Dr.

Mitja Zupančič, Prof. Dr. Nejc Jogan and Doc. Dr. Božo Frajman for their valuable advice and to Peter Strgar for his photographs and for accompanying us in the field. Iztok Sajko prepared Figure 1 for print. English translation by Andreja Šalamon Verbič.

REFERENCES – LITERATURA

- AESCHIMANN, D., K. LAUBER, D. M. MOSER & J.-P. THEURILLAT, 2004 b: *Flora alpina*. Bd. 2: *Gentianaceae–Orchidaceae*. Haupt Verlag, Bern, Stuttgart, Wien.
- BRAUN-BLANQUET, J., 1964: *Pflanzensoziologie. Grundzüge der Vegetationskunde*. 3. Auflage. Springer, Wien – New York.
- CEGNAR, T., 1998: *Temperatura zraka*. In: Fridl, J., D. Kladnik, M. Orožen Adamič & D. Perko: *Geografski atlas Slovenije. Država v prostoru in času*. Državna založba Slovenije, Ljubljana, pp. 100–101.

- DAKSKOBLER, I. 1998: *Pulmonaria australis*. V: Jogan, N. (ed.): *Nova nahajališča – New localities*. Hladnikia (Ljubljana) 10: 63.
- DAKSKOBLER, I., 2005: *Floristične novosti iz Posočja in sosednjih območij v zahodni in severozahodni Sloveniji – IV*. Hacquetia (Ljubljana) 4 (2): 173–200.
- EHRENDORFER, F. & U. HAMANN, 1965: *Vorschläge zu einer floristischen Kartierung von Mitteleuropa*. Ber. Deutsch. Bot. Ges. 78: 35–50.
- FISCHER M. A., W. ADLER & K. OSWALD, 2008: *Exkursionsflora von Österreich, Liechtenstein und Südtirol*. Land Oberösterreich, Biologiezentrum der OÖ Landesmuseen, Linz.
- GOBBO, G. & L. POLDINI, 2005: *La diversità floristica del parco delle Prealpi Giulie. Atlante corologico*. Università degli Studi di Trieste, Dipartimento di Biologia, Trieste.
- HORVATIĆ, S. & G. TOMAŽIČ, 1939: *Donos k spoznavanju flore Slovenije*. Hrvatski geografski glasnik (Zagreb) 8–9–10: 80–84.
- JOGAN, N., 2007: *Pulmonaria L. – pljučnik*. V: Martinčič, A. (ur.): *Mala flora Slovenije. Ključ za določanje praprotnic in semenk*. Tehniška založba Slovenije, Ljubljana: 536–538.
- JOGAN, N., T. BAČIČ, B. FRAJMAN, I. LESKOVAR, D. NAGLIČ, A. PODOBNIK, B. ROZMAN, S. STRGULC - KRAJŠEK & B. TRČAK, 2001: *Gradivo za Atlas flore Slovenije*. Center za kartografijo favne in flore, Miklavž na Dravskem polju.
- JURKOVŠEK, B., 1987 a: *Tolmač listov Beljak in Ponteba*. Osnovna geološka karta SFRJ 1:100 000. Zvezni geološki zavod, Beograd.
- JURKOVŠEK, B., 1987 b: *Osnovna geološka karta SFRJ. Beljak in Ponteba 1: 100 000*. Zvezni geološki zavod, Beograd.
- MAAREL van der, E., 1979: *Transformation of cover-abundance values in phytosociology and its effects on community similarity*. Vegetatio 39 (2): 97–114.
- MARTINČIČ, A., 2003: *Seznam listnatih mahov (Bryopsida) Slovenije*. Hacquetia (Ljubljana) 2 (1): 91–166.
- MARTINČIČ, A., 2011: *Seznam jetrenjakov (Marchanthiophyta) in rogovnjakov (Anthocerotophyta) Slovenije. Annotated Checklist of Slovenian Liverworts (Marchanthiophyta) and Hornworts (Anthocerotophyta)*. Scopolia (Ljubljana) 72: 1–38.
- MARTINČIČ, A., T. WRABER, N. JOGAN, A. PODOBNIK, B. TURK, B. VREŠ, V. RAVNIK, B. FRAJMAN, S. STRGULC KRAJŠEK, B. TRČAK, T. BAČIČ, M. A. FISCHER, K. ELER & B. SURINA, 2007: *Mala flora Slovenije. Ključ za določanje praprotnic in semenk*. Četrta, dopolnjena in spremenjena izdaja. Tehniška založba Slovenije, Ljubljana.
- MERXMÜLLER, H. & W. SAUER, 1973: *Pulmonaria L.* In: Tutin, T. G. et al.: *Flora Europaea* 3. University Press, Cambridge: 100–102.
- POLDINI, L. (s sodelovanjem G. Oriolo & M. Vidali), 2002: *Nuovo Atlante corologico delle piante vascolari nel Friuli Venezia Giulia*. Regione Autonoma Friuli Venezia Giulia, Azienda Parchi e Foreste Regionali & Università degli Studi di Trieste, Dipartimento di Biologia, Udine.
- PODANI, J., 2001: SYN-TAX 2000. *Computer Programs for Data Analysis in Ecology and Systematics*. User's Manual, Budapest.
- PRISTOV, J., N. PRISTOV & B. ZUPANČIČ, 1998: *Klima Triglavskega narodnega parka*. Hidrometeorološki zavod, Ljubljana in Triglavski narodni park, Bled.
- SAUER, W., 1975: *Karyo-systematische Untersuchungen an der Gattung Pulmonaria (Boraginaceae)*. Bibliotheca Botanica (Stuttgart) 131: 1–85.
- SELIŠKAR, T., B. VREŠ & A. SELIŠKAR, 2003: FloVegSi 2.0. *Računalniški program za urejanje in analizo bioloških podatkov*. Biološki inštitut ZRC SAZU, Ljubljana.
- SØRENSEN, Th., 1948: *A method of establishing groups of equal amplitude in plant sociology based on similarity of species content*. Det Kongelige Danske Videnskabernes Selskab, Biologiske Skrifter (København) 5 (4): 1–34.
- ŠILC, U. & A. ČARNI, 2012: *Conspectus of vegetation syntaxa in Slovenia*. Hacquetia (Ljubljana) 11 (1): 113–164.
- THEURILLAT, J.-P., 2004: *Pflanzensoziologisches System*. V: Aeschimann, D., K. Lauber, D. M. Moser & J.-P. Theurillat: *Flora alpina* 3: Register. Haupt Verlag, Bern, Stuttgart, Wien: 301–313.
- WRABER, M., 1969: *Pflanzengeographische Stellung und Gliederung Sloweniens*. Vegetatio 17: 176–199.
- ZUPANČIČ, B., 1998: *Padavine*. In: Fridl, J., D. Kladnik, M. Orožen Adamič & D. Perko: *Geografski atlas Slovenije. Država v prostoru in času*. Državna založba Slovenije, Ljubljana, pp. 98–99.
- ZUPANČIČ, M. & V. ŽAGAR, 2006: *Association Potentillo albae-Quercetum pubescentis in Slovenia*. Razprave 4. razreda SAZU (Ljubljana) 47–1: 247–261.



Figure 2: *Pulmonaria australis* under the peak Spodnja Brda. Photo P. Strgar

Slika 2: *Pulmonaruia australis* pod vzpetino Spodnja Brda. Foto P. Strgar



Figure 3: Ground leaves of *Pulmonaria australis* under the peak Spodnja Brda. Photo I. Dakskobler

Slika 3: Pritlični listi vrste *Pulmonaria australis* pod vzpetino Spodnja Brda. Foto I. Dakskobler



Figure 4: Site of *Pulmonaria australis* under the peak Spodnja Brda. Photo I. Dakskobler

Figure 4: Rastišče vrste *Pulmonaria australis* pod vzpetino Spodnja Brda. Foto I. Dakskobler

Table 1: Communities with *Pulmonaria australis* in the Triglav Mountains (the Julian Alps)**Table 1: Združbe z vrsto *Pulmonaria australis* v Triglavskem pogorju (Julijske Alpe)**

Number of relevé (Zaporedna številka popisa)		1	2	3	Pr.
Database number of relevé (Delovna številka popisa)		250801	250803	250800	
Elevation in m (Nadmorska višina v m)		1770	1775	1770	
Aspect (Lega)		NE	NE	SEE	
Slope in degrees (Nagib v stopinjah)		35	30	30	
Parent material (Matična podlaga)		A, Gr	A, Gr	A, Gr	
Soil (Tla)		R	R		
Stoniness in % (Kamnitost v %)		10	10	40	
Cover in % (Zastiranje v %) :					
Shrub layer (Grmovna plast)	E2	20	40	.	
Herb layer (Zeliščna plast)	E1	80	70	60	
Number of species (Število vrst)		61	51	40	
Relevé area (Velikost popisne ploskve)	m ²	20	10	20	
Date of taking relevé (Datum popisa)		8/1/2013	8/1/2013	8/1/2013	
Locality (Nahajališče)		Lipanca - Brda	Lipanca - Brda	Lipanca - Brda	
Quadrant (Kvadrant)		9649/2	9649/2	9649/2	
Coordinate GK Y (D-48)		418312	418326	418306	
Coordinate GK X (D-48)		5138208	5138206	5138200	
CF <i>Caricion ferrugineae</i>					
<i>Carex ferruginea</i>	E1	3	3	+	3
CA <i>Caricion austroalpinæ</i>					
<i>Laserpitium peucedanoides</i>	E1	+	+	+	3
<i>Heracleum austriacum</i> subsp. <i>siifolium</i>	E1	1	.	+	2
<i>Festuca calva</i>	E1	+	.	.	1
ES <i>Elyno-Seslerietea</i>					
<i>Betonica alopecuroides</i>	E1	1	2	1	3
<i>Helianthemum grandiflorum</i>	E1	1	1	2	3
<i>Campanula witasekiana</i>	E1	+	1	+	3
<i>Myosotis alpestris</i>	E1	+	+	1	3
<i>Euphrasia picta</i>	E1	+	+	+	3
<i>Phyteuma orbiculare</i>	E1	+	+	.	2
<i>Heliosperma alpestre</i>	E1	+	+	.	2
<i>Sesleria caerulea</i>	E1	.	1	2	2
<i>Thymus praecox</i> subsp. <i>polytrichus</i>	E1	.	+	2	2
<i>Achillea clavensis</i>	E1	.	+	+	2
<i>Galium anisophyllum</i>	E1	.	+	+	2
<i>Pulsatilla alpina</i> subsp. <i>austroalpina</i>	E1	.	+	+	2
SCf <i>Parnassia palustris</i>	E1	+	.	.	1
<i>Leucanthemum adustum</i>	E1	.	1	.	1
<i>Acinos alpinus</i>	E1	.	1	.	1
<i>Aster bellidiflorus</i>	E1	.	+	.	1
<i>Cerastium strictum</i>	E1	.	+	.	1
<i>Helictotrichon parlatorei</i>	E1	.	+	.	1
<i>Lotus alpinus</i>	E1	.	+	.	1
<i>Polygonum viviparum</i>	E1	.	+	.	1
<i>Scabiosa lucida</i> subsp. <i>stricta</i>	E1	.	+	.	1
FB <i>Carlina acaulis</i>	E1	.	+	.	1
<i>Thesium alpinum</i>	E1	.	.	+	1
MuA <i>Mulgedio-Aconitetea</i>					
<i>Veratrum album</i> subsp. <i>lobelianum</i>	E1	1	1	+	3
<i>Aconitum lycoctonum</i> subsp. <i>ranunculifolium</i>	E1	1	+	+	3
<i>Geranium sylvaticum</i>	E1	+	+	+	3
<i>Hypericum maculatum</i>	E1	1	+	+	3
<i>Carduus carduelis</i>	E1	+	+	.	2
<i>Chaerophyllum hirsutum</i>	E1	1	.	.	1
<i>Cirsium carniolicum</i>	E1	+	.	.	1
<i>Crepis paludosa</i>	E1	+	.	.	1
<i>Epilobium alpestre</i>	E1	+	.	.	1
<i>Primula elatior</i>	E1	+	.	.	1
<i>Rumex arifolius</i>	E1	+	.	.	1
<i>Saxifraga rotundifolia</i>	E1	+	.	.	1
<i>Silene dioica</i>	E1	+	.	.	1
<i>Silene vulgaris</i> subsp. <i>antelopum</i>	E1	+	.	.	1
BA <i>Betulo-Alnetea</i>					
<i>Sorbus chamaemespilus</i>	E2a	1	+	.	2
<i>Salix appendiculata</i>	E1	+	.	.	1
<i>Salix glabra</i>	E1	+	.	.	1

Number of relevé (Zaporedna številka popisa)		1	2	3	Pr.
TR	<i>Thlaspietea rotundifolii</i>				
	<i>Astrantia carniolica</i>	E1	+	+	3
	<i>Adenostyles glabra</i>	E1	2	.	2
	<i>Festuca nitida</i>	E1	1	.	2
	<i>Minuartia austriaca</i>	E1	+	.	2
	<i>Rhodiola rosea</i>	E1	+	.	2
	<i>Gymnocarpium robertianum</i>	E1	+	.	2
	<i>Biscutella laevigata</i>	E1	.	+	1
	<i>Hieracium bifidum</i>	E1	.	+	1
	<i>Valeriana montana</i>	E1	.	+	1
TG	<i>Trifolio-Geranietea</i>				
	<i>Pulmonaria australis</i>	E1	+	1	3
PaT	<i>Poo alpinae-Trisetalia</i>				
	<i>Ranunculus nemorosus</i>	E1	+	.	2
	<i>Poa alpina</i>	E1	.	.	1
EP	<i>Erico-Pinetea</i>				
	<i>Rubus saxatilis</i>	E1	1	1	3
	<i>Bupthalmum salicifolium</i>	E1	1	1	3
	<i>Rhodothamnus chamaecistus</i>	E1	+	+	3
	<i>Rhododendron hirsutum</i>	E2a	1	3	2
	<i>Erica carnea</i>	E1	.	2	2
	<i>Cirsium erisithales</i>	E1	+	+	2
	<i>Valeriana tripteris</i>	E1	+	.	2
	<i>Calamagrostis varia</i>	E1	1	.	1
	<i>Carex ornithopoda</i>	E1	.	.	1
VP	<i>Vaccinio-Piceetea</i>				
	<i>Luzula sylvatica</i>	E1	1	+	2
	<i>Homogyne sylvestris</i>	E1	+	+	2
	<i>Larix decidua</i>	E2a	+	.	1
	<i>Larix decidua</i>	E2b	.	+	1
	<i>Solidago virgaurea</i>	E1	r	.	2
	<i>Picea abies</i>	E2b	.	+	1
	<i>Vaccinium myrtillus</i>	E1	+	.	1
	<i>Polystichum lonchitis</i>	E1	+	.	1
	<i>Clematis alpina</i>	E1	.	.	1
FS	<i>Fagetalia sylvaticae</i>				
	<i>Daphne mezereum</i>	E1	+	.	1
	<i>Helleborus niger</i>	E1	.	+	1
	<i>Knautia drymeia</i>	E1	1	1	3
	<i>Lilium martagon</i>	E1	+	.	1
	<i>Melica nutans</i>	E1	1	.	2
	<i>Mercurialis perennis</i>	E1	1	1	3
	<i>Paris quadrifolia</i>	E1	+	.	1
	<i>Dryopteris filix-mas</i>	E1	+	.	1
QF	<i>Quercu-Fagetea</i>				
	<i>Anemone nemorosa</i>	E1	+	.	1
	<i>Hepatica nobilis</i>	E1	+	.	1
	<i>Listera ovata</i>	E1	+	.	1
AT	<i>Asplenietea trichomanis</i>				
	<i>Paederota lutea</i>	E1	.	+	2
	<i>Asplenium viride</i>	E1	.	+	2
	<i>Cystopteris fragilis</i>	E1	.	+	1
	<i>Bupleurum petraeum</i>	E1	.	.	1
M	Mosses (Mahovi)				
	<i>Marchantia polymorpha</i>	E0	+	.	1

A Limestone (apnenec)
 Gr Gravel (grušč)
 R Rendzina (rendzina)
 FB *Festuco-Brometea*
 SCf *Scheuchzerio-Caricetea fuscae*

Table 2: Groups of diagnostic species in the communities with *Pulmonaria australis* in the Triglav Mountains
 Preglednica 2: Skupine diagnostičnih vrst v sestojih z vrsto *Pulmonaria australis* v Triglavskem pogorju

Phytosociological group (Fitocenološka skupina)	Nr.	%
<i>Elyno-Seslerietea</i>	49	35,25
<i>Mulgedio-Aconitetea, Betulo-Alnetea</i>	27	19,42
<i>Thlaspietea rotundifolii</i>	16	11,51
<i>Trifolio-Geranietea</i>	3	2,16
<i>Poo alpinae-Trisetalia</i>	3	2,16
<i>Erico-Pinetea</i>	19	13,67
<i>Vaccinio-Piceetea</i>	12	8,63
<i>Quercu-Fagetea</i>	3	2,16
<i>Asplenietea trichomanis</i>	6	4,32
Mosses (Mahovi)	1	0,72
Total (Skupaj)	139	100