

Dry grasslands on fluvial terraces of the middle reaches of river Piave in the North East Italy

Stefano Tasinazzo¹ , Katia Zanatta² , Cesare Lasen³  & Roberto Fiorentin⁴ 

Key words: *Astragalo onobrychidis-Koelerietum pyramidatae, Chamaecytiso hirsuti-Chrysopogonetum grylli*, Po plain, river Brenta, *Scorzoneretalia villosae*, Veneto.

Ključne besede: *Astragalo onobrychidis-Koelerietum pyramidatae, Chamaecytiso hirsuti-Chrysopogonetum grylli*, Padska nižina, reka Brenta, *Scorzoneretalia villosae*, Benečija.

Abstract

Dry grassland vegetation on fluvial terraces along middle reaches of river Piave and river Brenta in Northeastern high Po plain were investigated through a phytosociological approach. Comparisons with ecologically analogous communities described from neighbouring territories lead to the description of a new association belonging to *Centaureion dichroanthae* alliance: *Astragalo onobrychidis-Koelerietum pyramidatae*. Residual fragments of *Chrysopogon gryllus*-dominated grassland on slightly deeper soils are referred to *Chamaecytiso hirsuti-Chrysopogonetum grylli*.

Izvleček

S fitocenološkim pristopom smo preučili suha travniča na aluvialnih terasah vzdolž srednjega tek rek Piave in Brente v višje ležeči severozahodni Padske nižini. Primerjava z ekološko podobnimi združbami, opisanimi na sosednjih območjih, je pokazala, da smo opisali novo asociacijo, ki jo uvrščamo v zvezo *Centaureion dichroanthae* alliance: *Astragalo onobrychidis-Koelerietum pyramidatae*. Preostale fragmente travnič s prevladujočo vrsto *Chrysopogon gryllus*, ki uspevajo na malo globljih tleh, pa uvrščamo v asociacijo *Chamaecytiso hirsuti-Chrysopogonetum grylli*.

Corresponding author:

Stefano Tasinazzo
E-mail:
stefano.tasinazzo@gmail.com

Received: 12. 2. 2020

Accepted: 19. 6. 2021

¹ Vicenza, Italy.

² Giavera del Montello (TV), Italy.

³ Arson di Feltre (BL), Italy.

⁴ Vicenza, Italy.

Introduction

Rivers are the most fitting example of ecological corridor concept in nature conservation. They mitigate negative effects of increasing habitat fragmentation by connecting patches or territories with high natural values and allowing species movements (e.g. Beier & Noss, 1998; Gonzalez et al., 1998). In man-made landscapes, such as Po plain, rivers in themselves constitute core areas by hosting animal and plant populations and habitat which are absent in the poor countryside outside their banks (Buffa & Lasen, 2010). Hydraulic safety reasons, natural disturbances connected to the hydraulic regime and inherent constraints due to poor soils often allow the development of mainly herbaceous vegetation with scattered shrubs and trees or groves. Prevailing communities are dry grasslands, which have been recognized at grain sizes below 100 m² as the most species-rich habitat worldwide (Wilson et al., 2012). On the other hand, they are assessed in Europe as the most threatened terrestrial and freshwater habitats, excluding mires and bogs, primarily because of the abandonment of traditional managements (Janssen et al., 2016). This concerns secondary grasslands, i.e. anthropogenic communities at sites whose potential vegetation is forest, whereas herbaceous vegetation investigated in the present work at least in part can be considered as extrazonal, spreading within forest

biomes but in too dry and shallow soils to support tree growth (Dengler et al., 2014).

As many long stretches of northeastern Italian rivers, also the middle river Piave (SAC IT3240023 – Grave del Piave) and Brenta (SCI-SAC IT3260018 – Grave e zone umide della Brenta) are protected in EU under Council Directives 92/43/EEC and/or 79/409/EEC. Military bondages have for a long-time ensured the preservation of some dry grassland surfaces along the Piave, as in Friuli (Nimis & Fonda, 1997), by avoiding land-use exploitations which are detrimental to natural environment (gravel extraction, vineyard planting etc.). A planned project of an overflow basin to retain flood waters could today compromise the preservation of the little but meaningful fragments remaining, contrasting with the principles of the EU environmental legislation (Habitats Directive). Hence, this contribution runs the risk of being reduced to mere documentary material before the complete disappearance of such natural emergencies, as has been advanced in recent years in Friuli-Venezia Giulia for some meadows and grasslands communities (Poldini & Oriolo, 1994).

Notwithstanding their naturalistic relevance, no analytical phytosociological study was carried out nor on Piave either along near Brenta Grave, where dry communities are very rare and more impoverished. Recently, the Grave di Ciano, in the core of Piave study area, was designated as a wilderness area (Zunino, 2018). On the basis of this



Figure 1: The landscape in the middle course of river Piave with the Venetian Prealps in the background. The investigated dry grasslands in the foregrounds.

Slika 1: Pokrajina v srednjem teku reke Piave z Beneškimi Predalpami v ozadju. Preučevana travnišča so v ospredju.

institution, the only available to date vegetation paper was published, in order to produce a vegetation map not supported by relevés (Pedrotti & Murra, 2020).

The relationship of Venetian terrace river grasslands with ecologically close communities described from neighbouring Friulian territories were tested by means of a phytosociological approach. In fact, Friuli-Venezia Giulia has a well-established tradition of vegetation studies concerning xeric grasslands on gravel bed deposited by pre-Alpine rivers (Zenari, 1928; Pignatti 1952; Lorenzoni, 1964; Lorenzoni, 1967; Feoli Chiapella & Poldini, 1994).

The aim of this research is to clarify floristic composition, structure and syntaxonomy of residual dry grass-

lands occurring in middle reaches (locally *grave*) of river Piave, before their possible disappearance or degradation due to human pressures (Figure 1). Observations have been partly extended to river Brenta in order to make the study more exhaustive.

Study area

The study area encompasses the middle courses of the river Piave and, to a lesser extent, of the river Brenta in the eastern high Venetian plain (western sector of the Venetian–Friulian plain, North East Italy; Figure 2).




Figure 2: Original relevés (●) in the middle reaches of river Brenta (A) and river Piave (B) in the eastern high Venetian plain. Also published relevés (■) from Natisone river (Lorenzoni, 1964) are given (see text).

Slika 2: Izvirni popisi (●) v srednjem teku rek Brente (A) in Piave (B) v višje ležeči severozahodni Padske nižini. Prikazani so tudi že objavljeni popisi (■) ob reki Natisone/Nadiža (Lorenzoni 1964) (glej besedilo).

The high Venetian-Friulian plain area is bounded to the north by the pre-Alpine chain, whereas the southern limit fades with the resurgence belt about 30 km south of the Alpine foothills. Large portions of the northeastern Po plain formed during the Last Glacial Maximum on late Pleistocene (12.000 b.p.). This period featured great sedimentary activity of the Adige, Brenta, Piave and Tagliamento Rivers, which received fluvio-glacial deposits from the eastern Alps glaciers and formed telescopic coalescent alluvial megafans (Fontana et al., 2014). These deposits are composed of gravels and small rocks, locally mixed with sands, according to a depositional model linked to a braided watercourse that alternates phases of flood, with huge solid transport, with lean phases in which it flows below the very permeable alluvial mattress. The gravel bedrock typically varies on size and composition: small-stone and gravel prevalence in the foothill sector, silt and clay widespread presence in correspondence of the resurgence belt (Michelutti et al., 2003). The lithological types reflect the prevalently carbonate nature of the mountain basins to which the two rivers belong. The Piave drains large part of the Dolomites, and its sediments are 50–70% carbonate, with a significant magmatic component, while Brenta sediments are 20–35% carbonate and have a high content of porphyric rocks (Fontana et al., 2014). Therefore, in the study area substrata are constituted by gravels and few, very permeable sand lenses. Currently, the Piave flows between 10 m high erosion scarps at Grave di Ciano, in

its medium course, where the riverbed is 3 km wide and includes some large fluvial islands.

Climatic parameters and bioclimatic indices of the study area are based on thermopluviometric data of Cittadella (48 m a.s.l., for river Brenta) and Volpago del Montello (94 m a.s.l., for river Piave) stations (<https://www.arpa.veneto.it>). Both stations, 45 km away, are in high Po plain. The mean annual temperature is 13.5 °C and the mean annual rainfall is 1180 mm. According to the ‘Worldwide Bioclimatic Classification System’ (Rivas-Martinez & Rivas-Saenz, 1996–2020), the study area is in a transition region between the temperate continental bioclimate, subtype weak subcontinental (Cittadella thermopluviometric station) and the temperate oceanic bioclimate, subtype strong semicontinental (Volpago del Montello thermopluviometric station); the thermotype is upper mesotemperate, the ombrotype lower humid (Figure 3). The humid trait of the bioclimate is mitigated by the porosity of the gravelly soil, which determines a rapid percolation of the rainwater with consequent edaphic aridity.

As regards landscape, Piave and Brenta riverbeds constitute the geosigmetum of glareicolous, lowland and hydrophilous riverbed vegetation of the high Venetian plain, which includes xeric grasslands (sub *Saturejion subspicatae*) on third-level terraces (Pedrotti & Murrja, 2020). In these consolidated and exceptionally flooded terraces, the original vegetation-plot records were carried out.




Figure 3: Climograms of Cittadella (PD) and Volpago del Montello (TV) thermopluviometric stations.
Slika 3: Klimadiagram za klimatski postaji Cittadella (PD) in Volpago del Montello (TV).

Materials and Methods

A total of 36 phytosociological relevés was carried out, 32 on river Piave terraces and 4 on the river Brenta terraces (Figure 2), ranging from 45 m to 140 m a.s.l. Data were collected according to Braun-Blanquet (1964) methodology in summer 2020, except for 5 stands, 2 dating back to 1994, one to 2012 and 2 to 2018. The modified Braun-Blanquet scale was used in 2020 vegetation plot records as regards '2a' and '2b' cover values (Barkman et al., 1964).

Comparisons with published data were restricted to materials describing communities sharing common ecological, altitudinal and phytogeographic traits. In particular, the following data was selected from literature:

- *Centaureo dichroanthae-Globularietum cordifoliae* (Tab. 1: rel. 1–14, in Feoli Chiapella & Poldini (1994))
- *Saturejo variegatae-Brometum condensati* (Tab. 2: rel. 1–18, in Feoli Chiapella & Poldini (1994); they include 3 relevés which Lorenzoni (1964) had attributed to *Astragalo-Stipetum*)
- *Saturejo variegatae-Brometum condensati* (Tab. 3: rel. 29–39, in Tasinazzo (2001))
- *Saturejo variegatae-Brometum condensati* (Tab. 1: rel. 2-3-4-20-21-22-26-27, in Lasen (1995))
- *Bromo condensati-Stipetum eriocaulis* (Tab. 1: rel. 5-6-7-8-10-11-12-13-14-15-16-17-18-19-33, in Lasen (1995))
- *Bromo condensati-Stipetum eriocaulis* (Tab. 1: tipologia 1, in Scortegagna & Curti (2000))
- *Schoeno nigricantis-Chrysopogonetum grylli* (Tab. 4: rel. 1–10, in Feoli Chiapella & Poldini (1994))
- *Chamaecytiso hirsuti-Chrysopogonetum grylli* (Tab. 5: rel. 1–7, in Feoli Chiapella & Poldini (1994))
- *Onobrychido arenariae-Brometum erecti avenuletosum pubescens* var. *a Chrysopogon gryllus* (Tab. 6: rel. 1–7, in Feoli Chiapella & Poldini (1994))
- *Astragalo onobrychidis-Artemisietum albae* (Tab. 28: rel. 1–17, in Biondi et al. (1997))

At first, binary (presence/absence) data were employed in multivariate analyses, and in a second step computation of similarity was indirectly tested from the site scores on the ordination axis of Principal Component Analysis (Jongman et al., 1995), obtained from a cover data matrix. As regards the latter and the classification of original relevés, before to perform cover data analysis, values were converted to van der Maarel (1979) ordinal scale. Also percentage cover values were used in multivariate analyses, adopting the approximations given in Gigante et al. (2012; Fig. 1: Braun Blanquet mod. by Barkman et al., 1964) and after square-root-transformation of input data.

All multivariate analyses were conducted with Syn-Tax 2000 package (Podani, 2001).

Due to difficult and often controversial determinations and to overcome possible bias in analysis results, some elementary taxa were grouped prior of analyses (e.g. *Koeleria pyramidata* aggr., *Thymus serpyllum* aggr., *Centaurea scabiosa* s.l.).

Biological and chorological spectra were weighted using percentage cover values with above approximations. Ecological indicator values and chorotypes (except for *Bromopsis condensata*, east-Alpic endemic instead of endemic) are in accordance with Pignatti (2005). For the comparison analyses, chorotypes were grouped (see Appendix 3).

The nomenclature of vascular plant species follows Bartolucci et al. (2018). The diagnostic species of alliance, suborder, order and class *Festuco-Brometea* are according to Terzi (2015); some integrations were derived from Mucina et al. (2016) for the *Festuco-Brometea* class. *Bromopsis condensata*, that in Terzi (2015) was not separated from *B. erecta*, and *Carex lyparocarpus*, absent in the same author's analysis, were considered diagnostic of *Centaureion dichroanthae* alliance (for the latter see also Feoli Chiapella & Poldini, 1994). *Campanula sibirica*, also lacking in Terzi (2015), was included amongst order diagnostic species (Sburlino et al., 2008). Bryophytes and lichens were disregarded. New syntaxon names are in accordance with the rules of the International Code of Phytosociological Nomenclature (Theurillat et al., 2020).

Results and Discussion

The binary numerical classification proved the independence of original relevés with respect to other stands and coenosis coming from neighbouring and ecologically comparable north-eastern Italian territories (cluster A in Figure 4). Only Lorenzoni's relevés (1964) from the river Natisone in Friuli-Venezia Giulia joined the cluster grouping the Venetian river terrace stands. According to Feoli Chiapella & Poldini (1994) these 3 relevés merge in *Saturejo variegatae-Brometum condensati* identifying within it the subass. *astragaletosum onobrychidis*, whereas Lorenzoni (1964) had attributed them to the independent coenosis *Astragalo-Stipetum pennatae*. Table of *Saturejo-Brometum* from Friuli includes also 3 Pignatti's relevés (subass. *epilobietosum dodonaei*) reported by the author sub "Aggr. ad *Artemisia campestris* ed *Epilobium dodonaei*" and referred to *Epilobion fleischeri* (Pignatti, 1952). On the basis of the overall floristic affinity, Feoli Chiapella & Poldini (1994) address the issue and attribute the Pignatti's relevés to *Saturejo-Brometum*. The ambiguity emerged again in our

analysis which enucleates these stands in a cluster totally separated not only from *Saturejo-Brometum* but also from the other compared xeric censes (cluster I in Figure 4). Our results are in some way in accordance with the initial Pignatti's interpretation (1952), dealing with 'a vegetation of sandy and stony gravel bed', but further focused studies may provide insights into the referring class of these *Festuco-Brometea*-species poor vegetation. *Artemisia*

campestris constitutes together with *Epilobium dodonaei* also on gravel bed of river Brenta a glareicolous community with strong pioneer traits on substrates that are not fully consolidated and subject to episodic flooding. The typical aspect of *Saturejo-Brometum* coincides with subass. *seslerietosum albantis* containing the holotypus (Feoli Chiapella & Poldini, 1994) and which merges with other relevés from Grappa Massif (cluster D in Figure 4).




Figure 4: Classification of original and Lorenzoni's relevés (A) in relation to other *Scorzoneretalia* relevés from NE-Italy (simplified dendrogram; average link, Jaccard index; binary data). B: stands from Berici Hills; C: *Schoeno-Chrysopogonetum* + *Chamaecytiso-Chrysopogonetum* + *Onobrychido-Brometum* + 1 rel. of *Saturejo-Brometum*, all from Friuli-Venezia Giulia; D: *Saturejo-Brometum* from Grappa Massif and Friuli-Venezia Giulia (including holotypus) + 3 rel. of *Bromo-Stipetum* from M. Summano; E: *Bromo-Stipetum* from Grappa Massif and M. Summano; F: *Centaureo-Globularietum*; G: 1 rel. of *Saturejo-Brometum* from Friuli-Venezia Giulia; H: 3 rel. of *Saturejo-Brometum* from Friuli-Venezia Giulia; I: 3 rel. of *Saturejo-Brometum* from Friuli-Venezia Giulia (corresponding to '*Artemisia campestris* ed *Epilobium dodonaei*' aggr. in Pignatti (1952)).

Slika 4: Klasifikacija izvirovih in Lorenzonijevih popisov (1964) (A) v primerjavi z ostalimi popisi reda *Scorzoneretalia* iz severovzhodne Italije (poenostavljen dendrogram; povprečna povezava, Jaccardov indeks; binarni podatki). B: stoji z gričevja Berici; C: *Schoeno-Chrysopogonetum* + *Chamaecytiso-Chrysopogonetum* + *Onobrychido-Brometum* + 1 popis *Saturejo-Brometum*, vsi iz Furlanije-Julijске krajine; D: *Saturejo-Brometum* z masivo Grappa in Furlanije-Julijске krajine (vključno s holotipom) + 3 popisi *Bromo-Stipetum* z gore Summano; E: *Bromo-Stipetum* z masivo Grappa in gore Summano; F: *Centaureo-Globularietum*; G: 1 popis *Saturejo-Brometum* iz Furlanije-Julijске krajine; H: 3 popisi *Saturejo-Brometum* iz Furlanije-Julijiske krajine; I: 3 popisi *Saturejo-Brometum* iz Furlanije-Julijiske krajine (ustreza '*Artemisia campestris* ed *Epilobium dodonaei*' aggr. iz Pignatti (1952)).




Figure 5: Classification of PCA coordinates of original relevés (□; this study), Lorenzoni's relevés (■; Lorenzoni, 1964), *Schoeno-Chrysopogonetum* + *Chamaecytiso-Chrysopogonetum* (○; Feoli Chiapella & Poldini 1994) and *Centaureo-Globularietum* (●; Feoli Chiapella & Poldini 1994) from Friuli-Venezia Giulia (cover data matrix).

Slika 5: Klasifikacija PCA koordinat izvirovih popisov (□; ta raziskava), Lorenzonijevih popisov z reke Natisone/Nadiže (■; Lorenzoni, 1964), *Schoeno-Chrysopogonetum* + *Chamaecytiso-Chrysopogonetum* (○; Feoli Chiapella & Poldini 1994) in *Centaureo-Globularietum* (●; Feoli Chiapella & Poldini 1994) iz Furlanije-Julijiske krajine (matrika podatkov s pokrovnostjo).

Indeed, according to the itself authors' statement the original description of *Saturejo-Brometum* is representative of a very articulated coenosis, appearing very heterogeneous also to our analysis.

Owing to the impossibility of ascribing original relevés to *Saturejo-Brometum* or to other hilly and submontane already typified coenosis, a close examination involved only ecologically comparable communities (known as 'magredi' grasslands) scattered on alluvial fans covering northern Friulian plain between 50 and 300 m elevation: the most pioneer stage *Centaureo-Globularietum*, the intermediate successional coenosis *Schoeno-Chrysopogonetum* and the most mature stage *Chamaecytiso-Chrysopogonetum*. The resulting classification of PCA coordinate values better describes evidences for observed ecological differences amongst original relevés (Figure 5). The Venetian stands are divided into two main groups: a numerically prevalent one on more primitive soils constitutes an independent cluster (A) including the already mentioned relevés from Natisone, the minority one (B1) on more well-developed soils merges with the other Friulian stands and particularly with *Schoeno-Chrysopogonetum* and *Chamaecytiso-Chrysopogonetum* (B2); *Centaureo-Globularietum* forms an independent cluster (C).

Astragalo onobrychidis-Koelerietum pyramidatae

As regards the cluster A, the framing in *Festuco-Brometea* class (Table 1) is clearly highlighted by a rich content of high frequency species often occurring also with high cover values, such as *Koeleria pyramidata*, *Helianthemum nummularium* subsp. *obscurum*, *Fumana procumbens* and *Thymus pulegioides* (V frequency class). Its membership in steppic submediterranean pastures of the SE-European–Illyrian order *Scorzoneretalia villosae* is clear despite it appears weak due to the marginality of the study area with respect to the distribution area of the order itself. In fact, character species are well-represented only in the relevés coming from eastern Friuli (cf in Figure 5; rel. 29–31 in Table 1), whereas they decrease sharply in westernmost surveys along the Brenta in Veneto (rel. 1–4 in Table 1). The most common diagnostic species of the order include *Plantago holosteum* and *Chrysopogon gryllus* (III frequency class) (Table 1). More evident is the framing in the lower syntaxonomical level. The presence, often with high frequency and cover values, of *Potentilla pusilla*, *Carex liparocarpos*, *Bromopsis condensata*, *Cytisus purpureus* assures the attribution to *Centaureion dichroanthae*, alliance encompassing prealpic submediterranean grasslands on shallow soils (Mucina et al., 2016). With regard to the association rank, because of phytogeographical reasons it is not possible to refer the vegetation plot records to *Astragalo-Stipetum pennatae*, as proposed by Lorenzoni

(1964). *Astragalo-Stipetum* is a community described for continental inner Alps and distinguished by mostly continental species such as *Poa perconcinna*, *Koeleria vallesiana*, *Minuartia rostrata*, *Androsace septentrionalis*, *Oxytropis halleri/velutina*, *Stipa capillata* etc. It belongs to steppic rocky grasslands of deep intramontane valleys of the Alps (*Stipo-Poion xerophilae*). Floristic composition of *Astragalo-Koelerietum* attributes unique traits with respect to the other *Scorzoneretalia* associations so far described. The comparison by means of chorological groups, extended to *Astragalo onobrychidis-Artemisietaum albae* of fluvial terraces of the Taro flowing in low Po plain in the Apennine area, shows closer relationship with Friulian *Centaureo-Globularietum* (Figure 6) as consequence of higher rates of orophytes attesting a marked primitiveness of the substratum. Lower percentages of the eurasian group constitute additional reason for their closeness (Table 2). *Centaureo-Globularietum* belongs to *Centaureion* alliance too, whereas *Schoeno-Chrysopogonetum* and *Chamaecytiso-Chrysopogonetum* are attributed to *Scorzonerion villosae* alliance (Terzi, 2015).




Figure 6: PCA of dry grassland of northeastern Italian plain (●) in relation to chorotype groups (□). First axis accounts for 48.9% of total variance, second axis for 31.6% (square-root-transformed percentage cover data). Community abbreviations as in Table 2, chorotype group as in Appendix 3.

Slika 6: PCA suhih travšč iz severovzhodne italijanske nižine (●) v odnosu do horoloških skupin (□). Prva os pojasni 48,9% skupne variancije, druga pa 31,6% (pokrovnost transformirana s korenjenjem). Okrajšave združb so enake kot v Tabeli 2, horološke skupine pa v Prilogi 3.

At a more detailed examination, the new community differs from *Centaureo-Globularietum* in lacking both all the character species of this association (*Euphorbia triflora/kernerii*, *Matthiola fruticulosa/valesiaca*, *Brassica glabrescens* and *Crambe tataria*) and many significant and often very frequent alliance species such as *Centaurea dichroantha*, *Carex mucronata*, *Sesleria caerulea*, *Polygala forojulensis*, *Lomelosia graminifolia*, *Hieracium porrifolium* ecc. Many of these are endemic elements (*Euphorbia triflora/kernerii*, *Matthiola fruticulosa/valesiaca*, *Brassica glabrescens*, *Centaurea dichroantha*) contributing to mark the absolute phytogeographic originality of the coenosis and more generally of the high Friulian Plain where it thrives (e.g. Poldini, 1973). From the synecological point of view Lorenzoni (1967) already outlined the independence of the two coenosis, however occurring in closely contact in Friuli-Venezia Giulia. He observed the hypothesized *Astragalo-Stipetum* on consolidated gravel deposits with initial humus accumulation, thriving between *Centaureo-Globularietum* closer to river bed where there are no evidence of organic matter and *Chrysopogonetum* s.l. lying on well-developed, decalcified soils. Ordination through soil indicator values confirms author's observations (Figure 7). *Centaureo-Globularietum* and *Astragalo-Koelerietum* share the presence of *Gypsophila repens*, a de-Alpine entity ingressive from *Thlaspietea*, that Feoli Chiapella & Poldini (1994) consider differential of the former. Thereby, this glareophyte should be regarded as diagnostic of *Centaurenion*. The exclusive entities *Potentilla pusilla*, *Astragalus onobrychidis* and *Erucastrum nasturtiifolium* may be considered the good differential entities of the new community (Table 3). *Astragalus onobrychidis* is also included amongst differential species of *Astragalo-Artemisietum*, a coenosis




Figure 7: PCA of dry grassland of northeastern Italian plain (■) in relation to soil indication values (○). M: moisture, N: nitrogen, R: reaction. First axis accounts for 72.9% of total variance, 2^o axis: 24.9%; weighted cover data). Community abbreviations as in Table 2. **Slika 7:** PCA suhih travničkih zajednica (■) v odnosu do indikatorskih vrednosti tal (○). M: vlažnost, N: dušik, R: reakcija. Prva os pojasnji 72,9% skupne variancije, druga os: 24,9%; tehtani podatki pokrovnosti). Okrajšave združb so enake kot v Tabeli 2.

very rich in *Festuco-Brometea* species, but lacking in diagnostic species of *Scorzoneralia* and lower syntaxonomical rank levels. In any case, it is attributed to *Ononido-Rosmarineta* (Biondi et al., 1997).

On the basis of the above, the relevés in Table 1 may relate to a new association, named *Astragalo onobrychidis-Koelerietum pyramidatae*.

Astragalo onobrychidis-Koelerietum pyramidatae ass. nova hoc loco (holotypus rel. 8 in Table 1)

Differential species. *Potentilla pusilla*, *Astragalus onobrychidis* and *Erucastrum nasturtiifolium*

Floristic composition. It is a discontinuous para-steppic dry grassland with largely dominant hemicryptophytes, with *Stipa eriocaulis* often structuring the herbaceous landscape (Figure 8). *Stipa eriocaulis*, *Koeleria pyramidata*, *Bromopsis condensata*, *Oreoselinum nigrum* and *Centaurea scabiosa* are the tall constant grasses and forbs forming the sparse high vegetation cover. The richer lower layer includes *Potentilla pusilla*, *Helianthemum nummularium/obscurum*, *Fumana procumbens*, *Thymus pulegioides*, *Globularia bisnagarica*, *Sanguisorba minor*, *Carex lippocarpos*, *Astragalus onobrychidis* etc. among the entities more frequent or abundant.

A cryptogamic layer is always present and sometimes significant in terms of coverage.

Life forms and chorotypes. Hemicryptophytes 72.2%, chamaephytes 19.5%, geophytes 5.3%, therophytes 2.3%, nanophanerophytes 0.2%, phanerophytes 0.5%.

Synecology. The coenosis thrives on consolidated gravel and pebble sediments, on flat ground or secondly gentle slopes, of river terraces, where it occupies Calcixerpts-Haploxerpts soils (Soil Survey Staff 2015). The small share of wide-ranging species agrees with initial stages of colonization. Dryness and stoniness of the habitat are also underlined by the occurrence of species ingressive from *Thlaspietea rotundifolii* (*Gypsophila repens*, *Erucastrum nasturtiifolium*).

Ecological spectrum. Light 5.07, temperature 5.89, continentality 5.82, moisture 2.91, soil reaction 5.95, nitrogen 2.20.

Syntaxonomy. The cluster analysis allows us to recognise three different aggregation types (Figure 9). The rel. 8–31 represent the typical one thriving on consolidated terraces of river Piave, with the exception of the last three stands collected along river Natisone by Lorenzoni (1964). Only rel. 8–31 include *Gypsophila repens* and *Erucastrum nasturtiifolium*. Diagnostic taxa of *Centaureion* alliance, *Koelerienalia* suborder and *Scorzoneralia* order are here numerically well represented. Here *Stipa eriocaulis* occurs, often attaining high cover values, in a such way to outline



Figure 8: The new association *Astragalo onobrychidis-Koelerietum pyramidatae* in the Grave di Ciano in the middle course of river Piave.
Slika 8: Nova asociacija *Astragalo onobrychidis-Koelerietum pyramidatae* v kraju Grave di Ciano v srednjem teku reke Piave.




Figure 9: Dendrogram of relevés of Table 1 (algorithm: average link, similarity ratio; cover data).

Slika 9: Dendrogram popisov iz Tabele 1 (algoritam: povprečna povezava, delež podobnosti; podatki s pokrovnostjo).

the structure of the habitat. The rel. 1–4, from terraces of river Brenta, are the most species-poor stands, with very scarce occurring alliance, suborder and order diagnostic species and with an important bryophyte cover. *Artemisia campestris* differentiate this western provenance. Habitat

appears highly localised, very fragmented and disappearing, so relevés are in partly degraded conditions. Further research can clarify the possibility of their framing in different vegetation types; at present, we think that the synecology and the occurrence of two differential species justifies their inclusion in *Astragalo-Koelerietum*. The remaining relevés (5–7) are representative of a degraded aspect differentiated by large cover of *Bothriochloa ischaemum* a species with apophyte traits. These stands are also liable to belong to another vegetation type due to the lacking of the greater part of the most significative species.

Synchorology. The community occurs on the middle reaches (grave) of rivers Piave and, partly, Brenta in Veneto, but according to Lorenzoni's observations (1964, 1967) its distribution area extended up to eastern Friuli-Venezia Giulia across rivers Tagliamento, Torre and Natisone. It is likely that it might occur also on river Isonzo, but all Friulian data need confirmation (see Appendix 4).

Chorological spectrum. Atlantic 25.4, Eurasiac 24.4, orophytes 14.8, Eastern 13.5, Mediterranean 9.4, Cosmopolitan 8.4, Boreal 2.2, alien 1.8, Endemic 0.1.

Natura 2000. 62A0

Chamaecytiso hirsuti-Chrysopogonetum grylli

Also original relevés in cluster B1 (Figure 5) can be easily attributed to *Festuco-Brometea* and *Scorzoneretalia villosae*, but inclusion in lower syntaxonomical levels is not equally clear. Stands were gathered on middle reaches of the river Piave, on residual fragments of well-developed soils representing higher terraces. The community is characterised by the large predominance of *Chrysopogon gryllus*, in place of *Stipa eriocalis*, but this vicariance is not associated with the occurring of *Hypocharidenion* and *Scorzonerion* species as it might be expected, diagnostic taxa of *Centaureion* prevail (Table 4). Small size patches are responsible for a reduction in vegetation homogeneity due to a mass effect induced by wider shallow-soil surrounding areas that favour the penetration of species better-tolerating more severe dryness conditions. Nevertheless, a set of mesophilous species enter the community by acting as good indicators of improved growth conditions (*Dactylis glomerata*, *Bromus erectus*, *Salvia pratensis* etc.). Despite the uncertain alliance position and following the cluster analysis outcome (Figure 5), the most viable option was the comparison with similar *Chrysopogon gryllus*-rich associations described from the high plain of Friuli-Venezia Giulia: *Schoeno-Chrysopogonetum* and *Chamaecytiso-Chrysopogonetum*, belonging to *Hypocharidenion* suballiance. The hypothesis to refer relevés to *Onobrychido-Brometum avenuletosum pubescens* var. *a Chrysopogon gryllus* was excluded by initial step of the analysis. Our relevés fit better with *Chamaecytiso-Chrysopogonetum* on the basis of a common chorological framework in which the high incidence of the eurasian group stands out (Figure 4). This is in line with Feoli Chiapella & Poldini (1994) postulate's according to which its distribution area should go beyond the western regional borders. Hence, relevés in Table 4 are referred to *Chamaecytiso-Chrysopogonetum*. Venetian material lacks the differential species of association reported by the authors for Friulian territory. However, *Anacamptis coriophora* shows its ecological optimum in this association occurring almost exclusively within it in Venetian stands and in no other dry semi-natural grassland community outside this one in Friuli-Venezia Giulia (Feoli Chiapella & Poldini, 1994). So, in our opinion, this species may be added to the differential entities of the association. *Cytisus pseudoprocumbens* and *Medicago prostrata* can be regarded as differential species of the Venetian provenance of the coenosis. This coenosis has suffered a drastic reduction in Friuli during last decades, because of intensive cultivation (maize). It appears highly likely that the coenosis was widespread outside the actual river banks of Piave, where now vineyards dominate the countryside.

Syntaxonomic scheme

Festuco-Brometea Br.-Bl. et Tx. ex Soó 1947

Scorzoneretalia villosae Kovačević 1959

Koelerienalia splendentis (Horvatić 1973) Terzi 2015

Centaureion dichroanthae Pignatti 1952

Astragalo onobrychidis-Koelerietum pyramidaliae ass. nova hoc loco (holotypus: Table 1, rel. 8) (syn. *Astragalo-Stipetum pennatae* Br.-Bl. 1961 sensu Lorenzoni 1964)

Scorzonerion villosae Horvatić ex Kovačević 1959

Hypochoeridenion maculatae Poldini & Feoli Chiapella ex Terzi 2011

Chamaecytiso hirsuti-Chrysopogonetum grylli Pignatti ex Feoli Chiapella & Poldini 1994

Conclusions

The present research from gravel-bed rivers Piave and Brenta in north-eastern Italian plain resulted in the identification of an already described coenosis and in the formalization of a new dry grassland association. The new knowledge provides a valuable tool for the geographical demarcation of the distribution area of orders *Scorzoneralia villosae*, *Brachypodietalia pinnati* and *Festucetalia valesiacae*, in a biogeographical context where the limits of these syntaxonomic levels have not yet been well clarified. The reported *Chamaecytiso-Chrysopogonetum* and *Astragalo-Koelerietum* are xeric coenosis hosting many grassland specialists. They belong to Palearctic grasslands, a series of habitat well known for their high biodiversity, but also for their strongly decline during last century due to practice intensification in agriculture, natural succession, artificial afforestation, transformation into arable lands, quarry opening (Dengler et al., 2014). The disappearance risk of most of remaining areas of *Chamaecytiso-Chrysopogonetum* and *Astragalo-Koelerietum* on gravel terraces of rivers Piave lies in the construction project of an overflow basin to retain flood waters. They both have to be referred to Habitat 62A0, are included in Natura 2000 network and deserve the attention of a society that wants to define itself cultured and civilised, at least for legal reasons. The very low rate of alien species underlines the good conservation status of the studied vegetation plot records, but the invasive *Amorpha fruticosa* spreading copiously along fluvial terraces threatens their biodiversity preservation. A regulated use of the most intact areas and the exclusion of land use changes as well as the maintenance of high proportion of ecologically valuable habitat in the surroundings (Janišová et al., 2014) could assure the conservation of these surviving communities.

Other syntaxa quoted in the text

- Astragalo onobrychidis-Artemisietum albae* Biondi, Vagge, Baldoni & Taffetani 1997
Astragalo onobrychidis-Stipetum pennatae Braun-Blanquet 1961
Bromo condensati-Stipetum eriocaulis Lasen ex Terzi 2015
Centaureo dichroanthae-Globularietum cordifoliae Pignatti 1952
Chamaecytiso hirsuti-Chrysopogonetum grylli Pignatti ex Feoli Chiapella & Poldini 1994
Onobrychido arenariae-Brometum erecti Poldini & Feoli Chiapella in Feoli Chiapella & Poldini 1994
Ononido-Rosmarinetea Br.-Bl. in A. Bolòs y Vayreda 1950
Saturejo variegatae-Brometum condensati Poldini & Feoli Chiapella in Feoli Chiapella & Poldini 1994
Schoeno nigricantis-Chrysopogonetum grylli Pignatti ex Feoli Chiapella & Poldini 1994
Stipo-Poion xerophilae Br.-Bl. et Richard 1950

Italy. *Plant Biosystems*, 152(2), 179–303. <https://doi.org/10.1080/11263504.2017.1419996>

Beier, P., & Noss, R. F. (1998). Do habitat corridors provide connectivity? *Conservation Biology*, 12(6), 1241–1252. <http://dx.doi.org/10.1111/j.1523-1739.1998.98036.x>

Buffa, G., & Lasen, C. (2010). *Atlante dei siti Natura 2000 del Veneto*. Regione del Veneto – Direzione Pianificazione Territoriale e Parchi.

Braun-Blanquet, J. (1964). *Pflanzensoziologie – Grundzüge der Vegetationskunde*. Springer Verlag.

Biondi, E., Vagge, I., Baldoni, M., & Taffetani, F. (1997). La vegetazione del Parco fluviale regionale del Taro (Emilia-Romagna). *Fitosociologia*, 34, 69–110.

Dengler, J., Janišova, M., Török, P., & Wellstein, C. (2014). Biodiversity of Palaearctic grasslands: a synthesis. *Agriculture Ecosystems & Environment*, 182, 1–14. <https://doi.org/10.1016/j.agee.2013.12.015>

Feoli Chiapella, L., & Poldini, L. (1994). Prati e pascoli del Friuli (NE Italia) su substrati basici. *Studia Geobotanica*, 13(1993), 3–140.

Gigante, D., Acosta, A. T. R., Agrillo, E., Attorre, F., Cambria, V. E., Casavecchia, S., Chiarucci, A., Del Vico, E., De Sanctis, M., Facioni, L., Geri, F., Guarino, R., Landi, S., Landucci, F., Lucarini, D., Panfili, E., Pesaresi, S., Prisco, I., Rosati, L., Spada, F., & Venanzoni, R. (2012). VegItaly: Technical features, crucial issues and some solutions. *Plant Sociology*, 49(2), 71–79. [10.7338/pls2012492/05](https://doi.org/10.7338/pls2012492/05)

Gonzalez, A., Lawton, J. H., Gilbert, F. S., Blackburn, T. M., & Evans-Freke, I. (1998). Metapopulation dynamics, abundance, and distribution in a microecosystem. *Science*, 281(5385), 2045–2047. <https://doi.org/10.1126/science.281.5385.2045>

Janišova, M., Michalcová, D., Bacaro, G., & Ghisla, A. (2014). Landscape effects on diversity of semi-natural grasslands. *Agriculture Ecosystems & Environment*, 182, 47–58. <http://dx.doi.org/10.1016/j.agee.2013.05.022>

Janssen, J. A. M., Rodwell, J. S., García Criado, M., Gubbay, S., Haynes, T., Nieto, A., Sanders, N., Landucci, F., Loidi, J., Ssymank A., Tahvanainen, T., Valderrabano, M., Acosta, A., Aronsson, M., Arts, G., Attorre, F., Bergmeier, E., Bijlsma, R.-J., Bioret, F., Bită-Nicolae, C., Biurrun, I., Calix, M., Capelo, J., Čarní, A., Chytrý, M., Dengler, J., Dimopoulos, P., Essl, F., Gardfjell, H., Gigante, D., Giusso del Galdo, G., Hájek, M., Jansen, F., Jansen, J., Kapfer, J., Mickolajczak, A., Molina J. A., Molnár, Z., Paternoster, D., Piernik, A., Poulin, B., Renaux, B., Schaminée, J. H. J., Šumberová, K., Toivonen, H., Tonteri, T., Tsiripidis, I., Tzanev, R., & Valachovič, M. (2016). *European Red List of Habitats. Part 2. Terrestrial and freshwater habitats*. Publications Office of the European Union.

Jongman, R. H. G., ter Braak, C. J. F., & van Tongeren, O. F. R. (1995). *Data analysis in community and landscape ecology*. Cambridge University Press.

Lasen, C. (1995). Note sintassonomiche e corologiche sui prati aridi del Grappa. *Fitosociologia*, 30, 181–199.

Lorenzoni, G. G. (1964). Un esempio di *Astragalo-Stipetum pennatae* Br.-Bl. 1961 nel Friuli nord-orientale. *Lavori di Botanica Istituto Botanico dell' Università Padova*, 23, 3–11.

Lorenzoni, G. G. (1967). Ricerche sui prati a “*Chrysopogon gryllus*” della pianura friulana. *“Udine”, Bollettino Biblioteca e Musei Civici e Biennali d'Arte Antica*, 4, 5–21.

Acknowledgements

We would like to thank Ballestrin S., Checuz R., Sebelin A. for their friendly joining during surveys and Ferrarese F. for valuable help in graphics; Checuz R. gave us the photographs included in the paper. We are grateful to dr. Vidali M. for supplying us with some bibliographic items. Prof. Pedrotti F. recently once again highlighted the naturalistic relevance of dry grasslands of the Piave. Two anonymous referees improved an early version of the manuscript. We thank the journal editorial team for translation into Slovenian.

Stefano Tasinazzo  <https://orcid.org/0000-0002-5829-0456>
Katia Zanatta  <https://orcid.org/0000-0002-9578-5802>
Cesare Lasen  <https://orcid.org/0000-0002-4182-7042>
Roberto Fiorentin  <https://orcid.org/0000-0003-3580-8985>

References

Barkman, J. J., Doing, H., & Segal, S. (1964). Kritische Bemerkungen und Vorschläge zur quantitativen Vegetationsanalyse. *Acta Botanica Neerlandica*, 13, 394–419. <https://doi.org/10.1111/j.1438-8677.1964.tb00164.x>

Bartolucci, F., Peruzzi, L., Galasso, G., Albano, A., Alessandrini, A., Ardenghi, N. M. G., Astuti, G., Bacchetta, G., Ballestri, S., Banfi, E., Barberis, G., Bernardo, L., Bouvet, D., Bovio, M., Cecchi, L., Di Pietro, R., Domina, G., Fascetti, S., Fenu, G., Festi, F., Foggi, B., Gallo, L., Gottschlich, G., Gubellini, L., Iamónico, D., Iberite, M., Jiménez-Mejías, P., Lattanzi, E., Marchetti, D., Martinetto, E., Masin R. R., Medagli, P., Passalacqua, N. G., Peccenini, S., Pennesi, R., Pierini, B., Poldini, L., Prosser, F., Raimondo, F. M., Roma-Marzio, F., Rosati, L., Santangelo, A., Scopolla, A., Scortegagna, S., Selvaggi, A., Selvi, F., Soldano, A., Stinca, A., Wagensommer, R. P., Wilhalm, T., & Conti, F. (2018). An updated checklist of the vascular flora native to

- Maarel, E. van der. (1979). Transformation of cover-abundance values in phytosociology and its effect on community similarity. *Vegetatio*, 39(2), 97–114.
- Michelutti, G., Zanolla, S., & Barbieri, S. (2003). *Suoli e paesaggi del Friuli Venezia Giulia. Pianura e colline del pordenonese*. Ersa.
- Mucina, L., Bültmann, H., Dierßen, K., Theurillat, J.-P., Raus, T., Čarni, A., Šumberová, K., Willner, W., Dengler, J., García, R. G., Chytrý, M., Hájek, M., Di Pietro, R., Iakushenko, D., Pallas, J., Daniëls, F. J. A., Bergmeier, E., Santos Guerra, A., Ermakov, N., Valachovič, M., Schaminée, J. H. J., Lysenko, T., Didukh, Y. P., Pignatti, S., Rodwell, J. S., Capelo, J., Weber, H. E., Solomeshch, A., Dimopoulos, P., Aguiar, C., Hennekens, S. M., & Tichý, L. (2016). Vegetation of Europe: hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *Applied Vegetation Science*, 19(suppl. 1), 3–264. <https://doi.org/10.1111/avsc.12257>
- Nimis, P. L., & Fonda, G. (1997). Phytogeography of parasteppeic vegetation in the high Friulian Plain (NE Italy). *Plant Ecology*, 132, 15–28.
- Pedrotti, F., & Murra, E. (2020). *Il paesaggio dell'area wilderness "Piave della Battaglia – settore Grave di Ciano" (Veneto)*. Tip. Ed. Temi.
- Pignatti, S. (1952). Introduzione allo studio fitosociologico della pianura veneta orientale. *Atti Istituto Botanico Pavia*, ser. 5, 11(1-3), 92–258.
- Pignatti, S. (2005). Valori di bioindicazione delle piante vascolari della flora d'Italia. *Braun-Blanquetia*, 39, 3–97.
- Podani, J. (2001). *Syn-Tax 2000. Computer program for data analysis in ecology and systematics. User's manual*. Scientia Publishing.
- Poldini, L. (1973). I magredi. *Informatore Botanico Italiano*, 5(2), 146–148.
- Poldini, L., & Oriolo, G. (1994). La vegetazione dei prati da sfalcio e dei pascoli intensivi (*Arrhenatheretalia* e *Poo-Trisetetalia*) in Friuli (NE Italia). *Studia Geobotanica*, 14(Suppl. 1), 3–48.
- Rivas-Martínez, S. (2008). *Global bioclimatics (Clasificación Bioclimática de la Tierra)*. Phytosociological Research Center. Retrieved January 15, 2021, from http://www.globalbioclimatics.org/book/bioc_global_bioclimatics-2008_00.htm
- Rivas-Martínez, S., & Rivas-Saenz, S. (1996–2020). *Worldwide Bioclimatic Classification System*. Phytosociological Research Center. Retrieved January 15, 2021, from <http://www.globalbioclimatics.org>
- Sburlino, G., Buffa, G., Filesi, L., & Gamper, U. (2008). Phytocoenotic originality of the N-Adriatic coastal sand dunes (Northern Italy) in the European context: The *Stipa veneta*-rich communities. *Plant Biosystems*, 142(3), 533–539. <https://doi.org/10.1080/11263500802410884>
- Scortegagna, S., & Curti, L. (2000). L'incespugliamento spontaneo dei prati aridi del Monte Summano (Prealpi Vicentine – Veneto). *Studi Trentini Scienze Naturali Acta Biologica*, 74(1997), 155–173.
- Soil Survey Staff. (2015). *Illustrated guide to soil taxonomy*. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center.
- Tasinazzo, S. (2001). I prati dei Colli Berici (Vicenza, NE Italia). *Fitosociologia*, 38(1), 103–116.
- Terzi, M. (2015). Numerical analysis of the order *Scorzoneretalia villosae*. *Phytocoenologia*, 45(1-2), 11–32. <http://dx.doi.org/10.1127/phyto/2015/0009>
- Theurillat, J. P., Willner, W., Fernández-González, F., Bültmann, H., Čarni, A., Gigante, D., Mucina, L., & Weber, H. (2020). International Code of Phytosociological Nomenclature. 4th edition. *Applied Vegetation Science*, 24, Article e12491. <https://doi.org/10.1111/avsc.12491>
- Wilson, J. B., Peet, R. K., Dengler, J., & Pärtel, M. (2012). Plant species richness: the world records. *Journal Vegetation Science*, 23, 796–802. <https://doi.org/10.1111/j.1654-1103.2012.01400.x>
- Zenari, S. (1928). La vegetazione dei 'magredi' nell'alta pianura del Friuli occidentale. *Atti Accademia Scientifica Veneto-Trentino-Istriana*, 19, 51–56.
- Zunino, F. (2018). Sul Fiume Piave di 100 anni fa designata la seconda Area Wilderness del Veneto! *Documenti Wilderness*, 33(4), 1.

Appendix 1:

Localities, dates and geographical coordinates (WGS84 system) of relevés of Table 1.

- Rel. 1: Ballin-Miotto (Fontaniva-PD), 9/06/2020, 45,640309 N – 11,726687 E; rel. 2: Ballin-Miotto (Fontaniva-PD), 9/06/2020, 45,641383 N – 11,725807 E; rel. 3: Ballin-Miotto (Fontaniva-PD), 9/06/2020, 45,642984 N – 11,723962 E; rel. 4: Ballin-Miotto (Fontaniva-PD), 9/06/2020, 45,639141 N – 11,727511 E; rel. 5: S.Lucia di Piave (TV), 9/07/2020, 45,80646 N – 12,26598 E; rel. 6: S. Lucia di Piave (TV), 9/07/2020, 45,8116 N – 12,27109 E; rel. 7: Salettuol (Maserada sul Piave-TV), 13/07/2020, 45,77922 N – 12,3071 E; rel. 8: Grave di Ciano (Crocetta del Montello-TV), 30/05/2020, 45,8392528 N – 12,06488611 E; rel. 9: Grave di Ciano (TV), 30/05/2020, 45,8334167 N – 12,06795278 E; rel. 10: Grave di Ciano (TV), 20/06/2020, 45,841191 N – 12,049816 E; rel. 11: Isola dei Morti (Moriago d. Battaglia-TV), 12/06/2020, 45,84907 N – 12,07862 E; rel. 12: Isola dei Morti (TV), 12/06/2020, 45,848436 N – 12,080083 E; rel. 13: Grave di Ciano (TV), 20/06/2020, 45,839898 N – 12,051321 E; rel. 14: Grave di Ciano (TV), 30/05/2020, 45,8347056 N – 12,07106389 E; rel. 15: Grave di Ciano (TV), 30/05/2020, 45,84363611 N – 12,0581722 E; rel. 16: Falzè di Piave (TV), 12/06/2020, 45,85712 N – 12,17706 E; rel. 17: Grave di Ciano (TV), 20/06/2020, 45,830689 N – 12,05247 E; rel. 18: Grave di Ciano (TV), 20/06/2020, 45,839245 N – 12,049357 E; rel. 19: Isola dei Morti (TV), 01/06/2012, 45,840876 N – 12,103676 E; rel. 20: Isola dei Morti (TV), 24/05/2018, 45,8427778 N – 12,0988889 E; rel. 21: Isola dei Morti (TV), 24/05/2018, 45,845 N – 12,1063889 E; rel. 22: Grave di Ciano (TV), 30/05/2020, 45,84363611 N – 12,0581722 E; rel. 23: Grave di Ciano (TV),

30/05/2020, 45,845688 N – 12,04533611 E; rel. 24: Salettuol (TV), 13/07/2020, 45,77205 N – 12,31414 E; rel. 25: Parabae (Maserada sul Piave-TV), 15/06/2020, 45,76498 N – 12,31571 E; rel. 26: Spresiano (TV), 15/06/2020, 45,78845 N – 12,28371 E; rel. 27: Spresiano (TV), 20/06/1994, 45,787821 N – 12,279095 E; rel. 28: Spresiano (TV), 20/06/1994, 45,787786 N – 12,283453 E; rel. 29: rel. 4 in Table 2 in Feoli Chiapella & Poldini 1994; rel. 30: rel. 5 in Table 2 in Feoli Chiapella & Poldini 1994; rel. 31: rel. 6 in Table 2 in Feoli Chiapella & Poldini 1994.

Appendix 2:

Localities, dates and geographical coordinates of relevés of Table 4.

Rel. 1: Spresiano (TV), 20/06/1994, 45,78882 N – 12,282104 E; rel. 2: Parabae (Maserada sul Piave-TV), 15/06/2020, 45,76314 N – 12,31719 E; rel. 3: Grave di Ciano (Crocetta del Montello-TV), 20/06/2020, 45,842808 N – 12,048247 E; rel. 4: Isola dei Morti (Moriago d. Battaglia-TV), 12/06/2020, 45,842974 N – 12,099714 E; rel. 5: Isola dei Morti (TV), 12/06/2020, 45,848223 N – 12,079289 E; rel. 6: Palazzon (Spresiano-TV), 15/06/2020, 45,78612 N – 12,28332 E; rel. 7: Isola dei Morti (TV), 12/06/2020, 45,842842 N – 12,102824 E; rel. 8: Palazzon (TV), 15/06/2020, 45,78771 N – 12.27907 E.

Appendix 3:

Chorotype grouping in statistical analysis.

cosmopolitan group: cosmopolitan, subcosmopolitan and thermocosmopolitan chorotypes

boreal group: circumboreal, Eurosiberian and Arctic-alpine chorotypes

eurasiatic group: paleotemperate, eurasianic, S-European-S-Siberian, European-W Asiatic, European-Caucasian, European, C-European, S-C-European and NC-European chorotypes

eastern group: NE-eurimediterranean, NE-Mediterranean-montane, SE-European-pontic, SE-European, SE-European-S-Siberian, E-Alpic endemic and SE-Alpic endemic chorotypes

Mediterranean group: stenomediterranean, W-stenomediterranean, SW-stenomediterranean, eurimediterranean, N-eurimediterranean, eurimediterranean-pontic and eurimediterranean-Turanian chorotypes

Atlantic group: W-European and subatlantic chorotypes

orophyte group: N-Mediterranean-montane, E-Mediterranean-montane, W-Mediterranean-montane, orophytic S-European, orophytic S-European-Caucasian,

orophytic SE-European, orophytic European and orophytic C-European chorotypes
endemic group: endemic and Alpic-endemic chorotypes
alien

Appendix 4:

After the acceptance of the article, the occurrence of *Astragalos onobrychidis-Koelerietum pyramidatae* was confirmed along river Tagliamento. The corresponding relévé is here reported. Locality: S. Martino al Tagliamento (PN); date: 12/06/2021; geographical coordinates: 46,02141 N – 12,90508 E; altitude: 70 m a.s.l.; area: 60 m²; cover: 65%; cover Musci: 25%; slope 0°. Species: *Stipa eriocaulis* 3, *Bothriochloa ischaemum* 2b, *Bromus condensatus* 2a, *Fumana procumbens* 2a, *Astragalus onobrychis* 1, *Euphorbia cyparissias* 1, *Helianthemum nummularium/obscurum* 1, *Sanguisorba minor* 1, *Asparagus officinalis* +, *Asperula purpurea* +, *Carex caryophyllea* +, *Carex liparocarpus* +, *Centaurea jacea/gaudinii* +, *Chrysopogon gryllus* +, *Erigeron annuus* +, *Globularia bisnagarica* +, *Hypericum perforatum* +, *Koeleria pyramidata* +, *Ligustrum vulgare* (pl+B2) +, *Linum tenuifolium* +, *Potentilla pusilla* +, *Reseda lutea* +, *Satureja montana/variegata* +, *Ophrys holosericea* r, *Oreoselinum nigrum* r.



Table 1: *Astragalus onobrychidis-Koelerietum pyramidatae* ass. nova. *: sub *K. vallesiana* (Sut.) Gaud.; pl: plantulae; B2: lower shrub layer; na: not available. Geographical area: B river Brenta, P river Piave, N river Natisone.

Tabela 1: *Astragalus onobrychidis-Koelerietum pyramidatae* ass. nova. *: sub *K. vallesiana* (Sut.) Gaud.; pl: plantulae; B2: spodnja grmovna plast; na: ni na voljo. Geografsko območje: B: reka Brenta, P: reka Piava, N: reka Natisone/Nadiža.

relež n°	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	
area (m ²)	30	50	100	70	60	60	100	100	80	60	60	80	80	100	50	60	70	100	100	60	25	30	50									
cover (%)	30	60	60	80	60	75	80	95	85	85	85	80	80	70	90	80	100	95	75	70	95	80	80	90	90	100	100	95				
cover Musci+lichens (%)	60	50	<5	80	15	35	15	15	10	10	25	50	15	20	30	30	nd	10	30	<5	10	25	35	30	nd	nd	nd	nd				
slope (°)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	30	70			
altitude (m a.s.l.)	40	40	40	62	63	44	127	125	132	125	125	132	124	131	91	124	131	116	118	116	131	134	39	37	52	51	na	na	na			
species number	20	21	19	23	28	27	25	28	32	31	37	32	23	31	26	26	35	37	36	40	42	28	33	30	31	26	30	38	37	37		
geographical area	B	B	B	B	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	N	N	N	N				
diff sp of Astragalo-Koelerietum																																
<i>Potentilla pusilla</i> (All)	+	1	+	1	1	+	2a	+	+	1	2a	2a	1	1	+	1	1	1	1	1	+	1	+	1	+	1	+	1	+	30	97	
<i>Astragalus onobrychidis</i> (Cl)	2a	2a	1	1																											74	
<i>Erucastrum nasturtiifolium</i>																															7	23
Centaurieion dichroanthae																																
<i>Carex liparocarpos</i>	2b	2a	2b	3	2a	1	1	1	+	1	2a	2b	1	1	+	+	+	+	+	+	+	+	1	1	1	1	1	1	1	24	77	
<i>Bromus condensatus</i>						2a	2b	2b	3	2a	1	1	2b	1	+								1	2a	2b	3	+	+	+	16	52	
<i>Cytisus purpureus</i>						2a	1	1	+	+	1	+	+	1	1	1	1	+	+	+	+	+	+	+	+	+	+	+	13	42		
<i>Gypsophila repens</i>						+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	11	35			
<i>Cytisus pseudoprocumbens</i>						+	+	+	+	+	+	+	+	1	1	1	1	1	1	1	1	1	1	1	1	1	1	6	19			
Koelerienalia splendens																																
<i>Stipa eriocaulis</i>																															24	77
<i>Teucrium montanum</i>						2a	1	+	2a	+			1	+	+	+																
<i>Trinia glauca</i>																																
<i>Imula ensifolia</i>																																
<i>Globularia cordifolia</i>																																
<i>Satureja monantha variegata</i>																																
<i>Potentilla heptaphylla austriaca</i>																																
Scorzoneralia villosae																																
<i>Plantago holosteum</i>																																
<i>Chrysopogon gryllus</i>																																
<i>Scabiosa triandra</i>																																
<i>Salvia pratensis</i>																																

relevé n°	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	pr.	fl.		
<i>Campanula sibirica</i>																				1	+														
<i>Eryngium amethystinum</i>																																			
<i>Centaurea jacea/gaudinii</i>																																			
<i>Medicago prostrata</i>																																			
<i>Thymus serpylloides</i>																																			
<i>Festuca valesiaca</i>																																			
<i>Hypochoeris maculata</i>																																			
<i>Rhinanthus freynii</i>																																			
<i>Filipendula vulgaris</i>																																			
<i>Lotus herbaceus</i>																																			
<i>Ononis spinosa</i>																																			
<i>Prunella grandiflora</i>																																			
<i>Pseudodysimachion barrelieri/nitens</i>																																			
<i>Ferulago campesiris</i>																																			
Festuco-Brometea																																			
<i>Koeleria pyramidata</i>																																			
<i>Helianthemum nummularium/obscurum</i>																																			
<i>Fumana procumbens</i>																																			
<i>Thymus pulegioides</i>																																			
<i>Globularia bisnagarica</i>																																			
<i>Sanguisorba minor</i>																																			
<i>Centaurea scabiosa</i> s.l.																																			
<i>Thesium humifissum</i>																																			
<i>Artemisia alba</i>																																			
<i>Bothriochloa ischaemum</i>																																			
<i>Euphorbia cyparissias</i>																																			
<i>Leontodon crispus</i>																																			
<i>Gallium verum</i>																																			
<i>Onobrychis arenaria/arenaria</i>																																			
<i>Bromus erectus</i>																																			
<i>Brachypodium rupestre</i>																																			
<i>Carex humilis</i>																																			
<i>Carex caryophyllea</i>																																			
<i>Festuca stricta/sulcata</i>																																			
<i>Hippocratea comosa</i>																																			
<i>Asperula cynanchica</i>																																			

relevé n°	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	pr.	fl.			
<i>Allium carinatum</i>	1	+	1	1																												6	19			
<i>Tenerrimus chamaedrys</i>																																+	6	19		
<i>Gallium lucidum</i>																																	4	13		
<i>Linum tenuifolium</i>																																	4	13		
<i>Lotus corniculatus</i>																																	4	13		
<i>Odonites luteus</i>																																	4	13		
<i>Anacamptis pyramidalis</i>																																+	3	10		
<i>Orobanche gracilis</i>																																	+	3	10	
<i>Trifolium montanum</i>																																	3	10		
<i>Anacamptis coriophora</i>																																	3	10		
<i>Anthyllis vulneraria</i> s.l.																																	2	6		
<i>Campanula glomerata</i>																																	2	6		
<i>Ononis natrix</i>	1	+																																2	6	
<i>Pimpinella saxifraga</i>																																+	2	6		
<i>Polygonia comosa</i>																																+	2	6		
<i>Thymus serpyllum</i> agg.																																+	2	6		
<i>Asperula purpurea</i>																																+	1	3		
<i>Kengia serotina</i>																																1	3			
<i>Medicago falcata</i>																																1	3			
<i>Ophrys holoserica</i>																																1	3			
<i>Orchis militaris</i>																																+	1	3		
<i>Seseli annuum</i>																																+	1	3		
<i>Stachys recta</i>																																+	1	3		
Trifolio-Geranietea																																				
<i>Oreselinum nigrum</i>	r	+																													1	1	20	65		
<i>Anthericum ranosum</i>																															1	1	1	7	23	
<i>Buphthalmum salicifolium</i>																															1	3				
<i>Clematis recta</i>																															+	1	3			
<i>Lilium bulbiferum</i>																															+	1	3			
<i>Trifolium rubens</i>																															+	1	3			
Koelerio-Corynephoretea+Sedo-Sclerantheeta																																				
<i>Petrorhagia saxifraga</i>																															+	1	+	1	15	48
<i>Sedum sexangulare</i>																															+	+	+	+	9	29
<i>Medicago minima</i>																															+	+	6	19		
<i>Dianthus sylvestris</i>																															+	+	3	10		



relevé n°	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	pr.	fl.
<i>Allium lusitanicum</i>																														1	3		
<i>Catapodium rigidum</i>	+																													1	3		
<i>Ceratistium semidecandrum</i>																														1	3		
<i>Trifolium campestre</i>		r																												1	3		
Other species																																	
<i>Erigeron annuus</i>	+	+	+	1	1	3	+	r	r	+	+	r	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	18	58				
<i>Asperagus officinalis</i>							+	r	+	+	+	r	+	+	r	+	r	+	r	+	r	+	r	+	r	+	12	39					
<i>Reseda lutea</i>		+					+	+	+	+	+	+	+	+	+	+	+	1	+	+	+	+	+	+	+	+	11	35					
<i>Echium vulgare</i>	+	+	r	+														r	r	r	r								8	26			
<i>Populus nigra</i> (pl+B2)	+					+	1	+					r	+				+											7	23			
<i>Amorpha fruticosa</i> (B2)						+	+	+										1	+	+	+								5	16			
<i>Ligustrum vulgare</i> (pl+B2)										+																			5	16			
<i>Arenaria campesiris</i>																													4	13			
<i>Cornus sanguinea</i> (pl+B2)																		+	r										4	13			
<i>Diplotaxis tenuifolia</i>	+		r									r																	4	13			
<i>Hypericum perforatum</i>													+					+	r										4	13			
<i>Rosa</i> sp. pl. (pl+B2)																													4	13			
<i>Sporobolus neglectus</i>																													4	13			
<i>Centaurium erythraea</i>																													3	10			
<i>Centaura stoebe</i>												r																	3	10			
<i>Plantago lanceolata</i>																													3	10			
<i>Rhamnus saxatilis</i> (B2)																		1											3	10			
<i>Salix eleagnos</i> (pl+B2)																													3	10			
<i>Arenaria serpyllifolia</i>																													3	10			
<i>Bromus squarrosus</i>																													2	6			
<i>Cephalanthera damasonium</i>												r																	2	6			
<i>Chondrilla juncea</i>																													2	6			
<i>Cytisus hirsutus</i>																													2	6			
<i>Erica carnea</i>																													2	6			
<i>Euphorbia maculata</i>																													2	6			
<i>Gymnadenia conopsea</i>																													2	6			
<i>Hieracium murorum</i> agg.																													2	6			
<i>Pilosella piloselloides</i>																													2	6			
<i>Lemnropis nigritans</i> (B2)																													2	6			
<i>Prunus spinosa</i> (pl+B2)																													2	6			
<i>Silene vulgaris</i>																													2	6			

relevé n°	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	pr.	fr.
<i>Thymelaea passerina</i>	r	+																													2	6	
<i>Biscutella laevigata</i>																														+	1	3	
<i>Calamintha nepeta</i>	+																													1	3		
<i>Berberis vulgaris</i> (B2)																														1	3		
<i>Carex sylvatica</i>	r																													1	3		
<i>Clematis vitalba</i>																														1	3		
<i>Crepis foersteri</i>	r																													1	3		
<i>Cynodon dactylon</i>																														1	3		
<i>Dactylis glomerata</i>	+																													1	3		
<i>Daucus carota</i>																														1	3		
<i>Epiptoleium dodoneae</i>																														1	3		
<i>Frangula alnus</i>																														1	3		
<i>Fraxinus ornus</i> (pl+B2)																														1	3		
<i>Molinia arundinacea</i>																														1	3		
<i>Poa compressa</i>																														1	3		
<i>Salvia verticillata</i>																														+	1	3	
<i>Teucrium botrys</i>																														1	3		
<i>Tordyli斯japonica</i>	+																													1	3		

Table 2: Percentage distribution of chorological groups in different comparison coenosis (percentage cover values). A-K: *Astragalos-Koelerietum* (this study); C-C-v: *Chamaecytiso-Chrysopogonetum* from Veneto (this study); C-G: *Centaureo-Globularietum* (Feoli Chiapella & Poldini 1994); S-C: *Schoeno-Chrysopogonetum* (Feoli Chiapella & Poldini 1994); C-C-f: *Chamaecytiso-Chrysopogonetum* from Friuli-Venezia Giulia (Feoli Chiapella & Poldini 1994); A-A: *Astragalo-Artemisiatum* (Biondi et al. 1997).

Tabela 2: Odstotni delež horoloških skupin v primerjanih združbah (pokrovnost v odstotkih) A-K: *Astragalos-Koelerietum* (ta raziskava); C-C-v: *Chamaecytiso-Chrysopogonetum* iz Benetceje (ta raziskava); C-G: *Centaureo-Globularietum* (Feoli Chiapella & Poldini 1994); S-C: *Schoeno-Chrysopogonetum* (Feoli Chiapella & Poldini 1994); C-C-f: *Chamaecytiso-Chrysopogonetum* iz Furlanije-Julijske krajine (Feoli Chiapella & Poldini 1994); A-A: *Astragalo-Artemisiatum* (Biondi et al. 1997).

	A-K	C-G	S-C	C-C_v	C-C_f	A-A
cosmopolitan	8,37	6,04	5,31	0,59	0,03	22,06
boreal	2,19	0,83	1,91	0,99	8,35	0,40
eurasitic	24,35	21,02	42,94	75,16	65,91	25,05
eastern	13,51	10,65	24,94	10,01	17,51	0,32
mediterranean	9,38	4,65	4,38	8,97	4,35	39,74
atlantic	25,44	5,44	14,48	1,64	0,93	0,24
orophytes	14,80	35,64	4,46	2,36	2,78	6,89
endemic	0,11	15,73	1,58	0,00	0,17	4,90
alien	1,84	0,00	0,00	0,27	0,00	0,40

Table 3: Synoptic table of dry grassland communities on fluvial terraces of gravel bed deposits from North-eastern Po plain. Abbreviations as in Table 2. Companion species occurring at least once in ≥ II frequency class were included.

Tabela 3: Sinoptična tabela združb suhih travnišč na rečnih prodnatih terasah v severovzhodni Padski nižini. Okrajšave so enake kot v Tabeli 2. Prikazane so spremeljevalne vrste, ki se pojavljajo vsaj enkrat s frekvenčnim razredom večjim od II.

relevés (n°)	A-K	C-G	C-C_v	C-C_f	S-C	A-A
	31	14	8	7	10	17
diff sp of Astragalo-Koelerietum						
<i>Astragalus onobrychis</i> (Cl)	74		38			59
<i>Potentilla pusilla</i> (All)	97		63			6
<i>Erucasterum nasturtiifolium</i>	23					
diff sp of Centaureo-Globularietum						
<i>Euphorbia triflora/kernerii</i> (All)		93			30	
<i>Matthiola fruticulosa/valesiaca</i>		57				
<i>Brassica glabrescens</i>		36				
<i>Crambe tataria</i>		36				
diff sp of Chamaecytiso-Chrysopogonetum						
<i>Dianthus sanguineus</i>				71		
<i>Prunella laciniata</i> (All2)				43		
<i>Serapias vomeracea</i>				43		
<i>Anacamptis coriophora</i>	6		63	14		
<i>Cytisus pseudoprocumbens</i> (All)	19	50	88	14	60	
<i>Medicago prostrata</i> (O)	16	7	75		10	
diff sp of Schoeno-Chrysopogonetum						
<i>Globularia bisnagarica</i> (Cl)	77	14	50	14	90	65
diff sp of Astragalo-Artemisietum						
<i>Artemisia alba</i> (Cl)	65	29	63			100
<i>Festuca robustifolia</i>						76
<i>Centaurea aplolepa/lunensis</i>						76
Centaureion dichroanthae						
<i>Bromopsis condensata</i>	52	71	50	57	100	
<i>Cytisus purpureus</i>	42	57	50		80	
<i>Centaurea dichroantha</i>		100			70	
<i>Carex liparocarpos</i>	77	14	50		20	6
<i>Gypsophila repens</i>	35	79	13		10	
<i>Carex mucronata</i>		93			10	
<i>Sesleria caerulea</i>		86				
<i>Polygala forojulensis</i>		43			20	
<i>Lomelosia graminifolia</i>		36			10	
<i>Hieracium porrifolium</i>		36				
<i>Euphrasia cuspidata</i>		14				
<i>Allium ochroleucum</i>		7				
<i>Gentiana clusii</i>		7				
<i>Leontodon incanus</i> s.l.		7				
Hypochoeridenion maculatae						
<i>Centaurea scabiosaffritschii</i>				86	50	
<i>Knautia ressmannii</i>				29	50	

relevés (n°)	A-K	C-G	C-C_v	C-C_f	S-C	A-A
<i>Rhinanthus freynii</i>	10			29	20	
<i>Carex montana</i>				43		
<i>Prunella grandiflora</i>	3			14	10	
<i>Cirsium pannonicum</i>					20	
<i>Scorzoneronion villosae</i> (All 2)						
<i>Hypochaeris maculata</i>	10	14		86	70	
<i>Filipendula vulgaris</i>	6			86	50	
<i>Plantago media</i>			25	14	40	
<i>Ononis spinosa</i>	3			43	10	18
<i>Lotus herbaceus</i>	3		13		20	
<i>Knautia illyrica</i>				14	10	
<i>Scorzonera villosa</i>				14		
<i>Euphorbia brittingeri</i>					10	
<i>Ferulago campestris</i>	3					
<i>Scorzoneretalia villosae</i>						
<i>Chrysopogon gryllus</i>	45	64	100	100	100	
<i>Plantago holosteum</i>	55	86	50	14	70	
<i>Salvia pratensis</i>	32		88	43	50	6
<i>Scabiosa triandra</i>	35	14	25	57	70	
<i>Centaurea jacea/gaudinii</i>	16	7	50	57	50	
<i>Betonica officinalis/serotina</i>			7	57	60	
<i>Eryngium amethystinum</i>	16	21	50			
<i>Campanula sibirica</i>	19	21	25		10	
<i>Plantago argentea/liburnica</i>			21		30	
<i>Noccaea praecox</i>				14		
<i>Centaurea triumfettii</i>					10	
<i>Festuca valesiaca</i>	10					
<i>Pseudolysimachion barrelieri/nitens</i>	3					
<i>Koelerienalia splendens</i>						
<i>Teucrium montanum</i>	52	93	50		10	24
<i>Stipa eriocalis/eriocalis</i>	77	93	50			
<i>Trinia glauca</i>	13	71	50		40	
<i>Globularia cordifolia</i>	10	100			30	
<i>Inula ensifolia</i>	13	86	13			
<i>Potentilla heptaphylla/australis</i>	3	43			50	
<i>Scorzonera austriaca</i>		57			20	
<i>Genista sericea</i>		64			10	
<i>Seseli kochii</i>		43			10	
<i>Satureja montana/variegata</i>	10	14				
<i>Festuco-Brometea</i>						
<i>Thymus serpyllum</i> agg.	90	71	100	86	90	88
<i>Koeleria pyramidata</i>	100	100	100	71	80	
<i>Helianthemum nummularium/obscurum</i>	94	64	100	71	70	29
<i>Sanguisorba minor</i>	77	21	63	29	90	82
<i>Galium verum</i>	48	21	88	100	100	
<i>Euphorbia cyparissias</i>	65	21	88	57	20	47

relevés (n°)	A-K	C-G	C-C_v	C-C_f	S-C	A-A
	31	14	8	7	10	17
<i>Carex humilis</i>	39	86	63	29	70	
<i>Fumana procumbens</i>	90	93	38			59
<i>Lotus corniculatus</i>	13	29	38	100	100	
<i>Thesium humifusum</i>	68	71	88		40	6
<i>Brachypodium rupestre</i>	39		50	71	100	12
<i>Festuca stricta/sulcata</i>	29	14	63	57	80	
<i>Leontodon crispus</i>	61	43	25	43	60	
<i>Asperula cynanchica</i>	26	43	13	57	80	
<i>Galium lucidum</i>	13	93	38	14	50	6
<i>Bothriochloa ischaemum</i>	65	7	25		10	100
<i>Hippocrepis comosa</i>	29	36	38	14	80	
<i>Teucrium chamaedrys</i>	19	29	63	14	60	12
<i>Bromopsis erecta</i>	42		88			65
<i>Carex caryophyllea</i>	29	14	25	43	70	
<i>Trifolium montanum</i>	10		50	71	50	
<i>Briza media</i>			13	86	60	
<i>Anthyllis vulneraria</i> s.l.	6	36		57	40	
<i>Pimpinella saxifraga</i>	6		13	71	40	
<i>Centaurea scabiosa</i>	71		50			
<i>Pilosella officinarum</i>			14		57	6
<i>Anacamptis morio</i>			43		14	50
<i>Linum catharticum</i>			7		29	70
<i>Anacamptis pyramidalis</i>	10	14	25	14	40	
<i>Asperula purpurea</i>	3	14			14	71
<i>Linum tenuifolium</i>	13	57	13			6
<i>Onobrychis arenaria/arenaria</i>	48		38			
<i>Campanula glomerata</i>	6				57	10
<i>Carex flacca</i>					14	40
<i>Odontites luteus</i>	13					41
<i>Plantago sempervirens</i>						53
<i>Ranunculus bulbosus</i>					29	20
<i>Stachys recta</i> s.l.	3					41
<i>Polygala comosa</i>	6			38		
<i>Ononis natrix</i>	6					35
<i>Eryngium campestre</i>						41
<i>Allium coloratum</i>					29	12
<i>Melica ciliata</i>			7			29
<i>Medicago lupulina</i>					14	20
<i>Orchis militaris</i>	3			25		
<i>Neotinea tridentata</i>						20
<i>Allium carinatum</i>	19					
<i>Ophrys apifera</i>			7			10
<i>Medicago falcata</i>	3			13		
<i>Kengia serotina</i>	3					12
<i>Asperula aristata</i>						12
<i>Carlina vulgaris</i>						12
<i>Orobanche gracilis</i>	10					
<i>Carlina acaulis</i>			7			
<i>Alyssum montanum</i>			7			

relevés (n°)	A-K	C-G	C-C_v	C-C_f	S-C	A-A
<i>Gentianella pilosa</i>		7				
<i>Ophrys holoserica</i>	3					
<i>Seseli annuum</i>	3					
<i>Molinio-Arrhenatheretea</i>						
<i>Dactylis glomerata</i>	3	29	63	43	90	59
<i>Gymnadenia conopsea</i>	6	29	13	29	20	
<i>Tragopogon pratensis</i> aggr.				43	30	12
<i>Leontodon hispidus</i>				14	70	
<i>Jacobaea vulgaris</i>		14		43	10	
<i>Prunella vulgaris</i>		14			50	
<i>Ranunculus tuberosus</i>				43		
<i>Rumex acetosa</i>				43		
<i>Linum bienne</i>						41
<i>Trifolium pratense</i>				14	20	
<i>Arrhenatherum elatius</i>				29		
<i>Holcus lanatus</i>				29		
<i>Trifolio-Geranietea</i>						
<i>Oreoselinum nigrum</i>	65	93	88	100	80	
<i>Anthericum ramosum</i>	23	43	38	43		
<i>Buphtalmum salicifolium</i>	3	7	25	57	40	
<i>Trifolium rubens</i>	3			71		
<i>Vincetoxicum hirundinaria</i>		7	13	14	30	
<i>Cruciata glabra</i>		7		29		
<i>Viola hirta</i>				29		
<i>Koelerio-Corynephoretea+Sedo-Scleranthesetea</i>						
<i>Sedum sexangulare</i>	29		25		10	35
<i>Petrorhagia saxifraga</i>	48		25		10	
<i>Catapodium rigidum</i>	3					24
<i>Sedum acre</i>						24
<i>Sedum sediforme</i>						24
<i>Iblaspietea rotundifolii+Asplenietea trichomanis</i>						
<i>Pilosella piloselloides</i>	6	50		71	60	
<i>Achnatherum calamagrostis</i>		14				24
<i>Epilobium dodonaei</i>	3					24
<i>Elyno-Seslerietea</i>						
<i>Biscutella laevigata</i>	3	50		86	40	
<i>Leucanthemum heterophyllum</i>					60	
<i>Rhinanthus glacialis</i>		14			40	
<i>Dianthus sternbergii</i>		36				
<i>Helianthemum alpestre</i>		36				
<i>Dryas octopetala</i>		21				
Other species						
<i>Plantago lanceolata</i>	10		25	57	30	53
<i>Erica carnea</i>	6	93			50	
<i>Schoenus nigricans</i>		79			60	

relevés (n°)	A-K	C-G	C-C_v	C-C_f	S-C	A-A
<i>Leucanthemum vulgare</i> agg.	31	14	8	7	10	17
<i>Hypericum perforatum</i>	13		13			88
<i>Cytisus hirsutus</i>	6			86	10	
<i>Asparagus officinalis</i>	39		38			18
<i>Anthoxanthum odoratum</i>				86		
<i>Erigeron annuus</i>	58		25			
<i>Reseda lutea</i>	35	21	25			
<i>Polygala vulgaris</i>				71	10	
<i>Genista tinctoria</i>				57	20	
<i>Blackstonia perfoliata</i>		7			50	18
<i>Potentilla erecta</i>				43	30	
<i>Diplotaxis tenuifolia</i>	13	21	13			18
<i>Agrostis capillaris</i>				57		
<i>Centaurea erythraea</i>	10	7	38			
<i>Amorpha fruticosa</i> (B2)	16		38			
<i>Chondrilla juncea</i>	6					47
<i>Dittrichia viscosa</i>						53
<i>Silene vulgaris</i>	6				20	18
<i>Potentilla alba</i>				43		
<i>Cota tinctoria</i>						41
<i>Avena barbata</i>						41
<i>Setaria viridis</i>						41
<i>Ligustrum vulgare</i> (pl+B2)	16		25			
<i>Echium vulgare</i>	26					12
<i>Erigeron canadensis</i>						35
<i>Erigeron sumatrensis</i>						35
<i>Sixalix atropurpurea</i>						35
<i>Rhamnus saxatilis</i> (B2)	10		25			
<i>Crepis froelichiana/dinarica</i>					30	
<i>Thalictrum minus</i> s.l.					30	
<i>Potentilla pedata</i>						29
<i>Scabiosa columbaria</i>						29
<i>Populus nigra</i> (pl+B2)	23					6
<i>Serratula tinctoria</i> s.l.		7			20	
<i>Helichrysum italicum</i>						24
<i>Inula spiraeifolia</i>						24
<i>Epipactis atrorubens</i>				20		
<i>Genista germanica</i>					20	
<i>Scirpoides holoschoenus</i>					20	

Table 4: *Chamaecytiso hirsuti-Chrysopogonetum grylli* Venetian provenance. Abbreviations as in Table 1.

Tabela 4: *Chamaecytiso hirsuti-Chrysopogonetum grylli* iz Benečije. Okrajšave so enake kot v Tabeli 1.

relevé n°	1	2	3	4	5	6	7	8		
area (m ²)	60	80	60	100	80	60	100	60		
cover (%)	100	100	95	95	100	95	100	90		
cover Musci+lichens (%)	na	5	-	5	-	15	-	10		
altitude (m a.s.l.)	52	36	132	118	125	51	117	51		
species number	32	26	36	36	30	30	47	30	pr	fr%
geographical area	P	P	P	P	P	P	P	P		
diff sp of Chamaecytiso-Chrysopogonetum										
<i>Cytisus pseudoprocumbens</i> (All)	1	2a		2b	+	+	1	+	7	88
<i>Medicago prostrata</i> (O)	+	+			+	1	+	2a	6	75
<i>Anacamptis coriophora</i>			+	+	1	+	+		5	63
Centaureion dichroanthae										
<i>Potentilla pusilla</i> agg.		+		1		+	+	1	5	63
<i>Bromus condensatus</i>	1				+	1		2a	4	50
<i>Cytisus purpureus</i>			1		+	+	2a		4	50
<i>Carex liparocarpos</i>	+	+				+		1	4	50
<i>Gypsophila repens</i>			+						1	13
Scorzonerion villosae										
<i>Plantago media</i>				+			+		2	25
<i>Lotus herbaceus</i>							+		1	13
Scorzoneretalia villosae										
<i>Chrysopogon gryllus</i>	5	3	4	3	4	3	4	2b	8	100
<i>Salvia pratensis</i>	r	+	+		+	+	+	+	7	88
<i>Plantago holosteum</i>	1	1		+				2a	4	50
<i>Centaurea jacea/gaudinii</i>			+	+	+		+		4	50
<i>Eryngium amethystinum</i>	+	+				+		+	4	50
<i>Scabiosa triandra</i>			+		+				2	25
<i>Campanula sibirica</i>				+			1		2	25
Koelerienalia splendentis										
<i>Teucrium montanum</i>	+	+				+		+	4	50
<i>Stipa eriocaulis/erioaulis</i>	+				r		+	+	4	50
<i>Trinia glauca</i>	1	+				+		1	4	50
<i>Inula ensifolia</i>				+					1	13
Festuco-Brometea										
<i>Thymus pulegioides</i>	1	1	+	1	1	1	1	2b	8	100
<i>Koeleria pyramidata</i>	1	1	1	1	1	+	1	+	8	100
<i>Helianthemum nummularium/obscurum</i>	1	1	1	1	+	1	1	1	8	100
<i>Bromus erectus</i>	+	3	2b	3	2a	2a	2a		7	88
<i>Euphorbia cyparissias</i>	+	+	+	+	1	2a	+		7	88
<i>Galium verum</i>	1	+		1	+	1	+	2b	7	88
<i>Thesium humifusum</i>	+	+		+	+	+	+	+	7	88
<i>Artemisia alba</i>	1		+	2a	+		1		5	63

relevé n°	1	2	3	4	5	6	7	8	pr.	fr.
<i>Carex humilis</i>			1	+		2b	1	1	5	63
<i>Festuca stricta/sulcata</i>	1	2b				2a	+	1	5	63
<i>Sanguisorba minor</i>	+	+				+	+	+	5	63
<i>Teucrium chamaedrys</i>	1	1	+			2a		2b	5	63
<i>Brachypodium rupestre</i>	+		+		1		2a		4	50
<i>Centaurea scabiosa</i> s.l.			+	+		+	+		4	50
<i>Globularia bisnagarica</i>	+			1	+	1			4	50
<i>Trifolium montanum</i>			+	1	+		1		4	50
<i>Fumana procumbens</i>	+		+		+				3	38
<i>Galium lucidum</i>	1	+						1	3	38
<i>Hippocratea comosa</i>			+		+		+		3	38
<i>Lotus corniculatus</i>			+	+	+				3	38
<i>Onobrychis arenaria</i>	+					+	+		3	38
<i>Polygala comosa</i>			+	+			+		3	38
<i>Astragalus onobrychis</i>			+			+		+	3	38
<i>Anacamptis pyramidalis</i>			+				+		2	25
<i>Bothriochloa ischaemum</i>						1		1	2	25
<i>Carex caryophyllea</i>				+			+		2	25
<i>Leontodon crispus</i>	+			+					2	25
<i>Orchis militaris</i>		r		+					2	25
<i>Asperula cynanchica</i>				+					1	13
<i>Briza media</i>							2a		1	13
<i>Gymnadenia conopsea</i>							+		1	13
<i>Linum tenuifolium</i>					+				1	13
<i>Medicago falcata</i>	+								1	13
<i>Pimpinella saxifraga</i>							r		1	13
Trifolio-Geranietea										
<i>Peucedanum oreoselinum</i>	+	+	+	+	+	+	+		7	88
<i>Anthericum ramosum</i>	+					r	+		3	38
<i>Buphtalmum salicifolium</i>			+				+		2	25
<i>Clematis recta</i>			+						1	13
<i>Vincetoxicum hirundinaria</i>			+						1	13
Other species										
<i>Dactylis glomerata</i>	+	+	1	1			1		5	63
<i>Amorpha fruticosa</i> (B2)		+		+			+		3	38
<i>Asparagus officinalis</i>		+	+				+		3	38
<i>Centaurium erythraea</i>	r		+			+			3	38
<i>Erigeron annuus</i>				+			+		2	25
<i>Ligustrum vulgare</i> (pl+B2)		+					+		2	25
<i>Petrorhagia saxifraga</i>	+							+	2	25
<i>Plantago lanceolata</i>				+	+				2	25
<i>Reseda lutea</i>	+							+	2	25
<i>Rhamnus saxatilis</i> (B2)					+		+		2	25
<i>Sedum sexangulare</i>	+			+					2	25
<i>Achillea millefolium</i> agg.					+				1	13
<i>Chenopodium album</i>								+	1	13
<i>Diplotaxis tenuifolia</i>								+	1	13

relevé n°	1	2	3	4	5	6	7	8	pr.	fr.
<i>Fraxinus ornus</i> (pl+B2)							+		1	13
<i>Hypericum perforatum</i>		r							1	13
<i>Inula salicina</i>							+		1	13
<i>Medicago minima</i>							+		1	13
<i>Pyrus communis</i> (B2)		r							1	13
<i>Salix eleagnos</i> (pl+B2)			1						1	13
<i>Solanum nigrum</i>							r		1	13
<i>Ulmus minor</i> (pl+B2)							r		1	13
<i>Vicia cracca</i> s.l.			+						1	13