

Phase angle as a prognostic indicator of surgical outcomes in patients with gastrointestinal cancer

Jana Gulin¹, Ester Ipavic¹, Denis Mlakar Mastnak², Erik Breclj², Ibrahim Edhemovic², Nada Rotovnik Kozjek^{2,3}

¹ Medical Centre Ljubljana, Ljubljana, Slovenia

² Institute of Oncology Ljubljana, Ljubljana, Slovenia

³ University of Ljubljana, Faculty of Medicine, Ljubljana, Slovenia

Radiol Oncol 2023; 57(4): 524-529.

Received 3 August 2023

Accepted 30 September 2023

Correspondence to: Asst. Prof. Nada Rotovnik Kozjek, M.D., Ph.D., Institute of Oncology Ljubljana, Zaloška 2, SI-1000 Ljubljana, Slovenia. E-mail: nkozjek1@gmail.com.

Disclosure: No potential conflicts of interest were disclosed.

This is an open access article distributed under the terms of the CC-BY license (<https://creativecommons.org/licenses/by/4.0/>).

Background. In patients with gastrointestinal cancer with planned elective surgery, malnutrition increases the risk of adverse outcomes in the postoperative period. The phase angle, measured by the bioelectrical impedance analysis is an indicator of the metabolic and functional status of the patient. It may be an important prognostic indicator for the clinical outcome of post-surgical treatment in patients with gastrointestinal cancer.

Patients and methods. In this prospective study, 70 patients with gastrointestinal cancer had their phase angles measured by the bioelectrical impedance analysis before the surgery. During the first month after the surgery, we documented the postoperative complications from the patient's records and classified them according to the Clavien Dindo classification of surgical complications. The time of hospitalization was also recorded. The data was statistically analysed in SPSS.

Results. We found a statistically significant difference ($p = 0.036$) in the average value of phase angles between the group of patients who had postoperative complications (phase angle 5.09°) and the group without postoperative complications (5.64°). We noted a correlating trend of decreasing phase angle values and increasing hospitalization time ($P_e R = -0.40$, $p = 0.001$). The phase angle cut-off value (5.5°) was calculated using the ROC curve method, predicting a higher risk of the postoperative complications ($p = 0.037$) in patients with lower phase angle.

Conclusions. Lower phase angle values before surgery were associated with more complications during the first month after surgery and longer hospitalization time. We found that a phase angle below than 5.5° could serve as a marker that predicts a greater risk of postoperative complications.

Key words: phase angle; colorectal cancer; postoperative complications; malnutrition

Introduction

Bioelectric Impedance Analysis (BIA) is a technique used to assess body composition and is becoming increasingly established as a tool to assess nutritional status in patients due to its ease of use, low cost, and non-invasiveness.^{1,2} BIA does not di-

rectly measure the body composition, but instead measures the drop in voltage of an alternating electric current, as it travels across the body. The phase angle is the quotient of measured resistance and reactance.¹⁻⁴ It is interpreted as an indicator of membrane integrity and water distribution between the intracellular and extracellular spaces.

Phase Angle is used to evaluate body cell mass, which serves as a tool to assess nutritional status in adults and children. Lower phase angle suggests cell degradation with a concomitant decrease in overall cell number with reduced integrity and functional capacity of cell membranes. On the other hand, higher phase angle points to the presence of more functional, intact cell membranes.⁵⁻⁹

Malnutrition is a common manifestation of advanced stage cancer and can severely affect the patient morbidity and mortality.¹⁰ On a cellular level malnutrition is reflected by the diminished integrity of the cell membranes and by altered water distribution throughout the body.¹¹ Body composition analysis is therefore an essential tool in assessing nutritional status in cancer patients.¹² The clinical role of measuring the patient's phase angle is becoming progressively more important. BIA-derived phase angle can help establish guidelines for preventive and supportive measures in patients with malnutrition, as it allows for early identification of high-risk patients with inadequate physiological reserves. The method has previously been shown to have prognostic value in patients with liver cirrhosis, haemodialysis, cancer, HIV/AIDS infection, and lung disease.^{5,13-18} For example, patients with stage III or IV colorectal cancer who had a phase angle less than or equal to 5.57° had a median survival of 8.6 months, while those who had a phase angle greater than 5.57° had a median survival of 40.4 months.¹⁴

The primary objective of this study was to determine if the phase angle can be useful as an independent predictor of post-surgical complications in gastrointestinal cancer patients.

Patients and methods

Setting and patients

Our study was a prospective observational study that was performed at the Institute of Oncology Ljubljana, Slovenia, between November 2018 and February 2019. During the study period, BIA was performed on every gastrointestinal cancer patient over 18 years of age that was admitted for elective surgery at the Institute of Oncology Ljubljana and agreed to participate in the study. Exclusion criteria were pregnancy or an implanted pacemaker. The study protocol was approved by the Ministry of Health Medical Ethics Committee and the Institute of Oncology (No. 0120-518/2018/7). Every included patient was fully informed of the study design and signed a written informed consent form.

Bioelectrical Impedance Analysis (BIA) and other measurements

BIA was performed with a Bodystat (R) Quadscan 4000 machine (Douglas, UK). This phase-sensitive BIA device uses an alternating current of 0.8 mA with frequencies of 5, 50, 100, and 2000 kHz to measure the body impedance. The BIA-derived phase angle was calculated from the impedance at 100 kHz.

Every patient had their phase angle measured the day before surgery and then a week and a month after the surgery. The patients were instructed to abstain from eating and drinking for at least 4 hours prior to the measurement and to abstain from any physical exercise for 8 hours prior to the measurement as well. Two pairs of electrodes were placed on the dorsal side of their right hand and right foot, and they were instructed to lie still in a supine position with no parts of the body touching one another. Clinical data, including gender, age, the exact location of malignant disease, complications in the postoperative course, and hospitalization time were obtained from the hospital information system.

Clavien Dindo Classification of surgical complications

The patients were categorized in different grades of the Clavien Dindo classification of surgical complications (revised 2004).¹⁹ The classification uses the degree of most severe pharmacological or surgical treatment needed after surgery, to distinguish between the grades of post-surgical complications. A normal postoperative course without any complications (use of analgesics is considered as normal postoperative course) is classified as degree 0. Degree 1 allows the use of antiemetics, antipyretics, potent analgesics, diuretics, physiotherapy, and treatment of wound infections. Degree 2 additionally includes the use of other drugs, blood transfusion, and total parenteral nutrition. Degree 3 allows surgical, endoscopic, or radiological interventions. Degree 4 includes life-threatening complications that require treatment in an intensive care setting. Fatal complications are classified as a degree 5.

Statistics

Numeric variables are expressed in terms of their mean values and a 95% confidence interval. Discrete variables are expressed as percentages.

TABLE 1. Characteristics of patients

Characteristics	All patients	Patients with no complications (G0)	Patients with complications (G1–G5)	p-value
Gender				
Male	50 (71 %)			
Female	20 (29 %)			
Age (years)^a	65.0 (10.7) [62.4–67.6]	61.7 (9.8) [56.9–66.6]	66.1 (10.9) [63.1–69.2]	0.137
Phase Angle (°)^a	5.23 (2.77) [5.0–5.5]	5.64 (0.72) [5.3–6.0]	5.09 (0.98) [4.8–5.4]	0.036*
Location of the primary tumor (%)				
Colon	17 (24.3)	4 (22.2)	13 (25.0)	
Rectosigmoid	2 (2.9)	1 (5.5)	1 (1.9)	
Rectum	45 (64.3)	11 (61.1)	34 (65.4)	
Anus	1 (1.4)	1 (5.5)	0 (0)	
Breast	1 (1.4)	0 (0)	1 (1.9)	
Ovary	4 (5.7)	1 (5.5)	3 (5.8)	

^a Arithmetic mean (standard deviation) [confidence interval].

* Statistically significant p-value

TABLE 2. Mean age and mean phase angle in each Clavien Dindo grade

	Number of patients (%)	Age (years) ^a	Phase angle (°) ^a
Grade 0	18 (26)	61.7 [56.9–66.6]	5.64 [5.28–6.00]
Grade I	5 (7)	56.0 [50.2–61.8]	6.22 [5.64–6.80]
Grade II	38 (54)	67.1 [63.4–70.7]	4.99 [4.67–5.31]
Grade III	4 (6)	62.2 [45.0–79.5]	5.15 [3.41–6.88]
Grade IV	5 (7)	72.2 [60.5–83.9]	4.64 [3.77–5.51]
Grade V	0 (0)		

^a Arithmetic mean (standard deviation) [95% confidence interval].

The Shapiro-Wilk test was used to check whether the data is normally distributed. The limit for statistical significance was set at $p < 0.05$. Logistic regression was used to test whether the initial pre-operative phase angle, as an independent variable, impacts the odds of post-surgical complications. The results are expressed as quotients. A cut-off value for the phase angle as a predictive factor for post-surgical complications was estimated with a ROC curve, and its sensitivity and specificity were calculated. The statistical analysis was done with the IBM SPSS 23.0 statistical program.

Results

During the study period (between November 2018 and February 2019), BIA analysis was performed

on every gastrointestinal cancer patient planned for elective surgery at the Institute of Oncology Ljubljana that agreed to participate in the study. Seventy patients were recruited. Characteristics of the included patients (column 1), of patients without any complications (column 2) and patients with complications (column 3) are presented in Table 1. The mean age of all patients was 65.0 ± 10.7 years, and 71% were male. Most patients were admitted because of rectal cancer (64.3%). Mean phase angle of all patients was $5.23^\circ \pm 2.77^\circ$. In total, 52 (74.3%) patients had a complication (Clavien Dindo grade 1-5) after surgery (Table 1 and Table 2).

Phase angle and the likelihood of complications

The phase angle of patients without a complication was significantly higher than that of the patients with a complication ($5.64^\circ \pm 0.72^\circ$ vs. $5.09^\circ \pm 0.98^\circ$, $p = 0.036$). Univariate logistic regression analysis showed that the phase angle was associated with the likelihood of a complication (phase angle: odds ratio = 0.48). The odds of a complication in a patient with a phase angle of 5.0 was 3.83, whereas it was only 1.84 in a patient with a phase angle of 6.0. The probabilities of the occurrence of a complication at different phase angles were calculated and are shown in Table 3. The area under the curve (AUC) of the ROC curve for phase angle for the likeli-




FIGURE 1. The ROC curve for the phase angle.

hood of a complication is 0.666 (CI: 0.529 - 0.803, $p = 0.037$), see Figure 1. The determination coefficient (Nagelkerke R square) is 0.104, which means that our model explains 10.4% of the complication likelihood variance. The Hosmer and Lemeshow test value for the model is 0.766.

The optimal cut-off value for the phase angle

The cut-off value of phase angle that was derived from the ROC-curve was 5.5° (Figure 1). It yielded a sensitivity of 63% and a specificity of 61% (Table 4). The likelihood of a complication was higher in patients with phase angle below 5.5° than in the patients with phase angle above that value (82.5% vs. 63.3%, Figure 1).

Discussion

The patients with a lower initial phase angle were shown to have a statistically significant higher likelihood of post-surgical complications. The mean phase angle in the group with complications (5.64°) was statistically higher than the mean phase angle in the group without complications (5.09°).

We used univariate logistic regression to calculate the likelihood of a complication at different initial phase angle values. At a phase angle of 5.0° , the odds of a complication are 3.83, and the likelihood of a complication is 79.3%. At a phase angle of

6.0° , the odds of a complication are reduced to 1.84, and the likelihood of a complication is reduced to 64.8%.

We are not aware of any prospective studies investigating the role of phase angle as a prognostic indicator of post-surgical complications in colorectal cancer patients. The retrospective data analysis from 210 elderly patients (aged ≥ 65 years) undergoing gastrectomy showed that preoperative low phase angle predicts a higher risk of postoperative complications.²⁰ Similar observations were made in a study among patients admitted to the ICU, where phase angle at admission was shown to be a predictor of 90-day mortality. The mean phase angle of survivors was significantly higher than that of the non-survivor group ($5.0^\circ \pm 1.3^\circ$ versus $4.1^\circ \pm 1.2^\circ$, $p < 0.001$).²¹ In a study that compared outcomes after cardiothoracic surgery between a group of low phase angle ($< 5.38^\circ$) patients and a group of normal phase angle patients, the participants from the first group had a higher number of post-operative infections, a larger percentage of them were ventilated for more than 12 hours, and had higher rates of mortality. However, after considering other parameters like sex, age, the type of operative procedure, risk assessment, inflammation activity, hypoalbuminemia, heart failure, time of cardiopulmonary bypass, and time of aortic cross-clamp, the phase angle was found not to be statistically significant in correlation with aforementioned complications. Still, the difference between the groups stayed statistically significant in regard to the hospitalization time and the time spent in the intensive care unit.²²

TABLE 3. Odds for developing a complication and probabilities of developing a complication at different values of phase angle

Phase Angle ($^\circ$)	Odds of a complication	Probability of a complication (%)
4.5	5.532	84.7
5.0	3.831	79.3
5.5	2.651	72.6
6.0	1.837	64.8

TABLE 4. Contingency Table with the sensitivity and a specificity of the phase angle

	Phase angle < 5.5	Phase angle ≥ 5.5	
Complication	33 (true positive)	19 (false negative)	52
No complication	7 (false positive)	11 (true negative)	18
	40	30	

In addition, we aimed to find a cut-off phase angle value that can be used to identify high-risk patients that are more likely to have a severe complication after gastrointestinal cancer surgery. Using 5.5° as a cut-off value, we were able to successfully identify these patients in 67% of the cases with a sensitivity of 63% and a specificity of 61%.

Different phase angle cut-off values have appeared in literature to identify patients with lower physiological reserves who are at risk for increased morbidity and mortality. The proposed PA values in literature are 5.5° for patients newly admitted to the hospital²³, 4.73° for patients newly diagnosed with head and neck cancer²⁴, and 5.2° for patients with various cancers before starting radiotherapy²⁵. The phase angle values put forward as a predictor of survival were 4.5° for patients with a non-small cell lung cancer¹⁵, 5.0° for patients with advanced pancreatic cancer²⁶, 5.4° for patients with liver cirrhosis⁵, and 5.57° for patients with colon and rectum cancer¹⁴.

One of the limitations of this study is a relatively heterogeneous group of patients. The patients included in our study had cancer in different locations and stages in abdomen and they were undergoing different treatment protocols. Additionally, our analysis only took into account the pre-operative value of the phase angle. The patients' phase angle might have changed in the peri-operative period, especially on account of nutritional support or further medical interventions. Therefore, the value might not have been representative of the patients overall physical state within the entire observed period. The obtained cut-off phase angle value of 5.5° needs to be considered as a tentative value as it was calculated using a specific sample in one population. Further research is needed to identify the cut-off value for different subtypes of colorectal cancer and to evaluate the validity of our obtained cut-off value in distinct clinical settings.

The primary clinical implication of this study is that the phase angle measurement can assist in identifying GI cancer patients with a higher risk of post-operative complications. This could benefit patients that would otherwise not have been identified. Further research is needed to investigate if nutritional or other medical interventions can significantly alter the phase angle and thus affect surgical outcomes.

References

1. Maggiore Q, Nigrelli S, Ciccarelli C, Grimaldi C, Rossi GA, Michelassi C. Nutritional and prognostic correlates of bioimpedance indexes in hemodialysis patients. *Kidney Int* 1996; **50**: 2103-8. doi: 10.1038/ki.1996.535
2. Baumgartner RN, Chumlea WC, Roche AF. Bioelectric impedance phase angle and body composition. *Am J Clin Nutr* 1988; **48**: 16-23. doi: 10.1093/ajcn/48.1.16
3. Kyle UG, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Gómez JM, et al. Bioelectrical impedance analysis - Part I: review of principles and methods. *Clin Nutr* 2004; **23**: 1226-43. doi: 10.1016/j.clnu.2004.09.012
4. Cornish B. Bioimpedance analysis: scientific background. *Lymphat Res Biol* 2006; **4**: 47-50. doi: 10.1089/lrb.2006.4.47
5. Selberg O, Selberg D. Norms and correlates of bioimpedance phase angle in healthy human subjects, hospitalized patients, and patients with liver cirrhosis. *Eur J Appl Physiol* 2002; **86**: 509-16. doi: 10.1007/s00421-001-0570-4
6. Selberg O, Böttcher J, Tusch G, Pichlmayr R, Henkel E, Müller MJ. Identification of high- and low-risk patients before liver transplantation: a prospective cohort study of nutritional and metabolic parameters in 150 patients. *Hepatology* 1997; **25**: 652-7. doi: 10.1002/hep.510250327
7. Friedl K, Johnson CL, Chumlea WC, Flegal KM, Lukaski HC, Heymsfield SB, et al. Development of bioelectrical impedance analysis prediction equations for body composition with the use of a multicomponent model for use in epidemiologic surveys. *Am J Clin Nutr* 2018; **77**: 331-40. doi: 10.1093/ajcn/77.2.331
8. Anja BW, Danielzik S, Dörhöfer RP, Later W, Wiese S, Müller MJ. Phase angle from bioelectrical impedance analysis: population reference values by age, sex, and body mass index. *JPEN J Parenter Enter Nutr* 2006; **30**: 309-16. doi: 10.1177/0148607106030004309
9. Barbosa-Silva MCG, Barros AJ, Wang J, Heymsfield SB, Pierson RN. Bioelectrical impedance analysis: population reference values for phase angle by age and sex. *Am J Clin Nutr* 2005; **82**: 49-52. doi: 10.1016/j.clnu.2005.05.005
10. Sarhill N, Mahmoud FA, Christie R, Tahir A. Assessment of nutritional status and fluid deficits in advanced cancer. *Am J Hosp Palliat Med* 2003; **20**: 465-73. doi: 10.1177/104990910302000610
11. Barbosa-Silva MCG, Barros AJ, Post CL, Waitzberg DL, Heymsfield SB. Can bioelectrical impedance analysis identify malnutrition in preoperative nutrition assessment? *Nutrition* 2003; **19**: 422-6. doi: 10.1016/s0899-9007(02)00932-2
12. Cox-Reijven PLM, Van Kreel B, Soeters PB. Bioelectrical impedance measurements in patients with gastrointestinal disease: validation of the spectrum approach and a comparison of different methods for screening for nutritional depletion. *Am J Clin Nutr* 2003; **78**: 1111-9. doi: 10.1093/ajcn/78.6.1111
13. Nescolarde L, Piccoli A, Román A, Núñez A, Morales R, Tamayo J, et al. Bioelectrical impedance vector analysis in haemodialysis patients: relation between oedema and mortality. *Physiol Meas* 2004; **25**: 1271-80. doi: 10.1088/0967-3334/25/5/016
14. Gupta D, Lammersfeld CA, Burrows JL, Dahlk SL, Vashi PG, Grutsch JF, et al. Bioelectrical impedance phase angle in clinical practice: implications for prognosis in advanced colorectal cancer. *Am J Clin Nutr* 2004; **80**: 1634-8. doi: 10.1088/0967-3334/25/5/016
15. Silvia T, Piccoli A, Gusella M, Bononi A, Crepaldi G, Ferrazzi E. Altered tissue electric properties in lung cancer patients as detected by bioelectric impedance vector analysis. *Nutrition* 2000; **16**: 120-4. doi: 10.1016/s0899-9007(99)00230-0
16. Schwenk A, Beisenherz A, Römer K, Kremer G, Salzberger B, Elia M. Phase angle from bioelectrical impedance analysis remains an independent predictive marker in HIV-infected patients in the era of highly active antiretroviral treatment. *Am J Clin Nutr* 2000; **72**: 496-501. doi: 10.1093/ajcn/72.2.496
17. Faisy C, Rabbat A, Kouchakji B, Laaban JP. Bioelectrical impedance analysis in estimating nutritional status and outcome of patients with chronic obstructive pulmonary disease and acute respiratory failure. *Intensive Care Med* 1999; **26**: 518-25. doi: 10.1007/s001340051198
18. Kyle UG, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Gómez JM, et al. Bioelectrical impedance analysis - Part II: utilization in clinical practice. *Clin Nutr* 2004; **23**: 1430-53. doi: 10.1016/j.clnu.2004.09.012

19. Katayama H, Kurokawa Y, Nakamura K, Ito H, Kanemitsu Y, Masuda N, et al. Extended Clavien-Dindo classification of surgical complications: Japan Clinical Oncology Group postoperative complications criteria. *Surg Today* 2016; **46**: 668-85. doi: 10.1007/s00595-015-1236-x
20. Chung HY. Bioelectrical impedance analysis for prediction of early complications after gastrectomy in elderly patients with gastric cancer: the phase angle measured using bioelectrical impedance analysis. *J Gastric Cancer* 2019; **19**: 278-89. doi: 10.5230/jgc.2019.19.e22
21. Stapel SN, Looijaard WGPM, Dekker IM, Girbes ARJ, Weijs PJM, Oudemans-Van Straaten HM. Bioelectrical impedance analysis-derived phase angle at admission as a predictor of 90-day mortality in intensive care patients. *Eur J Clin Nutr* 2018; **72**: 1019-25. doi: 10.1038/s41430-018-0167-1
22. Visser M, van Venrooij LMW, Wanders DCM, de Vos R, Wisselink W, van Leeuwen PAM, et al. The bioelectrical impedance phase angle as an indicator of undernutrition and adverse clinical outcome in cardiac surgical patients. *Clin Nutr* 2012; **31**: 981-6. doi: 10.1016/j.clnu.2012.05.002
23. Kyle UG, Soundar EP, Genton L, Pichard C. Can phase angle determined by bioelectrical impedance analysis assess nutritional risk? A comparison between healthy and hospitalized subjects. *Clin Nutr* 2012; **31**: 875-81. doi: 10.1016/j.clnu.2012.04.002
24. Matecka-Massalska T, Mlak R, Smolen A, Morshed K. Bioelectrical impedance phase angle and subjective global assessment in detecting malnutrition among newly diagnosed head and neck cancer patients. *Eur Arch Otorhinolaryngol* 2016; **273**: 1299-305. doi: 10.1007/s00405-015-3626-5
25. Motta RST, Castanho IA, Velarde LGC. Cutoff point of the phase angle in pre-radiotherapy cancer patients. *Nutr Hosp* 2015; **32**: 2253-60. doi: 10.3305/nh.2015.32.5.9626
26. Gupta D, Lis CG, Dahlk SL, Vashi PG, Grutsch JF, Lammersfeld CA. Bioelectrical impedance phase angle as a prognostic indicator in advanced pancreatic cancer. *Br J Nutr* 2004; **92**: 957-62. doi: 10.1079/BJN20041292