

Characterising CCA Sylow cyclic groups whose order is not divisible by four*

Luke Morgan[†]

Centre for the Mathematics of Symmetry and Computation, School of Mathematics and Statistics (M019), The University of Western Australia, Crawley, 6009, Australia

Joy Morris[‡]

Department of Mathematics and Computer Science, University of Lethbridge, Lethbridge, AB T1K 3M4, Canada

Gabriel Verret[§]

Centre for the Mathematics of Symmetry and Computation, School of Mathematics and Statistics (M019), The University of Western Australia, Crawley, 6009, Australia

Received 21 February 2017, accepted 5 April 2017, published online 10 May 2017

Abstract

A Cayley graph on a group G has a natural edge-colouring. We say that such a graph is CCA if every automorphism of the graph that preserves this edge-colouring is an element of the normaliser of the regular representation of G . A group G is then said to be CCA if every connected Cayley graph on G is CCA.

Our main result is a characterisation of non-CCA graphs on groups that are Sylow cyclic and whose order is not divisible by four. We also provide several new constructions of non-CCA graphs.

Keywords: CCA problem, Cayley graphs, edge-colouring, Sylow cyclic groups.

Math. Subj. Class.: 05C25

*We would like to thank Gordon Royle for his help with some of the computations. We would also like to thank the anonymous referee for a number of helpful suggestions.

[†]This research was supported by the Australian Research Council grant DE160100081.

[‡]This research was supported by the Natural Science and Engineering Research Council of Canada, and by the Cheryl E. Praeger Fellowship from The University of Western Australia.

[§]Current address: Department of Mathematics, The University of Auckland, Private Bag 92019, Auckland 1142, New Zealand. This research was supported by the Australian Research Council grant DE130101001.

E-mail addresses: luke.morgan@uwa.edu.au (Luke Morgan), joy.morris@uleth.ca (Joy Morris), g.verret@auckland.ac.nz (Gabriel Verret)

1 Introduction

All groups and all graphs in this paper are finite. Let G be a group and let S be an inverse-closed subset of G . The *Cayley graph* of G with respect to S is the edge-coloured graph $\text{Cay}(G, S)$ with vertex-set G and, for every $g \in G$ and $s \in S$, an edge $\{g, sg\}$ with colour $\{s, s^{-1}\}$. Its group of colour-preserving automorphisms is denoted $\text{Aut}_c(\text{Cay}(G, S))$. Let $\text{Aut}_{\pm 1}(G, S) = \{\alpha \in \text{Aut}(G) : s^\alpha \in \{s, s^{-1}\} \text{ for all } s \in S\}$. It is easy to see that $G_R \rtimes \text{Aut}_{\pm 1}(G, S) \leq \text{Aut}_c(\text{Cay}(G, S))$, where G_R is the right-regular representation of G .

Definition 1.1 ([5]). The Cayley graph $\text{Cay}(G, S)$ is *CCA* (Cayley colour automorphism) if $\text{Aut}_c(\text{Cay}(G, S)) = G_R \rtimes \text{Aut}_{\pm 1}(G, S)$. The group G is *CCA* if every connected Cayley graph on G is CCA.

In other words, a Cayley graph is CCA if and only if the colour-preserving graph automorphisms are exactly the “obvious” ones. The terminology we use for this problem largely comes from [5]. Other papers that study this problem include [2, 3, 4, 8].

Note that $\text{Cay}(G, S)$ is connected if and only if S generates G . It is also easy to see that $G_R \rtimes \text{Aut}_{\pm 1}(G, S)$ is precisely the normaliser of G_R in $\text{Aut}_c(\text{Cay}(G, S))$. In particular, $\text{Cay}(G, S)$ is CCA if and only if G_R is normal in $\text{Aut}_c(\text{Cay}(G, S))$, c.f. [5, Remark 6.2].

In Section 2, we introduce some basic terminology and recall a few previous results on the CCA property. In Section 3, we consider wreath products of permutation groups, and produce conditions that are sufficient to determine when such a product is a non-CCA group. This generalises results from [5]. In Section 4, we give some new constructions for non-CCA graphs.

Finally, in Section 5, we obtain a characterisation of non-CCA groups whose order is not divisible by four, in which every Sylow subgroup is cyclic. This generalises the work of [3], which dealt with the case of groups of odd squarefree order.

2 Preliminaries

The identity of a group G is denoted 1_G , or simply 1 if there is no risk of confusion. We denote a dihedral group of order $2n$ by D_n , while Q_8 denotes the quaternion group of order 8 with elements $\{\pm 1, \pm i, \pm j, \pm k\}$ and multiplication defined as usual.

We now state some preliminary results and introduce some terminology related to Cayley graphs. Let Γ be a graph and let v be a vertex of Γ . The neighbourhood of v is denoted by $\Gamma(v)$. If A is a group of automorphisms of Γ , then the permutation group induced by the vertex-stabiliser A_v on the neighbourhood of v is denoted $A_v^{\Gamma(v)}$.

Lemma 2.1 ([5, Lemma 6.3]). *The vertex-stabiliser in the colour-preserving group of automorphisms of a connected Cayley graph is a 2-group.*

Definition 2.2. Let G be a group, let $\Gamma = \text{Cay}(G, S)$ and let N be a normal subgroup of G . The *quotient graph* Γ/N is $\text{Cay}(G/N, S/N)$, where $S/N = \{sN : s \in S\}$.

Lemma 2.3 ([3, Lemma 3.4]). *Let A be a colour-preserving group of automorphisms of $\text{Cay}(G, S)$, let N be a normal subgroup of A and let K be the kernel of the action of A on the N -orbits. If $N \leq G$, then A/K is a colour-preserving group of automorphisms of Γ/N .*

Lemma 2.4. *Let $\Gamma = \text{Cay}(G, S)$, let A be a colour-preserving group of automorphisms of Γ , let N be a normal 2-subgroup of A and let K be the kernel of the action of A on the N -orbits. If $K_v \neq 1$, then S contains an element whose order is a power of 2 that is at least 4.*

Proof. Let $v_0 = v$. Since $K_{v_0} \neq 1$, $K_{v_0}^{\Gamma(v_0)} \neq 1$ and there exists $k \in K_{v_0}$ and a neighbour u_0 of v_0 such that $u_0^k \neq u_0$. Let $u_1 = u_0^k$. Note that $K_{u_1} \neq K_{v_0}$ hence there exists $\ell \in K_{u_1}$ such that $v_0^\ell \neq v_0$. Let $v_1 = v_0^\ell$. Repeating this process, we get a monochromatic cycle $C = (u_0, v_0, u_1, v_1, \dots)$ of length at least 3. By construction, $u_i \in u_0^K = u_0^N$ and $v_i \in v_0^K = v_0^N$ for all i . In particular, $|C \cap v_0^N| \in \{|C|, |C|/2\}$. Since each vertex of Γ lies in a unique monochromatic cycle of a given colour, C is a block for A . On the other hand, v_0^N is also a block for A and thus so is $C \cap v_0^N$. It follows that $|C \cap v_0^N|$ divides $|N|$ which is a power of 2. This implies that $|C|$ is also a power of 2. Since $|C| \geq 3$, $|C|$ is divisible by 4 and the result follows from the fact that C is monochromatic. \square

For a group H , let $H^2 := \langle x^2 \mid x \in H \rangle$. The following lemma is inspired by an argument contained within [5, Theorem 6.8].

Lemma 2.5. *Let $\Gamma = \text{Cay}(G, S)$ be connected, let A be a colour-preserving group of automorphisms of Γ that is normalised by G and let v be a vertex of Γ . If A_v has a subgroup U such that $U \leq (A_v)^2$ and no other subgroup of A_v is isomorphic to U , then $U = 1$. In particular, A_v is isomorphic to neither \mathbb{Z}_{2^n} for $n \geq 2$ nor isomorphic to D_{2^n} for $n \geq 3$.*

Proof. Since A is colour-preserving, $A_v^{\Gamma(v)}$ is an elementary abelian 2-group. Since $U \leq (A_v)^2$, it follows that U fixes all the neighbours of v . Let $s \in S$. Since A is normalised by G , we have $U^s \leq A$ and, by the previous observation, $U^s \leq A_v$. As A_v has a unique subgroup isomorphic to U , we must have $U = U^s$. Since this holds for every $s \in S$, U is normalised by G . As G is transitive and U fixes v , this implies that $U = 1$.

The second part of the lemma follows from the first. Indeed, if A_v is isomorphic to \mathbb{Z}_{2^n} for $n \geq 2$ or to D_{2^n} for $n \geq 3$, then $(A_v)^2$ is non-trivial and is the unique cyclic subgroup of its order. \square

3 Wreath products

Proposition 3.1. *Let H be a permutation group on a set Ω , let G be a group and let $X = G \wr_\Omega H$. If*

- (i) *there is an inverse-closed generating set S for G and a non-identity bijection $\tau: G \rightarrow G$ such that τ fixes 1, and $\tau(sg) = s^{\pm 1}\tau(g)$ for every $g \in G$ and every $s \in S$, and*
- (ii) *either H is nontrivial or $\tau \notin \text{Aut}(G)$,*

then X is non-CCA.

Proof. Let $m = |\Omega|$ and write $\Omega = \{1, \dots, m\}$ such that, if H is nontrivial, then 1 is not fixed by H .

Write $X = H \ltimes (G_1 \times \dots \times G_m)$. Note that, if $g \in G_i$ and $h \in H$, then $g^h \in G_{i^h}$. Without loss of generality, we may assume that $1_G \notin S$. Let S_i be the subset of G_i

corresponding to S , let $T = (H - \{1_H\}) \cup S_1 \cup \dots \cup S_m$ and let $\Gamma = \text{Cay}(X, T)$. Note that T generates X hence Γ is connected. We will show that Γ is non-CCA.

Define $\tau': X \rightarrow X$ by $\tau': hg_1g_2 \cdots g_m \mapsto h\tau(g_1)g_2 \cdots g_m$, where $g_i \in G_i$ and $h \in H$. Let v be a vertex of Γ and let $s \in T$. We will show that $\tau'(sv) = s^{\pm 1}\tau'(v)$ and hence τ' is a colour-preserving automorphism of Γ . Write $v = hg_1 \cdots g_m$ with $h \in H$ and $g_i \in G_i$. Let $g = g_1 \cdots g_m$. Note that $\tau'(v) = \tau'(hg) = h\tau'(g)$. If $s \in H$, then $\tau'(sv) = \tau'(shg) = sh\tau'(g) = s\tau'(v)$. Suppose now that $s \in S_i$ for some $i \in \Omega$. If $i^h \neq 1$, then

$$\begin{aligned}\tau'(s^h g) &= \tau'(g_1 \cdots s^h g_{i^h} \cdots g_m) = \tau(g_1)g_2 \cdots s^h g_{i^h} \cdots g_m = \\ &= s^h \tau(g_1)g_2 \cdots g_m = s^h \tau'(g).\end{aligned}$$

If $i^h = 1$, then $s^h \in S_1 \subseteq T$ and

$$\tau'(s^h g) = \tau'(s^h g_1 \cdots g_m) = \tau(s^h g_1)g_2 \cdots g_m = (s^h)^{\pm 1} \tau(g_1)g_2 \cdots g_m = (s^h)^{\pm 1} \tau'(g).$$

Either way, we have

$$\tau'(sv) = \tau'(hs^h g) = h\tau'(s^h g) = h(s^h)^{\pm 1} \tau'(g) = s^{\pm 1} h\tau'(g) = s^{\pm 1} \tau'(hg) = s^{\pm 1} \tau'(v).$$

This completes the proof that τ' is a colour-preserving automorphism of Γ . It remains to show that τ' is not a group automorphism of X . (Note that τ' fixes 1_X , so if $\tau' \in X_R \rtimes \text{Aut}_{\pm 1}(X, T)$, then $\tau' \in \text{Aut}(X)$.)

If H is nontrivial, then, since 1 is not fixed by H , there exists $h \in H$ such that $1^h \neq 1$. Let g be an element of G_1 that is not fixed by τ . We have $\tau'(gh) = \tau'(hg^h) = hg^h = gh$ but $\tau'(g)\tau'(h) = \tau(g)h$. Since $g \neq \tau(g)$, τ' is not an automorphism of X .

If H is trivial and $\tau \notin \text{Aut}(G)$, then there exist $g_1, g_2 \in G$ such that $\tau(g_1g_2) \neq \tau(g_1)\tau(g_2)$. Applying τ' to the corresponding elements of G_1 shows that τ' is not an automorphism of X . This completes the proof. \square

We now obtain a few corollaries of Proposition 3.1.

Corollary 3.2. *Let H be a permutation group on a set Ω and let G be a group. If G is non-CCA, then $G \wr_{\Omega} H$ is non-CCA.*

Proof. Since G is non-CCA, there exists a colour-preserving graph automorphism τ of a Cayley graph $\text{Cay}(G, S)$ such that $\tau(1_G) = 1_G$ but τ does not normalise G_R . Since τ is colour-preserving, $\tau(sg) = s^{\pm 1}\tau(g)$ for every $g \in G$ and every $s \in S$. Finally, since τ does not normalise G_R , we have $\tau \notin \text{Aut}(G)$ and the result follows from Proposition 3.1. \square

Corollary 3.3. *Let H be a nontrivial permutation group on a set Ω and let G be a group. If $G = B \rtimes A$, where A is abelian of exponent greater than 2, then $G \wr_{\Omega} H$ is non-CCA.*

Proof. Every element of G can be written uniquely as ba with $a \in A$ and $b \in B$. Let τ be the permutation of G mapping ba to ba^{-1} . Clearly, τ fixes 1_G but, since A has exponent greater than 2, τ is not the identity. Let $S = (A \cup B) - \{1_G\}$. Note that S is an inverse-closed generating set for G . Let $g \in G$, let $s \in S$ and write $g = ba$ with $a \in A$ and $b \in B$.

If $s \in B$, then $\tau(sg) = \tau(sba) = sba^{-1} = s\tau(ba) = s\tau(g)$. Otherwise, $s \in A$, $s^b \in A$ and

$$\tau(sg) = \tau(sba) = \tau(bs^b a) = b(s^b a)^{-1} = b(s^b)^{-1} a^{-1} = s^{-1} b a^{-1} = s^{-1} \tau(g).$$

The result then follows from Proposition 3.1, since H is nontrivial. \square

Corollary 3.4. *Let H be a permutation group on a set Ω and let G be a group. If*

- (i) G has exponent greater than 2,
- (ii) H is nontrivial when G is abelian, and
- (iii) G has a generating set S with the property that $s^g = s^{\pm 1}$ for every $s \in S$ and $g \in G$,

then $G \wr_{\Omega} H$ is non-CCA.

Proof. We can assume without loss of generality that S is inverse-closed. Let τ be the permutation of G that maps every element to its inverse. For every $s \in S$ and $g \in G$, we have $s^g = s^{\pm 1}$ and thus $\tau(sg) = g^{-1} s^{-1} = s^{\pm 1} g^{-1} = s^{\pm 1} \tau(g)$. Since G has exponent greater than 2, τ is not the identity. If H is trivial, then G is non-abelian so that τ is not an automorphism of G . The result then follows from Proposition 3.1. \square

In view of Corollary 3.4, it would be interesting to determine the groups G such that G has a generating set S with the property that $s^g = s^{\pm 1}$ for every $s \in S$ and $g \in G$. This family of groups includes abelian groups and Q_8 . This family is closed under central products but it also includes examples which do not arise as central products of smaller groups in the family, for example the extraspecial group of order 32 and minus type.

4 A few constructions for non-CCA graphs

In this section, we will describe a few constructions which yield non-CCA Cayley graphs. For a group G , let K_G denote $\text{Cay}(G, G - \{1\})$, the complete Cayley graph on G . We will need a result which tells us when $G_R < \text{Aut}_c(K_G)$. First we state some definitions.

Definition 4.1. Let A be an abelian group of exponent greater than 2, and define a map $\iota: A \rightarrow A$ by $\iota(a) = a^{-1}$ for every $a \in A$. The *generalised dihedral group* over A is $\text{Dih}(A) = A \rtimes \langle \iota \rangle$.

Definition 4.2. Let A be an abelian group of even order and of exponent greater than 2, and let y be an element of A of order 2. The *generalised dicyclic group* over A is $\text{Dic}(A, y) := \langle A, x \mid x^2 = y, a^x = a^{-1} \forall a \in A \rangle$. Let ι be the permutation of $\text{Dic}(A, y)$ that fixes A pointwise and maps every element of the coset Ax to its inverse.

It is not hard to check that ι is an automorphism of $\text{Dic}(A, y)$.

Definition 4.3. For $\alpha \in \{i, j, k\}$, let $S_\alpha = \{\pm \alpha\} \times \mathbb{Z}_2^n$ and let σ_α be the permutation of $Q_8 \times \mathbb{Z}_2^n$ that inverts every element of S_α and fixes every other element.

Theorem 4.4 ([2], Classification Theorem). *If G is a group, then $G_R < \text{Aut}_c(K_G)$ if and only if one of the following occurs:*

- 1. G is abelian but is not an elementary abelian 2-group, and $\text{Aut}_c(K_G) = \text{Dih}(G)$,

2. G is generalised dicyclic but not of the form $Q_8 \times \mathbb{Z}_2^n$, and $\text{Aut}_c(K_G) = G_R \rtimes \langle \iota \rangle$, where ι is as in Definition 4.2, or
3. $G \cong Q_8 \times \mathbb{Z}_2^n$ and $\text{Aut}_c(K_G) = \langle G_R, \sigma_i, \sigma_j, \sigma_k \rangle$, where $\sigma_i, \sigma_j, \sigma_k$ are as in Definition 4.3.

Definition 4.5. Let B be a permutation group and let G be a regular subgroup of B . We say that (G, B) is a *complete colour pair* if G is as in the conclusion of Theorem 4.4 and $B \leq \text{Aut}_c(K_G)$.

For a graph Γ , let $\mathcal{L}(\Gamma)$ denote its line graph.

Proposition 4.6. Let Γ be a connected bipartite G -edge-regular graph. If H is a group of automorphisms of Γ such that:

- (i) $G \leq H$,
- (ii) the orbits of H on the vertex-set of Γ are exactly the biparts, and
- (iii) for every vertex v of Γ , either

- (a) $G_v^{\Gamma(v)} = H_v^{\Gamma(v)}$, or
- (b) $(G_v^{\Gamma(v)}, H_v^{\Gamma(v)})$ is a complete colour pair,

then H is a colour-preserving group of automorphisms of $\mathcal{L}(\Gamma)$ viewed as a Cayley graph on G .

Proof. Since G acts regularly on edges of Γ , its induced action on $\mathcal{L}(\Gamma)$ is regular on vertices. Vertices of Γ induce cliques in $\mathcal{L}(\Gamma)$, which we call *special*. Clearly, H has exactly two orbits on special cliques. Moreover, special cliques partition the edges of $\mathcal{L}(\Gamma)$, and each vertex of $\mathcal{L}(\Gamma)$ is in exactly two special cliques, one from each H -orbit. Since $G \leq H$, the set of edge-colours appearing in special cliques from different H -orbits is disjoint.

Let v be a vertex of Γ and let C be the corresponding special clique of $\mathcal{L}(\Gamma)$. Note that $H_v^{\Gamma(v)}$ is permutation isomorphic to H_C^C , while $G_v^{\Gamma(v)} \cong G_C^C \cong G_C$. Since G is vertex-regular on $\mathcal{L}(\Gamma)$, G_C is regular on C and thus C can be viewed as a complete Cayley graph on G_C . If $(G_v^{\Gamma(v)}, H_v^{\Gamma(v)})$ is a complete colour pair, Theorem 4.4 implies that H_C^C is colour-preserving. If $G_v^{\Gamma(v)} = H_v^{\Gamma(v)}$, then since G is colour-preserving, so is H_C^C . Since G acts transitively on the special cliques within an H -orbit and G is colour-preserving, it follows that H is colour-preserving. \square

Remark 4.7. In the proof of Proposition 4.6, we only use one direction of Theorem 4.4, namely that if G appears in Theorem 4.4, then $G_R < \text{Aut}_c(K_G)$. The converse is not used here, but it can help to identify situations where Proposition 4.6 can be used to construct non-CCA graphs.

Example 4.8. Let Γ be the Heawood graph and let H be the bipart-preserving subgroup of $\text{Aut}(\Gamma)$. Note that $H \cong \text{PSL}(2, 7)$ and H contains an edge-regular subgroup G isomorphic to F_{21} , the Frobenius group of order 21. Moreover, for every vertex v of Γ , we have $G_v^{\Gamma(v)} \cong \mathbb{Z}_3$ while $H_v^{\Gamma(v)} \cong D_3$ and (\mathbb{Z}_3, D_3) is a complete colour pair. By Proposition 4.6,

H is a colour-preserving group of automorphisms of $\mathcal{L}(\Gamma)$ viewed as a Cayley graph on G . Since G is not normal in H , it follows that $\mathcal{L}(\Gamma)$ is a non-CCA graph and so F_{21} is a non-CCA group.

Example 4.8 was previously studied in [3] and [5], under a slightly different guise.

Example 4.9. Let $A \cong Q_8 \times \mathbb{Z}_2^m$ and $B = \text{Aut}_c(K_A) = \langle A_R, \sigma_i, \sigma_j, \sigma_k \rangle$. Then A is not normal in B , and by Theorem 4.4(3), (A, B) is a complete colour pair. Let $n = |A|$ and let $K_{n,n}$ be the complete bipartite graph of order $2n$. Let $G = A \times A$ and let $H = B \times B$. By Proposition 4.6, H is a colour-preserving group of automorphisms of $\mathcal{L}(K_{n,n})$ viewed as a Cayley graph on G . Since A is not normal in B , G is not normal in H hence $\mathcal{L}(K_{n,n})$ is a non-CCA graph and so G is a non-CCA group.

For a graph Γ , let $\mathcal{S}(\Gamma)$ denote its subdivision graph.

Corollary 4.10. *Let Γ be a connected G -arc-regular graph. If H is a group of automorphisms of Γ such that:*

- (i) $G \leq H$, and
- (ii) $(G_v^{\Gamma(v)}, H_v^{\Gamma(v)})$ is a complete colour pair for every vertex v of Γ ,

then H is a colour-preserving group of automorphisms of $\mathcal{L}(\mathcal{S}(\Gamma))$ viewed as a Cayley graph on G .

Proof. Let $\Gamma' = \mathcal{S}(\Gamma)$. We show that Proposition 4.6 applies to Γ' . Clearly, Γ' is bipartite and G acts on it faithfully and edge-regularly. It is also obvious that, in its induced action on Γ' , H must preserve the biparts of Γ' . Finally, let x be a vertex of Γ' . If x arose from a vertex v of Γ , then we have that $A_v^{\Gamma(v)}$ is permutation isomorphic to $A_x^{\Gamma'(x)}$ for every $A \leq \text{Aut}(\Gamma)$. Since $(G_v^{\Gamma(v)}, H_v^{\Gamma(v)})$ is a complete colour pair, so is $(G_x^{\Gamma'(x)}, H_x^{\Gamma'(x)})$. If x arose from an edge of Γ , then x has valency 2 and, since G is arc-transitive, $G_x^{\Gamma'(x)} = H_x^{\Gamma'(x)} \cong \mathbb{Z}_2$ and $(G_x^{\Gamma'(x)}, H_x^{\Gamma'(x)})$ is a complete colour pair. \square

Example 4.11. Let Γ be the Heawood graph and let $H = \text{Aut}(\Gamma)$. Note that H contains an arc-regular subgroup G isomorphic to $\text{AGL}(1, 7)$. Moreover, for every vertex v of Γ , we have $G_v^{\Gamma(v)} \cong \mathbb{Z}_3$ while $H_v^{\Gamma(v)} \cong D_3$. By Corollary 4.10, H is a colour-preserving group of automorphisms of $\mathcal{L}(\mathcal{S}(\Gamma))$ viewed as a Cayley graph on G . Since G is not normal in H , it follows that Γ is a non-CCA graph and so $\text{AGL}(1, 7)$ is a non-CCA group.

Remark 4.12. In fact, $\text{AGL}(1, 7)$ is a Sylow cyclic group whose order is not divisible by four, so Example 4.11 will appear again in our characterisation of non-CCA groups of this sort, in Section 5. However, the construction we have just presented is very different from the approach we use in that section.

5 Sylow cyclic and order not divisible by four

We first introduce some notation that will be useful throughout this section. Recall that $\text{PGL}(2, 7)$ has a unique conjugacy class of subgroups isomorphic to $\text{AGL}(1, 7)$. The intersection of such a subgroup with the socle $\text{PSL}(2, 7)$ is a Frobenius group of order 21 which we will denote F_{21} . We say that a group G is *Sylow cyclic* if, for every prime p , the Sylow p -subgroups of G are cyclic.

Our aim in this section is to characterise both the non-CCA Sylow cyclic groups whose order is not divisible by four, and the structure of the corresponding colour-preserving automorphism groups for non-CCA graphs.

Theorem 5.1. *Let G be a Sylow cyclic group whose order is not divisible by four, let $\Gamma = \text{Cay}(G, S)$, let A be a colour-preserving group of automorphisms of Γ , let R be a Sylow 2-subgroup of G and let r be a generator of R . If G is not normal in A , then $G = (F \times H) \rtimes R$ and $A = (T \times J) \rtimes R$, where the following hold:*

- (i) $\text{PSL}(2, 7) \cong T \trianglelefteq A$,
- (ii) $T \cap G = F \cong \text{F}_{21}$,
- (iii) $J \cap G = H \trianglelefteq J \trianglelefteq A$,
- (iv) H is self-centralising in J ,
- (v) J splits over H ,
- (vi) H is normal in A .

Proof. To avoid ambiguity, for $g \in G$, we write $[g]$ for the vertex of Γ corresponding to g and, for $X \subseteq G$, we write $[X]$ for $\{[x] : x \in X\}$.

Let P be a Sylow 2-subgroup of A containing R . By Lemma 2.1, $A_{[1]}$ is a 2-group. Up to relabelling, we may assume that $A_{[1]} \leq P$. Since G is regular, we have $A = GA_{[1]}$ and $|A| = |G||A_{[1]}|$. Note that (v) and (vi) follow from the rest of the claims. Indeed, H must have odd order and, since $|A : G|$ is a power of 2, so is $|J : H|$ and thus $J = H \rtimes (P \cap J)$. As H has odd order and is normal in J , it must be characteristic in J and thus normal in A .

Since $|G|$ is not divisible by 4, it follows that G has a characteristic subgroup G_2 of odd order such that $G = G_2 \rtimes R$. By order considerations, we have $A = G_2P$.

Case 1: There is no minimal normal subgroup of A of odd order.

In this case, we have that $\text{soc}(A) = T_1 \times \cdots \times T_k \times B$, where $\text{soc}(A)$ is the socle of A , the T_i s are non-abelian simple groups, and B is an elementary abelian 2-group. Recall that $A = G_2P$, that is, A has a 2-complement. Since this property is inherited by normal subgroups, $\text{soc}(A)$ and T_i also have 2-complements for every i . This implies that, for every i , $T_i \cong \text{PSL}(2, p)$ for some Mersenne prime p (see [7, Theorem 1.3] for example). Now, $|T_i|$ is divisible by 3 but the Sylow 3-subgroup of $\text{soc}(A)$ is cyclic (since $|A : G|$ is a power of 2 and G is Sylow cyclic) so that $k = 1$. Let $T = T_1$. Suppose that $p > 7$ and hence $p \geq 31$. Note that $T_{[1]}$ has index at most 2 in some Sylow 2-subgroup of T which is isomorphic to $D_{(p+1)/2}$. It follows that $T_{[1]}$ has order at least $(p+1)/2$ and is either dihedral or cyclic. Since $p \geq 31$, this implies that $T_{[1]}$ contains a unique cyclic subgroup of order $(p+1)/8$, say U , and U is contained in $(T_{[1]})^2$. By Lemma 2.5, $U = 1$, which is a contradiction. It follows that $p = 7$ and $T \cong \text{PSL}(2, 7)$.

Let $O_2(A)$ be the largest normal 2-subgroup of A . If $O_2(A) = 1$, then $\text{soc}(A) = T \cong \text{PSL}(2, 7)$ and A is isomorphic to one of $\text{PSL}(2, 7)$ or $\text{PGL}(2, 7)$. If $A \cong \text{PSL}(2, 7)$, then, as F_{21} is the only proper subgroup of $\text{PSL}(2, 7)$ with index a power of 2, $G \cong \text{F}_{21}$ and the theorem holds. If $A \cong \text{PGL}(2, 7)$, then, for the same reason, G is isomorphic to either F_{21} or $\text{AGL}(1, 7)$. If $G \cong \text{F}_{21}$, then $A_{[1]}$ must be a Sylow 2-subgroup of A and thus isomorphic to D_8 . In particular, $A_{[1]}$ contains a unique cyclic subgroup of order 4 and this subgroup is contained in $(A_{[1]})^2$. This contradicts Lemma 2.5. We must therefore have $G \cong \text{AGL}(1, 7)$ and again the theorem holds.

We now assume that $O_2(A) \neq 1$. In particular, the orbits of $O_2(A)$ are of equal length, which is a power of 2 greater than 1. It follows that $|O_2(A) : O_2(A)_{[1]}| = |[1]^{O_2(A)}| = 2$. Let K be the kernel of the action of A on the $O_2(A)$ -orbits. By Lemma 2.4, K is semiregular hence so is $O_2(A)$. It follows that $|O_2(A)| = 2$ and $O_2(A)$ is central in A . This implies that $B = O_2(A)$, hence $\text{soc}(A) = T \times O_2(A)$. Now, $A_{[1]}$ is a complement for $O_2(A)$ in P , so by Gaschutz' Theorem (see for example [6, 3.3.2]), $O_2(A)$ has a complement in A .

Clearly, $O_2(A) \leq C_A(T)$. We show that equality holds. Suppose, on the contrary, that $O_2(A) < C_A(T)$. Since $C_A(T)$ is normal in A , $C_A(T)/O_2(A)$ must contain a minimal normal subgroup of $A/O_2(A)$, say $Y/O_2(A)$. Since $O_2(A)$ has a complement in A , $O_2(A)$ has a complement in Y , say Z . Thus $Y = O_2(A) \times Z$ and Z is isomorphic to $Y/O_2(A)$ which is a minimal normal subgroup of $A/O_2(A)$ and therefore either an elementary abelian group of odd order, or a product of non-abelian simple groups. It follows that Z is characteristic in Y and thus normal in A . Since the action of A by conjugation on Z and on $Y/O_2(A)$ are equivalent, we see that Z is a minimal normal subgroup of A . The only possibility is that $Z = T$ but, since T has trivial centre, this contradicts the fact that $Z \leq Y \leq C_A(T)$. This concludes our proof that $C_A(T) = O_2(A) \cong \mathbb{Z}_2$.

As $O_2(A)$ has a complement in A , it follows that A is isomorphic to one of $\text{PSL}(2, 7) \times \mathbb{Z}_2$ or $\text{PGL}(2, 7) \times \mathbb{Z}_2$. Suppose first that $A \cong \text{PSL}(2, 7) \times \mathbb{Z}_2$. Since G has even order, is not normal in A and has index a power of 2, we must have $G \cong F_{21} \times \mathbb{Z}_2$ and the theorem holds with $H = J = 1$. Finally, suppose that $A \cong \text{PGL}(2, 7) \times \mathbb{Z}_2$. In particular, $P = Q \times O_2(A)$ where $Q \cong D_8$. Note that $|P : A_{[1]}| = 2$ and $A_{[1]} \cap O_2(A) = 1$ hence $A_{[1]} \cong P/O_2(A) \cong D_8$. This contradicts Lemma 2.5.

Case 2: There exists a minimal normal subgroup of A of odd order.

Let N be a minimal normal subgroup of odd order, that is, $|N|$ is a power of some odd prime p . Let K be the kernel of the action of A on the set of N -orbits. Since the N -orbits have odd size and $K_{[1]} \leq A_{[1]}$ is a 2-group, $K_{[1]}$ must fix at least one point in every N -orbit. For each N -orbit B , pick $b \in G$ such that $K_{[1]} = K_{[b]}$ and $B = [b]^N$. Now the kernel of the action of K on $[1]^N$ is $K_{([1]^N)} = \bigcap_{n \in N} (K_{[1]})^n = \bigcap_{n \in N} (K_{[b]})^n = K_{([b]^N)}$. It follows that $K_{([1]^N)}$ fixes every vertex of Γ , and so K acts faithfully on $[1]^N$. Moreover, $N_{[1]} = 1$ hence $K = N \rtimes K_{[1]}$. As $|A : G|$ is a power of 2, G contains a Sylow p -subgroup of A . Since N is normal in A , it is contained in every Sylow subgroup of A , and thus $N \leq G$. We therefore have $GK = GNK_{[1]} = GK_{[1]}$. Since G is Sylow cyclic and N is elementary abelian, we must have $|N| = p$.

If $K_{[1]} \neq 1$, then $K_{[1]}$ must move a neighbour of $[1]$, say $[s]$ for some non-involution $s \in S$. It follows that $K_{[s]}$ must move $[1]$, necessarily to $[s^2]$ since K is colour-preserving, and thus $[s^2] \in [1]^N$. Let C be the cycle containing $[1]$ with edge-label $\{s, s^{-1}\}$. We have shown that $[1], [s^2] \in [1]^N \cap C$ and hence $|[1]^N \cap C| \geq 2$. Since $[1]^N$ and C are both blocks for the action of G , the former of prime order, it follows that $[1]^N \cap C = [1]^N$, that is $[1]^N \subseteq C$. Since K acts faithfully on $[1]^N$, $K_{[1]}$ acts faithfully on C and thus $|GK : G| = |K_{[1]}| \leq 2$. It follows that G is normal in GK , $GK = G \rtimes K_{[1]}$ and either $K = N \cong \mathbb{Z}_p$ or $K \cong D_p$.

Suppose that GK/K is normal in A/K and hence GK is normal in A . We show that this implies that G is normal in A , which is a contradiction. This is trivial if $G = GK$ hence we assume that $|GK : G| = 2$ and $K \cong D_p$. If G has odd order, then it is characteristic in GK and thus normal in A . We may thus assume that G has even order. Recall that G_2 is a characteristic subgroup of index 2 in G , hence G_2 is normal in GK and, since $|GK : G_2| = 4$, we have that G_2 is characteristic in GK and thus normal in A . Note that

G and $G_2 \rtimes K_{[1]}$ both have index two in GK but $G_2 \rtimes K_{[1]}$ is not semiregular, hence they are not conjugate in A . In particular, $G_2 \rtimes K_{[1]}$ and G are distinct index two subgroups of GK and thus GK/G_2 is elementary abelian of order 4. Let X be the centraliser of N in GK . Since $N \cong \mathbb{Z}_p$, $\text{Aut}(N)$ is cyclic hence GK/X is cyclic and X is not contained in G_2 . Since N , G_2 and GK are normal in A , so is XG_2 . If $XG_2 = G$, then we are done. We thus assume that this is not the case. Note that $|XG_2 : X| = |G_2 : C_{G_2}(N)|$ is odd, hence every Sylow 2-subgroup of XG_2 centralises N . Since $K_{[1]}$ has order 2 but does not centralise N , $G_2 \rtimes K_{[1]}$ is not contained in XG_2 . We thus conclude that G , $G_2K_{[1]}$ and XG_2 are the three index two subgroups of GK containing G_2 . One of them is normal in A , and we have seen that the other two are not conjugate in A . It follows that all three are normal in A . In particular, G is normal in A , a contradiction.

We may thus assume that GK/K is not normal in A/K . Again, we use the bar notation with respect to the mapping $A \mapsto A/K$. By Lemma 2.3, \bar{A} is a colour-preserving group of automorphisms of Γ/N . By induction, we have $\bar{G} = (\bar{F} \times \bar{H}) \rtimes \bar{D}$ and $\bar{A} = (\bar{T} \times \bar{J}) \rtimes \bar{D}$, where \bar{D} is a Sylow 2-subgroup of \bar{G} , $\text{PSL}(2, 7) \cong \bar{T} \trianglelefteq \bar{A}$, $\bar{T} \cap \bar{G} = \bar{F} \cong F_{21}$, $\bar{J} \cap \bar{G} = \bar{H} \trianglelefteq \bar{J} \trianglelefteq \bar{A}$ and \bar{H} is self-centralising in \bar{J} . Further, since $R \cap K = 1$ we may assume $\bar{R} = \bar{D}$. Note that $T/C_T(K) \leq \text{Aut}(K)$ is soluble since K is either cyclic or dihedral. As $T/K \cong \text{PSL}(2, 7)$, it follows that $T = KC_T(K)$ and hence $C_T(K)/Z(K) \cong \text{PSL}(2, 7)$. If $K \cong D_p$, then $Z(K) = 1$. Set $T_0 = C_T(K)$ in this case. Otherwise, $Z(K) = N = K \cong \mathbb{Z}_p$ and, since the Schur multiplier of $\text{PSL}(2, 7)$ has order 2, we have $C_T(K) = N \times T_0$ for some T_0 . In both cases, $T_0 \cong \text{PSL}(2, 7)$ and, since both T and K are normal in A , so is T_0 , which proves (i). Now, $TJ = T_0KJ = T_0J$, both T_0 and J are normal in A and $T_0 \cap J = 1$ hence $A = (T_0 \times J) \rtimes R$. Since $\bar{T}_0 = \bar{T}$ there is $F_0 \leq T_0$ such that $\bar{F}_0 = \bar{F}$. Since $F_0 \cap K = 1$ we have $F_0 \cong \bar{F}_0 \cong F_{21}$. Further, $F = F_0K = F_0 \times K$. Since $|GK : G| \leq 2$, we have that $F_0 \leq G$ and, since F_0 is maximal in T_0 , we have $T_0 \cap G = F_0$ which is (ii).

Note that $|H|$ is not divisible by 4, hence $H = H_0 \rtimes K_{[1]}$ for some characteristic subgroup H_0 of H . In particular, $\bar{H} = \bar{H}_0$. Now $GK = FHR = F_0H_0K_{[1]}R$ so $G = F_0H_0R = (F_0 \times H_0) \rtimes R$. Since H_0 is characteristic in H , it is normal in J . Recall that $K \cap G = N \leq H_0$, hence $K \cap F_0R = 1$. As $\bar{J} \cap \bar{F}_0\bar{R} = 1$, this implies that $J \cap F_0R = 1$. Since $H_0 \leq J$, we have $J \cap G = H_0(J \cap F_0R) = H_0$, which is (iii). Note that $H_0/N \cong \bar{H}$. Since \bar{H} contains its centraliser in \bar{J} , we have $C_J(H_0) \leq HK = H_0K_{[1]}$. As $N \leq H_0$ and N is self-centralising in K , we have $C_J(H_0) \leq H_0$, which is (iv).

This concludes the proof. \square

We now build on the previous result and give some information about the structure of the connection set.

Theorem 5.2. *Let G be a Sylow cyclic group whose order is not divisible by four, let $\Gamma = \text{Cay}(G, S)$ be a connected non-CCA graph and let $A = \text{Aut}_c(\Gamma)$. Using the notation of Theorem 5.1, write $A = (T \times J) \rtimes R$ and $G = (F \times H) \rtimes R$. Let r be the generator of R , let $Y = S \setminus (F \cup (H \rtimes R))$ and let*

$$\Gamma' = \text{Cay}(F \rtimes R, (F \cap S) \cup \{r\} \cup \{s^2 : s \in Y\}).$$

Then

1. Γ' is connected and non-CCA,

2. $Y \subseteq \{fz : f \in F, z \in Hr, |f| = 3, |z| = 2\}$, and
3. if $Y \neq \emptyset$, then $|R| = 2$, and T commutes with R .

Proof. Since Γ is non-CCA, G is not normal in A . This yields the conclusion of Theorem 5.1. As in the proof of that theorem, for $g \in G$, we write $[g]$ for the vertex of $\text{Cay}(G, S)$ corresponding to g and, for $X \subseteq G$, we write $[X]$ for $\{[x] : x \in X\}$.

Let P be a Sylow 2-subgroup of A containing R . Up to relabelling, we may assume that $A_{[1]} \leq P$ and thus $P = A_{[1]}R$. It follows that $[1]^{PH} = [H \rtimes R]$ is a block for A . As T is normal in A , its orbits are also blocks. One such block is $[1]^T = [F]$. As $[F] \cap [H \rtimes R] = [1]$, we find that the two block systems induced by $[F]$ and by $[H \rtimes R]$ are transverse.

The action of T on $[F]$ is equivalent to the action of $\text{PSL}(2, 7)$ by conjugation on its 21 Sylow 2-subgroups. In particular, if $f \in F$ and $T_{[1]} = T_{[f]}$, then $f = 1$. This observation, together with the previous paragraph, yields that the set of fixed points of $T_{[1]}$ is exactly $[H \rtimes R]$.

We first show (2) and (3). Let $s \in Y$. Note that $[Fs]$ is the orbit of T that contains s . Since $s \notin H \rtimes R$, $[s]$ is not fixed by $T_{[1]}$. Since T is colour-preserving, we have that $[s] \neq [s^{-1}] \in [Fs]$. It follows that $1 \neq s^2 \in F$. In particular, $|s^2| \in \{3, 7\}$. Since A is colour-preserving, the cycles coloured $\{s, s^{-1}\}$ form a block system for A . This means that $[\langle s^2 \rangle]$ is also a block for A , contained in the block $[F]$. Now, $\text{PSL}(2, 7)$ on its action on 21 points does not admit blocks of size 7, therefore $|s^2| = 3$. Since $s \notin F$ this implies $|s| = 6$. Notice also that $\{[1, s^3]\}$ is a block of A . Thus, $[s^3]$ is a fixed point of $T_{[1]}$, so $[s^3] \in [H \rtimes R]$. Since $|s^3| = 2$ but $|H|$ is odd, $s^3 \notin H$ hence $|R| = 2$ and $s^3 \in Hr$. Since H and F centralise each other and $s^3r \in H$ and $s^2 \in F$, it follows that s^2 commutes with r . Note that $[1]^P = [R]$ is a block for A . Now, $[\langle s^2 \rangle]$ is also a block for A , being a set of vertices of even distance contained in one of the monochromatic hexagons coloured $\{s, s^{-1}\}$. It follows that $[\langle s^2, r \rangle]$ is also a block for A , of size 6 and contained in the block $[1]^{TP} = [F \rtimes R]$. Note that $\text{PGL}(2, 7)$ does not have blocks of size 6 in its transitive action on 42 points. It follows that $T \rtimes R \not\cong \text{PGL}(2, 7)$, and hence T commutes with R . Writing $f = s^4$ and $z = s^3$ concludes the proof of (2) and (3).

Let $\pi : G \mapsto F \rtimes R$ be the natural projection and let $s \in Y$. By the previous paragraph, we have $s^{-1} = s^2s^3 = s^2hr$, where $s^2 \in F$ and $h \in H$. It follows that $\pi(s^{-1}) = s^2r$. As S is inverse-closed, we have $\pi(Y) = \{s^2r : s \in Y\}$. Since $\langle S \rangle = G$, we have $F \rtimes R = \langle \pi(S) \rangle \leq \langle F \cap S, r, s^2r : s \in Y \rangle = \langle F \cap S, r, s^2 : s \in Y \rangle$ and thus Γ' is connected. Note that $[1]^{T \rtimes R} = [F \rtimes R]$ hence $T \rtimes R \leq \text{Aut}_c(\Gamma')$. Since $F \rtimes R$ is not normal in $T \rtimes R$, Γ' is not CCA. \square

The following result is, in some sense, a converse to Corollary 5.2.

Proposition 5.3. *Let G be a Sylow cyclic group whose order is not divisible by four such that $G = (F \times H) \rtimes R$ where $F \cong F_{21}$, R is a Sylow 2-subgroup of G , and F and H are normal in G . Let r be the generator of R , let S be a generating set for G , let $Y = S \setminus (F \cup (H \rtimes R))$, let $S' = (F \cap S) \cup \{r\} \cup \{s^2 : s \in Y\}$, and let*

$$\Gamma' = \text{Cay}(F \rtimes R, S').$$

If

1. Γ' is connected and non-CCA,

2. $Y \subseteq \{fz : f \in F, z \in Hr, |f| = 3, |z| = 2\}$, and
3. if $Y \neq \emptyset$, then $|R| = 2$, and F commutes with R ,

then $\text{Cay}(G, S)$ is connected and non-CCA.

Proof. Since S generates G , $\text{Cay}(G, S)$ is connected. Since Γ' is a connected and non-CCA Cayley graph on $F \rtimes R$, it follows from Theorem 5.1 that there exists a group $T \rtimes R$ of colour-preserving automorphisms of Γ' , with $F \leq T$ and $T \cong \text{PSL}(2, 7)$.

This yields an action of T on $F \rtimes R$. We extend this action to the vertex-set of $\text{Cay}(G, S)$ in the following way: for $t \in T$ and $xh \in G$, with $x \in F \rtimes R$ and $h \in H$, let $(xh)^t = x^t h$.

Notice that if $x \in F \rtimes R$, then, since $r \in S'$ is an involution and $T \rtimes R$ is colour-preserving on Γ' , for any $t \in T$ we have $(rx)^t = rx^t$.

Note that $F \leq T \cap G < T$. Since T is simple, it follows that $T \cap G$ is not normal in T . We claim that T is a colour-preserving group of automorphisms of $\text{Cay}(G, S)$. By the previous comment, this will show that $\text{Cay}(G, S)$ is non-CCA.

Let $t \in T$, let $v \in G$ and write $v = xh$ with $x \in F \rtimes R$ and $h \in H$. We will show that, for all $s \in S$, we have $(sv)^t = s^{\pm 1}v^t$.

Suppose first that $s \in S'$. (This includes the case when $s \in F$.) Since T is colour-preserving on Γ' , we have $(sx)^t = s^{\pm 1}x^t$. Since $sx \in F \rtimes R$, we have $(sv)^t = (sxh)^t = (sx)^t h = s^{\pm 1}x^t h = s^{\pm 1}v^t$, as required.

Suppose next that $s \in H \rtimes R$. Write $s = h'r^i$ and $x = r^j f$, where $h' \in H$, $f \in F$ and $i, j \in \mathbb{Z}$. Let $h'' \in H$ be such that $r^{i+j}h'' = h'r^{i+j}$. Then

$$\begin{aligned} (sv)^t &= (h'r^{i+j}fh)^t = (r^{i+j}h''fh)^t = (r^{i+j}fh''h)^t = (r^{i+j}f)^t h''h = r^{i+j}f^t h''h \\ &= r^{i+j}h''f^t h = h'r^{i+j}f^t h = sr^j f^t h = s(r^j f)^t h = sx^t h = sv^t, \end{aligned}$$

as desired.

Finally, suppose $s \in Y$. We can write $s = fz$ where $f \in F$, $z \in Hr$, $|f| = 3$ and $|z| = 2$. By (3), we have $s^3 = z$, $s^2 = f^2$ and $|s| = 6$. Since $s^3 \in H \rtimes R$, the argument of the previous paragraph shows $(s^3v)^t = s^3v^t$. On the other hand, since $s^2 \in S'$, we have $(s^3v)^t = (s^2(sv))^t = s^{\pm 2}(sv)^t$. Combining these gives $(sv)^t = s^{3\pm 2}v^t = s^{\pm 1}v^t$, as desired. \square

We view Theorem 5.2 as a reduction of the CCA problem for groups of the kind appearing in its statement to the determination of non-CCA graphs on F_{21} and $\text{AGL}(1, 7)$. It therefore becomes of significant interest to understand the structure of such graphs.

Let x and y be elements of order 7 and 6 in $\text{AGL}(1, 7)$, respectively, and let $d = (y^3)^x$. Note that $\langle x, y^2 \rangle = F_{21}$. Let

$$S_{21} = \{y^{\pm 2}, (xy^2)^{\pm 1}\}, \quad S_{42,1} = \{y^{\pm 2}, d\} \quad \text{and} \quad S_{42,2} = \{y^{\pm 2}, (y^{\pm 2})^d, d\}.$$

Note that $\text{Cay}(F_{21}, S_{21})$ is isomorphic to the line graph of the Heawood graph (see Example 4.8), while $\text{Cay}(\text{AGL}(1, 7), S_{42,1})$ is isomorphic to the line graph of the subdivision of the Heawood graph (see Example 4.11).

Proposition 5.4.

1. The graph $\text{Cay}(F_{21}, S)$ is connected but not CCA if and only if S is conjugate in $\text{AGL}(1, 7)$ to S_{21} .

2. The graph $\text{Cay}(\text{AGL}(1, 7), S)$ is connected but not CCA if and only if S is conjugate in $\text{AGL}(1, 7)$ to one of $S_{42,1}$ or $S_{42,2}$.
3. The graph $\text{Cay}(\text{F}_{21} \times \mathbb{Z}_2, S)$ is connected but not CCA if and only if S is conjugate in $\text{AGL}(1, 7) \times \mathbb{Z}_2$ to some inverse-closed subset of

$$\{y^{\pm 2}, (xy^2)^{\pm 1}, y^{\pm 2}r, (xy^2)^{\pm 1}r, r\}$$

that generates $\text{F}_{21} \times \mathbb{Z}_2$, where $\mathbb{Z}_2 = \langle r \rangle$.

Proof. This was verified using MAGMA [1]. The proof of the first claim can also be found in [3, Proposition 2.5, Remark 2.6]. \square

Remark 5.5. It can be checked that Proposition 5.4(3) yields eleven generating sets for $\text{F}_{21} \times \mathbb{Z}_2$, up to conjugacy in $\text{AGL}(1, 7) \times \mathbb{Z}_2$.

References

- [1] W. Bosma, J. Cannon and C. Playoust, The Magma algebra system. I: The user language, *J. Symbolic Comput.* **24** (1997), 235–265, doi:10.1006/jsc.1996.0125.
- [2] D. P. Byrne, M. J. Donner and T. Q. Sibley, Groups of graphs of groups, *Beitr. Algebra Geom.* **54** (2013), 323–332, doi:10.1007/s13366-012-0093-7.
- [3] E. Dobson, A. Hujdurović, K. Kutnar and J. Morris, On color-preserving automorphisms of Cayley graphs of odd square-free order, *J. Algebraic Combin.* **45** (2017), 407–422, doi:10.1007/s10801-016-0711-9.
- [4] E. Dobson, A. Hujdurović, K. Kutnar and J. Morris, Vertex-transitive digraphs with extra automorphisms that preserve the natural arc-colouring, *Australas. J. Combin.* **67** (2017), 88–100, https://ajc.maths.uq.edu.au/pdf/67/ajc_v67_p088.pdf.
- [5] A. Hujdurović, K. Kutnar, D. W. Morris and J. Morris, On colour-preserving automorphisms of Cayley graphs, *Ars Math. Contemp.* **11** (2016), 189–213, <http://amc-journal.eu/index.php/amc/article/view/771>.
- [6] H. Kurzweil and B. Stellmacher, *The Theory of Finite Groups: An Introduction*, Universitext, Springer-Verlag, New York, 2004, doi:10.1007/b97433, translated from the 1998 German original.
- [7] C. Martínez-Pérez and W. Willems, The trivial intersection problem for characters of principal indecomposable modules, *Adv. Math.* **222** (2009), 1197–1219, doi:10.1016/j.aim.2009.05.018.
- [8] J. Morris, Automorphisms of circulants that respect partitions, *Contrib. Discrete Math.* **11** (2016), 1–6, <http://cdm.ucalgary.ca/cdm/index.php/cdm/article/view/376>.