

review article
received: 2004-11-14

UDC 597.5:591.134(262-18)

LENGTH-WEIGHT RELATIONSHIPS OF FISHES FROM TURKISH SEAS: A REVIEW

Hatice TORCU KOÇ & Zeliha AKA ERDOĞAN

University of Balikesir, Faculty of Science and Arts, Department of Biology, TR-10100 Balikesir, Turkey

Jakov DULČIĆ

Institute of Oceanography and Fisheries, HR-21000 Split, P.O. BOX 500, Croatia
E-mail: dulcic@izor.hr

ABSTRACT

The paper presents 360 length-weight relationships gathered from available literature pertaining to 90 fish species belonging to 40 families occurring throughout Turkish seas. The value of the slope b ranged from 0.409 for *Thunnus thynnus* to 4.343 for *Dentex macrophthalmus*. The mean value of b was 3.088 ($\pm SD = 0.898$) and did not differ significantly from 3 (t-test, $p < 0.05$). The median value of b was 3.062 and 50% of b values ranged between 2.941 and 3.190.

Key words: length-weight relationship, fishes, Turkish seas, review

RAPPORTO PESO/LUNGHEZZA DEI PESCI NEI MARI DELLA TURCHIA: UNA RASSEGNA

SINTESI

Sulla base dei dati messi a disposizione dall'attuale letteratura, l'articolo elenca 360 esempi di rapporto peso/lunghezza di pesci appartenenti a 90 specie e 40 famiglie che dimorano nei mari della Turchia. La curva (b) va dallo 0,409 della specie *Thunnus thynnus* al 4,343 di quella del *Dentex macrophthalmus*. Il valore medio di b è 3,088 ($\pm SD = 0,898$) e si discosta pochissimo dal 3 (t-test, $p < 0,05$). La mediana del coefficiente b è di 3,062, mentre il 50 % del valore di b oscilla tra 2,941 e 3,190.

Parole chiave: rapporto peso/lunghezza, pesci, mari della Turchia, rassegna

INTRODUCTION

Recording length and weight data is a standard procedure of fish sampling programs. Such data are essential for studies of fish population dynamics. Length-weight relationships have a number of uses including: (a) the estimation of standing-crop biomass when the length-frequency distribution is known; (b) estimation of biomass from length observations because of technical difficulties (e.g. bobbing motion of the boat) and the amount of time required to record weight in the field; (c) the calculation of condition indexes in the analysis of ontogenetic changes; and (d) comparing the life stories and morphological variations of fish populations from different regions (Cailliet et al., 1986; Safran, 1992; Petrakis & Stergiou, 1995; Goncalves et al., 1996; Martin-Smith, 1996; Morato et al., 2001; Stergiou & Motopoulos, 2001; Filiz & Bilge, 2004).

In this review, we gathered 336 length-weight relationships from the available literature for 70 fish species from Turkish seas.

MATERIAL AND METHODS

All length-weight relationships presented here are the product of field studies conducted during 1972–2004 in Turkish seas. For the majority of the original length-weight relationship ($W=aL^b$), length was expressed in cm and weight in g (356 relationships out of 360; 98.8%), whereas for 4 (1.2%) relationships, length and weight were expressed in mm and g.

RESULTS AND DISCUSSION

Overall, 360 length-weight relationships, gathered from the literature, represent 90 species and 40 families occurring in Turkish marine waters (Tab. 1). Overall, 50 out of the 360 length-weight relationships refer to *Merluccius merluccius* (14.9%), 35 (10.4%) to *Mullus barbatus*, and 26 (7.7%) to *Pagellus erythrinus*, three of the most commercially important demersal species in Turkish seas. Regarding the number of studies, of which length-weight relationship of an individual family is recorded, Sparidae dominated the overall records (84, 25.0%), and followed by Mullidae (60, 17.9%), Merlucciidae (50, 14.9%), Engraulidae (17, 5.1%) and Mugilidae (14, 4.2%). Each of the remaining 28 families was represented by less than one relationship.

The value of the slope b in the plot of $\log W$ against $\log L$ ranged from 0.409 for *Thunnus thynnus* in Aegean and Mediterranean Sea to 4.343 for *Dentex mac-*

rophthalmus in Saros Bay. The mean value of b was 3.088 ($\pm SD = 0.8985$) and did not differ significantly from 3 (*t*-test, $p < 0.05$). The median value of b was 3.062 and 50 % of b values ranged between 2.941 and 3.19.

According to Stergiou & Moutopoulos (2001), a plot of a versus b for all known length-weight relationships of a species results in a linear relationship, and this relationship can be used to identify outliers. We have applied this method to all species with more than 10 length-weight relationships. This led to the detection of outliers, where the respective point deviated more than two standard deviations from the regression line. These length-weight relationships were marked as "questionable" in Table 1.

The length-weight relationship in fishes is affected by a number of factors, including season, habitat, gonad maturity, sex, diet and stomach fullness, health and preservation techniques (Dulčić & Kraljević, 1996), all of which were not accounted for in this review. As stressed by Bagenal & Tesch (1978) and Petrakis & Stergiou (1995), the use of relationships should be limited to the size range used to estimate the parameters. Extrapolation to juvenile or immature stages may be particularly unwise given the potential differences in growth patterns between adults and earlier life history stages (Bagenal & Tesch, 1978). The parameters of the length-weight relationship may vary significantly according to season, habitat and even on a daily basis (*ibid*). In fishes, size is generally more biologically relevant than age, mainly because several ecological and physiological factors are more size-dependent than age-dependent (Dulčić & Kraljević, 1996). Consequently, variability in size has important implications for diverse aspects of fisheries science and population dynamics. Length-weight relationships have several applications, namely on fish biology, physiology, ecology and fisheries assessment. In biological studies, length-weight relationships enable seasonal variations in fish growth to be followed and the calculation of condition indexes. In fisheries studies, length-weight relationships have many different uses, including the estimation of weight from length, the estimation of weight-at-age and the conversion of growth-in-length equations to growth-in-weight. The establishment of length-weight relationships is also fundamental for the calculation of production and biomass of a fish population, being very useful for biomass estimations based on visual census of fish populations.

This review of length-weight parameters reported here will be of considerable use in ongoing studies of the catch in Turkish commercial fisheries.

Tab. 1: Parameters of the length-weight relationship $W = aL^b$ of fish species from Turkish marine waters (length (cm), weight (g)).

Legend: Sex (M, male; F, female; C, combined; U, unidentified); Year = year of sampling; S = sampling season (AUT, Autumn; WI, Winter; SP, Spring; SU, Summer; C, All seasons); L = length (TL, Total length; FL, Fork length; SL, Standard length); a = intercept (from the literature); b = slope (of the relationship $W=aL^b$); r^2 = coefficient of correlation; N = sampling size; min and max = minimum and maximum sample lengths (cm); SD = standard deviation.

Species are listed in alphabetical order.

Tab. 1: Parametri razmerja $W = aL^b$ med dolžino in težo ribjih vrst, živečih v turških morjih (dolžina (cm), teža (g)).

Legenda: Sex = spol (M, samec; F, samica; C, oba skupaj; U, neidentificiran); Year = leto vzorčenja; S = sezona vzorčenja (AUT, jesen; WI, zima; SP, pomlad; SU, poletje; C, vse sezone); L = dolžina (TL, celotna dolžina; FL, dolžina razcepa repa; SL, standardna dolžina); a = presečišče (na podlagi virov); b = naklon (iz razmerja $W=aL^b$); r^2 = korelačni koeficient; N = število primerkov; min in max = najmanjše in največje vzorčevalne velikosti (cm); SD = standardni odklon.

Vrste so navedene v abecednem redu.

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Alosa</i> sp.	Black Sea	C	1988–94	C	TL	0.0081	3.103	0.921*	65	—	—	—	—	Erkoyuncu et al. (1994)
<i>Arnoglossus laterna</i>	İzmir Bay	C	1996–97	C	TL	0.00932	2.897	0.940	643	12.1	6.0	15.3	—	Mater & Bayhan (2000)
<i>Argentína sphyraena</i>	Sigacik trawl area	C	2003	C	TL	0.0062	2.93	0.93	238	—	7.5	20.7	—	Filiz & Bilge (2004)
<i>Atherina hepsetus</i>	Eceabat Dardanelles	C	1997–99	C	FL	0.01287	2.890	0.999	564	—	5.8	13.0	—	Altun (2000)
<i>Belone belone</i>	Black Sea	C	1988–94	C	TL	0.0005	3.203	0.940*	65	—	—	—	—	Erkoyuncu et al. (1994)
<i>Belone belone euxini</i>	Black Sea	C	1995–96	C	TL	0.00055	3.177	0.940*	643	37.55	31.2	52.2	0.17	Samsun (1995)
<i>Capros aper</i>	Yeşilova Bay	F	1992	SU	TL	0.0000294	2.912	0.883*	170	86.35	—	—	0.7	Kaya & Ozaydin (1996)
<i>Capros aper</i>	Yeşilova Bay	M	1992	SU	TL	0.0000204	2.915	0.883*	165	84.72	—	—	0.6	Kaya & Ozaydin (1996)
<i>Capros aper</i>	Sigacik trawl area	C	2003	WI, SU	TL	0.0232	2.83	0.98	455	—	2.9	10.1	—	Filiz & Bilge (2004)
<i>Cepola rubescens</i>	İzmir Bay	F	1996–97	C	TL	0.32388	1.269	0.792	131	34.43	11.5	45.6	6.76	Kaya et al. (2001)
<i>Caelorinchus caelorinchus</i>	Sigacik trawl area	C	2003	C	TL	0.0065	2.74	0.78	208	—	4.0	21.6	—	Filiz & Bilge (2004)
<i>Cepola rubescens</i>	İzmir Bay	M	1996–97	C	TL	0.2154	1.384	0.810	144	29.17	19.8	47.1	6.26	Kaya et al. (2001)
<i>Chelidonichthyes lucerna</i>	İskenderun Bay	F	1999–00	C	TL	0.0095	2.990	0.960*	199	—	8.0	30.3	—	Ismen et al. (2004)
<i>Chelidonichthyes lucerna</i>	İskenderun Bay	M	1999–00	C	TL	0.0089	3.010	0.980*	143	—	8.3	21.2	—	Ismen et al. (2004)
<i>Chelon labrosus</i>	Güllük Lagoon	C	1993–94	C	TL	0.0075539	3.067	0.962*	45	26.4	20.0	35.0	3.5	Hossucu (2001)
<i>Chlorophthalmus agassizi</i>	Sigacik trawl area	C	2003	C	TL	0.0027	3.37	0.98	378	—	7.7	17.5	—	Filiz & Bilge (2004)

Species	Area	Sex	Year	S	L	a	b	r ²	N	mean	min	max	SD	Source
<i>Chimaera monstrosa</i>	Sigacik trawl area	C	2003	C	TL	0.0028	2.82	0.98	17	—	13.1	75.3	—	Filiz & Bilge (2004)
<i>Conger conger</i>	Sigacik trawl area	C	2003	C	TL	0.0005	3.24	0.96	22	—	32.2	65.4	—	Filiz & Bilge (2004)
<i>Dasyatis pastinaca</i>	İskenderun Bay	F	1999–00	C	TL	0.00091	3.440	0.940	110	—	20.5	88.0	—	Ismen (2003)
<i>Dasyatis pastinaca</i>	İskenderun Bay	M	1999–00	C	TL	0.00237	3.170	0.950	146	—	20.0	73.0	—	Ismen (2003)
<i>Dasyatis pastinaca</i>	İskenderun Bay	C	1999–00	C	TL	0.00144	3.310	0.940	256	—	20.0	88.0	—	Ismen (2003)
<i>Dasyatis pastinaca</i>	Sigacik trawl area	C	2003	WI, SU	TL	0.0149	2.81	0.85	29	—	37.3	74.2	—	Filiz & Bilge (2004)
<i>Dentex dentex</i>	İskenderun Bay	C	2000	AUT	TL	0.0861	2.495	0.921	16	42.41	31.40	51.0	6.35	Can et al. (2000)
<i>Dentex gibbosus</i>	İskenderun Bay	C	2000	AUT	TL	0.0341	2.714	0.857	34	27.29	17.68	47.3	6.83	Can et al. (2000)
<i>Dentex macrophthalmus</i>	Saros Bay	C	1991	SU	TL	0.01760	3.071	0.864*	20	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	İzmir Bay	C	1991	SU	TL	0.02248	2.966	0.889*	20	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	Güllük Bay	C	1991	SU	TL	0.00782	3.292	0.974*	20	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	Gökova Bay	C	1991	SU	TL	0.02404	2.932	0.874*	60	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	Saros Bay	C	1992	WI	TL	0.01974	3.027	0.988*	12	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	Saros Bay	C	1992	SP	TL	0.00002	4.343	0.980*	58	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	Gökova Bay	C	1992	SP	TL	0.01280	3.107	0.970*	40	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	Gökova Bay	C	1993	SP	TL	0.01731	3.084	0.986*	46	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	İzmir Bay	C	1993	AUT	TL	0.02968	2.859	0.975*	9	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	Güllük Bay	C	1993	AUT	TL	0.02946	2.812	0.974*	10	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	Gökova Bay	C	1993	AUT	TL	0.02117	2.977	0.978*	82	—	—	—	—	Benli et al. (2000)
<i>Dentex macrophthalmus</i>	Aegean Sea	F	1991–93	C	FL	0.01308	3.171	0.955*	285	14.10	8.2	22.0	0.18	Ozaydin (1997)
<i>Dentex macrophthalmus</i>	Aegean Sea	M	1991–93	C	FL	0.00921	3.282	0.941*	203	14.65	8.2	22.0	0.37	Ozaydin (1997)
<i>Dentex macrophthalmus</i>	Aegean Sea	C	1991–93	C	FL	0.01191	3.198	0.941*	507	14.23	7.8	22.0	0.20	Ozaydin (1997)
<i>Dentex macrophthalmus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000099	3.143	0.962*	87	—	10.7	19.5	—	Anonymous (1993)
<i>Diplodus annularis</i>	Edremit Bay	F	1997–98	C	FL	0.0367	2.797	0.865*	322	10.33	7.3	13.8	0.57	Koc et al. (2002)
<i>Diplodus annularis</i>	Edremit Bay	M	1997–98	C	FL	0.0455	2.696	0.846*	330	10.18	8.0	14.0	0.55	Koc et al. (2002)
<i>Diplodus annularis</i>	İzmir Bay	C	1997–99	C	FL	0.014	3.190	—	160	—	7.8	15.6	—	Kıncıgil & Akyol (2001)
<i>Diplodus annularis</i>	İzmir Bay	F	1997–99	C	FL	0.018	3.090	—	72	—	—	—	—	Kıncıgil & Akyol (2001)
<i>Diplodus annularis</i>	İzmir Bay	M	1997–99	C	FL	0.012	3.260	—	68	—	—	—	—	Kıncıgil & Akyol (2001)

Species	Area	Sex	Year	S	L	a	b	r ²	N	mean	min	max	SD	Source
<i>Diplodus annularis</i>	Çandarlı Bay	C	1992	WI	TL	0.04792	2.643	0.886*	24	—	—	—	—	Benli et al. (2000)
<i>Diplodus annularis</i>	Güllük Bay	C	1992	WI	TL	0.02056	3.016	0.957*	20	—	—	—	—	Benli et al. (2000)
<i>Diplodus annularis</i>	Güllük Bay	C	1993	SP	TL	0.02791	2.926	0.945*	20	—	—	—	—	Benli et al. (2000)
<i>Diplodus annularis</i>	Çandarlı Bay	C	1993	AUT	TL	0.01509	3.187	0.927*	36	—	—	—	—	Benli et al. (2000)
<i>Diplodus annularis</i>	İzmir Bay	C	1993	AUT	TL	0.01449	3.196	0.931*	32	—	—	—	—	Benli et al. (2000)
<i>Diplodus annularis</i>	Gökova Bay	C	1993	AUT	TL	0.09692	2.473	0.893*	18	—	—	—	—	Benli et al. (2000)
<i>Diplodus annularis</i>	Gülbahçe Bay	C	1996–1997	C	FL	0.021	3.023	0.961	205	6.5	15.0	—	Tosunoglu et al. (1997)	
<i>Diplodus annularis</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000050	3.299	0.929*	214	—	8.7	17.3	—	Anonymus (1993)
<i>Diplodus annularis</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0000248	2.959	0.895*	381	—	8.4	15.4	—	Anonymus (1993)
<i>Diplodus annularis</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000327	2.905	0.918*	193	—	8.2	19.3	—	Anonymus (1993)
<i>Diplodus annularis</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000102	3.150	0.976*	104	—	9.1	16.6	—	Anonymus (1993)
<i>Diplodus sargus</i>	İskenderun Bay	C	2000	AUT	TL	0.0342	2.807	0.850	33	19.29	14.90	26.7	2.94	Can et al. (2000)
<i>Diplodus vulgaris</i>	Saros Bay	C	1992	WI	TL	0.02840	2.929	0.943*	20	—	—	—	—	Benli et al. (2000)
<i>Diplodus vulgaris</i>	İzmir Bay	C	1993	AUT	TL	0.02413	2.983	0.952*	35	—	—	—	—	Benli et al. (2000)
<i>Diplodus vulgaris</i>	İskenderun Bay	C	2000	AUT	TL	0.0131	3.124	0.933	105	16.66	13.20	27.0	2.36	Can et al. (2000)
<i>Diplodus vulgaris</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000334	2.929	0.943*	20	—	12.1	17.5	—	Anonymus (1993)
<i>Diplodus vulgaris</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.000365	2.908	0.992*	20	—	10.5	18.9	—	Anonymus (1993)
<i>Diplodus vulgaris</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0000680	2.795	0.984*	11	—	8.8	16.7	—	Anonymus (1993)
<i>Diplodus vulgaris</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000016	3.538	0.960*	18	—	13.0	18.0	—	Anonymus (1993)
<i>Dipturus oxyrinchus</i>	Sigacik trawl area	C	2003	WI, SU	TL	0.0007	3.40	0.99	8	—	17.9	62.2	—	Filiz & Bilge (2004)
<i>Engraulis encrasiculus</i>	Eastern Black Sea	C	2001–02	WI	SL	0.032	2.507	0.895	50	10.48	9.45	11.7	0.066	Aka (2003)
<i>Engraulis encrasiculus</i>	Mid Black Sea	C	2001–02	WI	SL	0.0214	2.673	0.913	50	10.04	8.75	11.35	0.088	Aka (2003)
<i>Engraulis encrasiculus</i>	Western Black Sea	C	2001–02	WI	SL	0.0111	2.979	0.865	50	10.28	9.35	11.05	0.061	Aka (2003)

Species	Area	Sex	Year	S	L	a	b	r ²	N	mean	min	max	SD	Source
<i>Engraulis encrasiculus</i>	Marmara Sea	C	2001–02	WI	SL	0.0094	2.999	0.942	50	11.34	10.5	12.1	0.060	Aka (2003)
<i>Engraulis encrasiculus</i>	İzmir Bay	C	2001–02	WI	SL	0.0102	2.937	0.852	50	10.34	9.5	11.2	0.049	Aka (2003)
<i>Engraulis encrasiculus</i>	Edremit Bay	C	2001–02	WI	SL	0.009	2.991	0.866	50	10.14	9.15	11.45	0.062	Aka (2003)
<i>Engraulis encrasiculus</i>	Eastern Black Sea	F	1997–98	—	TL	0.0054	3.040	0.891	907	11.71	8.8	13.8	3.42	Gözler & Ciloglu (1998)
<i>Engraulis encrasiculus</i>	Eastern Black Sea	M	1997–98	—	TL	0.0049	3.071	0.640	339	11.68	8.8	13.8	1.74	Gözler & Ciloglu (1998)
<i>Engraulis encrasiculus</i>	Eastern Black Sea	C	1997–98	—	TL	0.0057	3.015	0.855	1664	11.22	7.0	13.8	4.43	Gözler & Ciloglu (1998)
<i>Engraulis encrasiculus</i>	Eastern Black Sea	C	1996–97	—	TL	0.00569	3.117	—	—	—	6.2	13.5	—	Kayalı (1998)
<i>Engraulis encrasiculus</i>	Eastern Black sea	C	1993–94	—	TL	0.0051	3.048	—	—	10.43	—	—	—	Düzungünes et al. (1995)
<i>Engraulis encrasiculus</i>	Black Sea	C	1988–94	C	TL	0.0053	3.038	0.941*	43	—	—	—	—	Erkoyuncu et al. (1994)
<i>Engraulis encrasiculus</i>	Black Sea	C	—	—	—	0.0047	3.100	—	—	—	—	—	—	Ozdamar et al. (1991a)
<i>Engraulis encrasiculus</i>	Black Sea	C	—	—	—	0.023	3.412	—	—	—	—	—	—	Ozdamar et al. (1991b)
<i>Engraulis encrasiculus</i>	Black Sea	C	—	—	—	0.0025	3.388	—	—	—	—	—	—	Karacan & Düzungünes (1990)
<i>Engraulis encrasiculus</i>	Black Sea	C	1987–88	—	TL	-2.5941	3.386	—	956	9.34	6.66	15.25	—	Düzungünes & Karacan (1989)
<i>Engraulis encrasiculus</i>	Black Sea	C	1988–89	C	TL	0.00643	2.974	—	1172	—	75	130	—	Unsal (1989)**
<i>Epinephelus costae</i>	İskenderun Bay	C	2000	AUT	TL	0.0885	2.391	0.930	53	29.76	14.20	55.4	8.78	Can et al. (2000)
<i>Gadus euxinus</i>	Eastern Black Sea	C	—	—	—	0.0272	2.573	—	—	—	—	—	—	Düzungünes & Karacan (1990)
<i>Gadus merlangus euxinus</i>	Black Sea	C	1995–96	C	TL	0.0039	3.238	0.931	1302	14.53	9.0	24.0	0.07	Samsun & Erkoyuncu (1998)
<i>Gadus merlangus euxinus</i>	Black Sea	C	1988–94	C	TL	0.0034	3.299	0.922*	54	—	—	—	—	Erkoyuncu et al. (1994)
<i>Gadus merlangus euxinus</i>	Mid Black Sea	C	—	C	—	0.0043	3.195	—	—	—	—	—	—	Samsun et al. (1993)
<i>Gadiculus argenteus argenteus</i>	Sigacik trawl area	C	2003	C	TL	0.0056	3.24	0.89	110	—	6.4	10.5	—	Filiz & Bilge (2004)
<i>Gymnura alatvela</i>	Sigacik trawl area	C	2003	C	TL	0.0268	2.96	0.98	9	—	37.5	72.0	—	Filiz & Bilge (2004)
<i>Helicolenus dactylopterus</i>	Sigacik trawl area	C	2003	C	TL	0.0079	3.28	0.92	178	—	5.5	13.5	—	Filiz & Bilge (2004)
<i>Hoplostethus mediterraneus</i>	Sigacik trawl area	C	2003	C	TL	0.0149	2.95	0.98	137	—	8.0	18.0	—	Filiz & Bilge (2004)

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Lesueurigobius friessi</i>	Sigacik trawl area	C	2003	C	TL	0.0392	2.13	0.72	17	—	6.2	8.1	—	Filiz & Bilge (2004)
<i>Lepidopus caudatus</i>	Sigacik trawl area	C	2003	C	TL	0.0004	3.11	0.99	40	—	21.9	81.5	—	Filiz & Bilge (2004)
<i>Liza ramada</i>	Güllük Lagoon	C	1993–94	C	TL	0.0066894	3.052	0.946*	86	23.5	17.0	33.8	2.8	Hossucu (2001)
<i>Liza saliens</i>	Güllük Lagoon	C	1993–94	C	TL	0.008859	2.997	0.965*	38	25.3	18.3	39.0	3.7	Hossucu (2001)
<i>Liza saliens</i>	İzmir Bay	U	1997–98	C	FL	0.0082	3.090	0.990	15	—	8.4	19.6	3.29	Akyol (1999)
<i>Liza saliens</i>	İzmir Bay	F	1997–98	C	FL	0.0088	3.080	0.953	294	—	13.4	33.1	—	Akyol (1999)
<i>Liza saliens</i>	İzmir Bay	M	1997–98	C	FL	0.0079	3.120	0.986	121	—	13.2	28.7	—	Akyol (1999)
<i>Liza saliens</i>	İzmir Bay	C	1997–98	C	FL	0.0078	3.120	0.985	430	—	8.4	33.1	—	Akyol (1999)
<i>Macroramphosus scolopax</i>	Sigacik trawl area	C	2003	C	TL	0.0079	3.28	0.92	43	—	7.1	11.4	—	Filiz & Bilge (2004)
<i>Merlangius merlangus euxinus</i>	Black Sea	F	1990–93	C	—	0.0040	3.250	0.980*	4176	—	—	—	—	İşmen (2002)
<i>Merlangius merlangus euxinus</i>	Black Sea	M	1990–93	C	—	0.0044	3.220	0.980*	3181	—	—	—	—	İşmen (2002)
<i>Merlangius merlangus euxinus</i>	Eastern Black Sea	F	1997	C	—	0.004856	3.151	—	1349	—	—	—	—	Sahin & Akbulut (1997)
<i>Merlangius merlangus euxinus</i>	Eastern Black Sea	M	1997	C	—	0.005450	3.110	—	864	—	—	—	—	Sahin & Akbulut (1997)
<i>Merlangius merlangus euxinus</i>	Eastern Black Sea	C	1990	C	—	0.0070	3.007	—	—	—	—	—	—	Uysal (1990)
<i>Merluccius merluccius</i>	Saros Bay	C	1991	SU	TL	0.00669	3.013	0.736*	275	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1991	SU	TL	0.00933	2.617	0.64*	78	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1991	SU	TL	0.01038	2.862	0.674*	53	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1991	SU	TL	0.00732	2.977	0.681*	203	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Saros Bay	C	1992	WI	TL	0.00382	3.221	0.960*	74	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1992	WI	TL	0.00507	3.097	0.986*	156	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1992	WI	TL	0.00383	3.188	0.995*	35	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1992	WI	TL	0.00573	3.025	0.997*	21	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Saros Bay	C	1992	SP	TL	0.00297	3.164	0.960*	170	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1992	SP	TL	0.00572	3.029	0.941*	73	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1992	SP	TL	0.00670	3.101	0.980*	9	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1992	SP	TL	0.00570	3.050	0.960*	68	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Saros Bay	C	1993	SP	TL	0.00297	3.075	0.970*	60	—	—	—	—	Benli et al. (2000)

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1993	SP	TL	0.00561	3.069	0.991*	35	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1993	SP	TL	0.00665	3.037	0.903*	37	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Güllük Bay	C	1993	SP	TL	0.00291	3.257	0.943*	17	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1993	SP	TL	0.00618	3.049	0.983*	51	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Saros Bay	C	1993	AUT	TL	0.00642	3.028	0.997*	76	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1993	AUT	TL	0.00312	3.266	0.989*	68	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1993	AUT	TL	0.00432	3.175	0.986*	58	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Güllük Bay	C	1993	AUT	TL	0.00731	3.017	0.993*	20	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1993	AUT	TL	0.00396	3.173	0.977*	26	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Saros Bay	C	1991	SU	TL	0.00669	3.013	0.736*	275	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1991	SU	TL	0.00933	2.617	0.640*	78	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1991	SU	TL	0.01038	2.862	0.674*	53	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1991	SU	TL	0.00732	2.977	0.681*	203	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Saros Bay	C	1992	WI	TL	0.00382	3.221	0.960*	74	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1992	WI	TL	0.00507	3.097	0.986*	156	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1992	WI	TL	0.00383	3.188	0.995*	35	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1992	WI	TL	0.00573	3.025	0.997*	21	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Saros Bay	C	1992	SP	TL	0.00297	3.167	0.960*	170	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1992	SP	TL	0.00572	3.029	0.941*	73	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1992	SP	TL	0.00670	3.014	0.980*	9	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1992	SP	TL	0.00570	3.050	0.960*	68	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Saros Bay	C	1993	SP	TL	0.00572	3.075	0.970*	60	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1993	SP	TL	0.00561	3.069	0.991*	35	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1993	SP	TL	0.00665	3.037	0.903*	37	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Güllük Bay	C	1993	SP	TL	0.00291	3.257	0.943*	17	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1993	SP	TL	0.00618	3.049	0.983*	51	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Saros Bay	C	1993	AUT	TL	0.00642	3.028	0.977*	76	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Çandarlı Bay	C	1993	AUT	TL	0.00312	3.266	0.989*	68	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	İzmir Bay	C	1993	AUT	TL	0.00432	3.175	0.983*	58	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Güllük Bay	C	1993	AUT	TL	0.00731	3.017	0.993*	20	—	—	—	—	Benli et al. (2000)
<i>Merluccius merluccius</i>	Gökova Bay	C	1993	AUT	TL	0.00396	3.173	0.977*	26	—	—	—	—	Benli et al. (2000)

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Merluccius merluccius</i>	Aegean Sea	C	1994–95	C	TL	0.0045	3.194	0.974*	336	—	13.6	43.5	—	Uckun et al. (2000)
<i>Merluccius merluccius</i>	Edremit Bay	C	1996–97	—	TL	0.0095	2.928	0.960*	165	—	15.8	37.2	—	Torcu et al. (1997)
<i>Merluccius merluccius</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0000048	3.065	0.984*	1164	—	10.5	69.5	—	Anonymous (1993)
<i>Merluccius merluccius</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000037	3.117	0.990*	1114	—	7.8	73.5	—	Anonymous (1993)
<i>Merluccius merluccius</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000043	3.090	0.988*	614	—	14.2	79.0	—	Anonymous (1993)
<i>Merluccius merluccius</i>	İzmir Bay	C	1989	AUT	FL	0.00246	2.960	—	74	26.10	20.50	36.14	—	Kara & Gurbet (1990)
<i>Mugil cephalus</i>	Güllük Lagoon	C	1993–94	C	TL	0.008198	3.048	0.976*	132	31.0	17.6	46.2	6.6	Hossucu (2001)
<i>Mugil cephalus</i>	İzmir Bay	U	1997–98	C	FL	0.0149	2.950	0.998	29	—	10.2	45.3	—	Akyol (1999)
<i>Mugil cephalus</i>	İzmir Bay	F	1997–98	C	FL	0.0075	3.160	0.989	98	—	25.7	54.5	—	Akyol (1999)
<i>Mugil cephalus</i>	İzmir Bay	M	1997–98	C	FL	0.0109	3.040	0.965	77	—	28.2	55.5	—	Akyol (1999)
<i>Mugil cephalus</i>	İzmir Bay	C	1997–98	C	FL	0.0101	3.070	0.991	204	—	10.2	55.5	—	Akyol (1999)
<i>Mugil so-iuy</i>	Eastern Black Sea	C	1995	C	TL	0.010	2.980	0.970*	174	—	22.5	68.9	—	Başçınar & Okumus (1997)
<i>Mugil</i> sp.	Black Sea	C	1988–94	C	TL	0.0068	3.077	0.980*	75	—	—	—	—	Erkoyuncu et al. (1994)
<i>Mullus barbatus</i>	İzmir Bay	C	1997	WI	FL	0.0071	3.290	—	221	—	10.3	19.1	—	Kıncıgil et al. (2001)
<i>Mullus barbatus</i>	Saroz Bay	C	1991	SU	TL	0.01493	2.968	0.965*	222	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Çandarlı Bay	C	1991	SU	TL	0.00775	3.234	0.918*	40	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	İzmir Bay	C	1991	SU	TL	0.01540	2.988	0.905*	118	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Güllük Bay	C	1991	SU	TL	0.02774	2.768	0.704*	122	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Gökova Bay	C	1991	SU	TL	0.01578	2.998	0.857*	60	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Saroz Bay	C	1992	WI	TL	0.00942	3.188	0.977*	80	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Çandarlı Bay	C	1992	WI	TL	0.001261	3.067	0.973*	166	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	İzmir Bay	C	1992	WI	TL	0.00871	3.204	0.980*	39	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Güllük Bay	C	1992	WI	TL	0.01442	3.051	0.980*	53	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Gökova Bay	C	1992	WI	TL	0.00824	3.219	0.967*	40	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Saroz Bay	C	1992	SP	TL	0.00321	3.309	0.922*	170	—	—	—	—	Benli et al. (2000)

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Mullus barbatus</i>	Çandarlı Bay	C	1992	SP	TL	0.00328	3.313	0.960*	100	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	İzmir Bay	C	1992	SP	TL	0.00346	3.305	0.980*	120	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Güllük Bay	C	1992	SP	TL	0.00110	3.051	0.941*	60	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Gökova Bay	C	1992	SP	TL	0.04830	2.801	0.884*	11	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Saroz Bay	C	1993	SP	TL	0.01421	3.029	0.990*	38	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	İzmir Bay	C	1993	SP	TL	0.01382	3.038	0.938*	91	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Güllük Bay	C	1993	SP	TL	0.01205	3.115	0.981*	40	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Gökova Bay	C	1993	SP	TL	0.00736	3.314	0.960*	68	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Saroz Bay	C	1993	AUT	TL	0.03087	2.738	0.899*	48	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Çandarlı Bay	C	1993	AUT	TL	0.02738	2.824	0.963*	40	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	İzmir Bay	C	1993	AUT	TL	0.01723	2.984	0.902*	115	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	Gökova Bay	C	1993	AUT	TL	0.00767	3.292	0.950*	99	—	—	—	—	Benli et al. (2000)
<i>Mullus barbatus</i>	İskenderun Bay	C	1992–93	C	FL	-1.77494	3.026	0.956*	348	—	7.5	15.12	—	Türeli & Erdem (1997)
<i>Mullus barbatus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000033	3.320	0.960*	533	—	8.5	22.4	—	Anonymus (1993)
<i>Mullus barbatus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000091	3.106	0.970*	636	—	9.0	20.5	—	Anonymus (1993)
<i>Mullus barbatus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.00000201	2.941	0.955*	1084	—	8.5	21.0	—	Anonymus (1993)
<i>Mullus barbatus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000078	3.143	0.968*	1090	—	8.8	22.3	—	Anonymus (1993)
<i>Mullus barbatus</i>	Mid Black sea	C	—	—	—	0.0054	3.209	—	—	—	—	—	—	Anonymus (1991a)
<i>Mullus barbatus</i>	İzmir Bay	C	1989	AUT	FL	0.00542	3.344	—	231	13.30	10.83	19.50	—	Kara & Gurbet (1990)
<i>Mullus barbatus</i>	Aliağa-Çandarlı Bay	C	1989	AUT	FL	0.13243	2.310	—	199	14.68	11.20	19.12	—	Kara & Gurbet (1990)
<i>Mullus barbatus</i>	Edremit Bay	C	1989	AUT	FL	0.011782	2.960	—	196	15.34	11.50	18.75	—	Kara & Gurbet (1990)
<i>Mullus barbatus</i>	Gülbahçe Bay (İzmir)	C	1973	C	FL	0.0165	2.923	0.960*	6054	—	7.6	22.0	—	Togulga (1977)
<i>Mullus barbatus</i>	Edremit Bay	C	1970	SU	FL	0.022	—	—	204	—	10.0	19.0	—	Kınikaslan (1972)
<i>Mullus barbatus ponticus</i>	Eastern Black Sea	F	1997	C	—	0.0052653	3.217	—	—	—	—	—	—	Sahin & Akbulut (1997)

Species	Area	Sex	Year	S	L	a	b	r ²	N	mean	min	max	SD	Source
<i>Mullus barbatus ponticus</i>	Eastern Black Sea	M	1997	C	—	0.0053736	3.220	—	—	—	—	—	—	Sahin & Akbulut (1997)
<i>Mullus barbatus ponticus</i>	Black Sea	C	1988–89	C	TL	0.006855	3.156	—	2116	12.0	6.9	25.3	0.004	Samsun & Erkoyuncu (1992)
<i>Mullus surmuletus</i>	Saros Bay	C	1991	SU	TL	0.0078	3.089	0.950*	57	—	—	—	—	Benli et al. (2000)
<i>Mullus surmuletus</i>	Gökova Bay	C	1991	SU	TL	0.02168	2.920	0.926*	20	—	—	—	—	Benli et al. (2000)
<i>Mullus surmuletus</i>	Saros Bay	C	1992	WI	TL	0.03224	2.819	0.830*	6	—	—	—	—	Benli et al. (2000)
<i>Mullus surmuletus</i>	Güllük Bay	C	1992	WI	TL	0.00984	3.242	0.913*	20	—	—	—	—	Benli et al. (2000)
<i>Mullus surmuletus</i>	Çandarlı Bay	C	1992	SP	TL	0.00510	3.221	0.810*	27	—	—	—	—	Benli et al. (2000)
<i>Mullus surmuletus</i>	İzmir Bay	C	1992	SP	TL	0.00580	3.194	0.980	28	—	—	—	—	Benli et al. (2000)
<i>Mullus surmuletus</i>	İzmir Bay	C	1993	AUT	TL	0.01177	3.131	0.958	37	—	—	—	—	Benli et al. (2000)
<i>Mullus surmuletus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000087	3.122	0.947*	170	—	11.5	32.1	—	Anonymus (1993)
<i>Mullus surmuletus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0000488	2.784	0.929*	131	—	114.0	205.0	—	Anonymus (1993)
<i>Mullus surmuletus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000103	3.104	0.990*	67	—	97.0	238.0	—	Anonymus (1993)
<i>Mullus surmuletus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000209	2.971	0.968*	30	—	100.0	253.0	—	Anonymus (1993)
<i>Mustelus mustelus</i>	Sigacik trawl area	C	2003	WI, SU	TL	0.0011	3.25	0.99	35	—	38.3	97.5	—	Filiz & Bilge (2004)
<i>Myliobatis aquila</i>	Sigacik trawl area	C	2003	WI, SU	TL	0.0008	3.34	0.93	14	—	47.5	76.5	—	Filiz & Bilge (2004)
<i>Oblada melanura</i>	İskenderun Bay	C	2000	AUT	TL	0.0322	2.697	0.964	22	20.45	15.60	27.0	3.32	Can et al. (2000)
<i>Pagellus sp.</i>	İzmir Bay	C	1989	AUT	FL	0.01592	3.008	—	111	14.59	8.66	21.43	—	Kara & Gurbet (1990)
<i>Pagellus sp.</i>	Edremit Bay	C	1989	AUT	FL	0.14723	2.289	—	74	16.75	13.71	22.0	—	Kara & Gurbet (1990)
<i>Pagellus acerna</i>	Aegean Sea	F	1991–93	C	FL	0.00527	3.447	0.850*	85	14.74	14.7	21.0	0.34	Ozaydin (1997)
<i>Pagellus acerna</i>	Aegean Sea	M	1991–93	C	FL	0.01087	3.172	0.927*	139	13.68	10.5	17.8	0.22	Ozaydin (1997)
<i>Pagellus acerna</i>	Aegean Sea	C	1991–93	C	FL	0.01420	3.075	0.962*	302	13.27	7.8	21.0	0.24	Ozaydin (1997)
<i>Pagellus acerna</i>	Gülbahçe Bay	C	1996–97	C	FL	0.082	3.288	0.935	107	—	9.8	14.8	—	Tosunoglu et al. (1997)
<i>Pagellus acerna</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000876	2.646	0.806*	40	—	8.0	15.0	—	Anonymus (1993)

Species	Area	Sex	Year	S	L	a	b	r ²	N	mean	min	max	SD	Source
<i>Pagellus bogaraveo</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000018	3.472	0.951*	73	—	9.0	20.0	—	Anonymus (1993)
<i>Pagellus bogaraveo</i>	Marmara, Aegean and	C	1991–93	SU	FL	0.0000083	3.150	0.978*	5	—	11.0	15.0	—	Anonymus (1993)
<i>Pagellus bogaraveo</i>	Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000709	2.709	0.792*	20	—	11.0	14.0	—	Anonymus (1993)
<i>Pagellus bogaraveo</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000061	3.214	0.968*	81	—	9.0	19.0	—	Anonymus (1993)
<i>Pagrus coeruleostictus</i>	İskenderun Bay	C	2000	AUT	TL	0.0671	2.521	0.908	311	18.52	12.50	38.8	2.93	Can et al. (2000)
<i>Pagellus erythrinus</i>	Saros Bay	C	1991	SU	TL	0.01299	2.972	0.832*	124	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Çandarlı Bay	C	1991	SU	TL	0.01190	3.156	0.959*	20	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	İzmir Bay	C	1991	SU	TL	0.04097	2.889	0.857*	60	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Güllük Bay	C	1991	SU	TL	0.01816	2.968	0.908*	40	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Gökova Bay	C	1991	SU	TL	0.0255	2.932	0.822*	40	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Saros Bay	C	1992	WI	TL	0.03184	2.773	0.896*	20	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Çandarlı	C	1992	WI	TL	0.01459	3.079	0.978*	29	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Gökova Bay	C	1992	WI	TL	0.01356	3.101	0.945*	20	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Saros Bay	C	1992	SP	TL	0.00109	3.537	0.941*	46	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Çandarlı Bay	C	1992	SP	TL	0.01619	3.011	0.960*	40	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	İzmir Bay	C	1992	SP	TL	0.06402	2.752	0.960*	56	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Güllük Bay	C	1992	SP	TL	0.02065	2.527	0.689*	42	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Gökova Bay	C	1992	SP	TL	0.04277	2.838	0.960*	7	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Saros Bay	C	1993	SP	TL	0.01249	3.140	0.988*	16	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	İzmir Bay	C	1993	SP	TL	0.01498	3.077	0.988*	57	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Güllük Bay	C	1993	SP	TL	0.06999	2.539	0.975*	17	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Gökova Bay	C	1993	SP	TL	0.01754	2.993	0.956*	17	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Saros Bay	C	1993	AUT	TL	0.00606	3.391	0.984*	17	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Çandarlı Bay	C	1993	AUT	TL	0.06999	2.539	0.975*	17	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	İzmir Bay	C	1993	AUT	TL	0.01699	3.001	0.944*	57	—	—	—	—	Benli et al. (2000)
<i>Pagellus erythrinus</i>	Aegean Sea	F	1991–93	C	FL	0.02370	2.907	0.978*	512	15.36	11.1	24.5	0.18	Ozaydin (1997)
<i>Pagellus erythrinus</i>	Aegean Sea	M	1991–93	C	FL	0.2722	2.865	0.956*	163	18.27	8.8	25.6	0.41	Ozaydin (1997)
<i>Pagellus erythrinus</i>	Aegean Sea	C	1991–93	C	FL	0.02423	2.985	0.972*	692	16.04	8.8	25.6	0.20	Ozaydin (1997)

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Pagellus erythrinus</i>	Gülbahçe Bay	C	1996–97	C	FL	0.018	3.012	0.976	204	—	8.0	24.0	—	Tosunoglu et al. (1997)
<i>Pagellus erythrinus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000287	2.917	0.966*	387	—	7.0	25.0	—	Anonymous (1993)
<i>Pagellus erythrinus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000122	3.080	0.966*	117	—	11.0	24.0	—	Anonymous (1993)
<i>Peristedion cataphractum</i>	Sigacil trawl area	C	2003	C	TL	0.0048	2.97	0.99	11	—	8.1	21.2	—	Filiz & Bilge (2004)
<i>Pleuronectes platessa</i>	Black Sea	C	1988–94	C	TL	0.0078	3.109	0.960*	48	—	—	—	—	Erkoyuncu et al. (1994)
<i>Phycis blennoides</i>	Sigacik trawl area	C	2003	C	TL	0.0017	3.55	0.89	12	—	12.3	15.0	—	Filiz & Bilge (2004)
<i>Pomadasys incisus</i>	İskenderun Bay	C	2000	AUT	TL	0.0465	2.604	0.910	106	16.12	13.40	21.2	1.75	Can et al. (2000)
<i>Pomatomus saltatrix</i>	Black Sea	C	1988–94	C	TL	0.0388	2.558	0.846*	19	—	—	—	—	Erkoyuncu et al. (1994)
<i>Pomatomus saltator</i>	İzmir Bay	C	1987	AUT —WI	TL	0.02017	2.955	—	400	—	15.0	40.5	—	Alpbaz & Kinacıgil (1988)
<i>Raja clavata</i>	Sinop	C	—	—	—	0.0026	3.200	0.990	54	—	27.3	83.2	—	Erdem et al. (2001)
<i>Raja clavata</i>	Sigacik trawl area	C	2003	WI, SU	TL	0.0016	3.30	0.94	37	—	20.5	99.0	—	Filiz & Bilge (2004)
<i>Raja miraletus</i>	Sigacik trawl area	C	2003	WI, SU	TL	0.0001	4.15	0.93	13	—	30.0	50.5	—	Filiz & Bilge (2004)
<i>Raja</i> sp.	Black Sea	C	1988–94	C	TL	0.0090	2.920	0.922*	40	—	—	—	—	Erkoyuncu et al. (1994)
<i>Sarda sarda</i>	Black Sea	C	1995–96	AUT —WI	TL	0.0058	3.176	0.980*	4104	34.42	20.0	40.0	2.21	Samsun (1997)
<i>Sarda sarda</i>	Black Sea	C	1988–94	C	TL	0.0297	2.679	0.865*	14	—	—	—	—	Erkoyuncu et al. (1994)
<i>Sardina pilchardus</i>	İzmir Bay	F	1996–97	C	TL	0.0053	3.275	0.872*	187	—	9.3	13.0	—	Karakayıs & Togulga (2000)
<i>Sardina pilchardus</i>	İzmir Bay	M	1996–97	C	TL	0.0081	3.108	0.781*	75	—	—	—	—	Karakayıs & Togulga (2000)
<i>Sardina pilchardus</i>	İzmir Bay	C	1996–97	C	TL	0.0062	3.214	0.855*	262	—	9.3	14.6	—	Karakayıs & Togulga (2000)
<i>Sardina pilchardus</i>	İzmir Bay	F	1997–98	C	FL	0.0042	3.382	0.850	216	—	9.6	14.7	—	Mater & Bayhan (1999)
<i>Sardina pilchardus</i>	İzmir Bay	M	1997–98	C	FL	0.0025	3.600	0.870	148	—	9.8	14.9	—	Mater & Bayhan (1999)
<i>Sardina pilchardus</i>	İzmir Bay	C	1997–98	C	FL	0.0045	3.359	0.830	364	—	9.6	14.9	—	Mater & Bayhan (1999)

Species	Area	Sex	Year	S	L	a	b	r ²	N	mean	min	max	SD	Source
<i>Saurida undosquamis</i>	İskenderun	F	1999–00	C	TL	0.088	3.190	0.980	368	—	9.0	35.0	—	Ismen (2002)
<i>Saurida undosquamis</i>	İskenderun	M	1999–00	C	TL	0.120	2.950	0.960	234	—	5.0	32.0	—	Ismen (2002)
<i>Saurida undosquamis</i>	Fethiye Bay	C	1993	C	FL	-1.3279	3.295	0.828*	430	—	17.3	30.8	—	Mater & Torcu (1997)
<i>Saurida undosquamis</i>	Mersin Bay	C	1993	C	FL	-0.9596	2.616	0.656*	100	—	13.5	32.2	—	Mater & Torcu (1997)
<i>Saurida undosquamis</i>	İskenderun Bay	C	1992–93	C	FL	-2.06171	3.022	0.998*	333	—	8.3	21.92	—	Türeli & Erdem (1997)
<i>Scomberomorus commerson</i>	Güllük and Gökova Bay	C	1994	C	TL	0.1567	2.223	0.885*	70	—	52.0	87.0	—	Buhan et al. (1997)
<i>Scophthalmus maeoticus</i>	Black Sea	C	—	C	—	0.0047	3.418	—	—	—	40.0	—	—	Erdem (2001)
<i>Scophthalmus maximus</i>	Eastern Black Sea	F	1990–91	C	TL	0.0099	3.140	0.980	86	—	196	630	—	Avsar (1999)**
<i>Scophthalmus maximus</i>	Eastern Black Sea	M	1990–91	C	TL	0.0112	3.110	0.980	63	—	181	595	—	Avsar (1999)**
<i>Scophthalmus maximus</i>	Eastern Black Sea	C	1990–91	C	TL	0.0085	3.180	0.980	194	—	181	630	—	Avsar (1999)**
<i>Scorpaena porcus</i>	Sinop	C	1996–97	—	TL	0.054	2.590	—	633	17.26	11.49	23.63	0.083	Koca (2002)
<i>Scorpaena</i> sp.	Black Sea	C	1988–94	C	TL	0.0180	3.080	0.980*	31	—	—	—	—	Erkoyuncu et al. (1994)
<i>Scyliorhinus canicula</i>	Northern Aegean Sea	F	1991–96	SP	TL	0.001	3.453	0.991	95	—	—	51.7	—	Cihangir et al. (1997)
<i>Scyliorhinus canicula</i>	Northern Aegean Sea	M	1991–96	SP	TL	0.005	2.004	0.989	121	—	—	54.6	—	Cihangir et al. (1997)
<i>Scyliorhinus canicula</i>	Northern Aegean Sea	C	1991–96	SU-AUT	TL	0.001	3.205	0.919	370	—	—	54.6	—	Cihangir et al. (1997)
<i>Scyliorhinus canicula</i>	Siagacik trwal area	C	2003	WI, SU	TL	0.0012	3.26	0.99	637	—	10.5	50.9	—	Filiz & Bilge (2004)
<i>Serranus cabrilla</i>	Edremit Bay	C	1997–98	C	FL	0.0311	2.670	0.880	595	14.43	8.6	22.3	1.573	Torcu et al. (2004)
<i>Serranus cabrilla</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000225	2.879	0.937*	100	—	94.0	200.0	—	Anonymus (1993)
<i>Serranus cabrilla</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0000081	3.059	0.929*	193	—	94.0	255.0	—	Anonymus (1993)
<i>Serranus cabrilla</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000066	3.121	0.966*	176	—	10.4	20.3	—	Anonymus (1993)
<i>Serranus cabrilla</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.000001	3.043	0.980*	75	—	11.6	17.5	—	Anonymus (1993)
<i>Serranus scriba</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0000252	2.878	0.719	20	—	8.0	24.0	—	Anonymus (1993)

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Siganus rivulatus</i>	Antalya Bay	F	1996–98	C	TL	0.0064	3.221	0.903*	292	15.12	7.0	21.5	0.32	Bilecenoglu & Kaya (2002)
<i>Siganus rivulatus</i>	Antalya Bay	M	1996–98	C	TL	0.007945	3.135	0.903*	229	16.09	7.1	20.6	0.34	Bilecenoglu & Kaya (2002)
<i>Siganus rivulatus</i>	Antalya Bay	C	1996–98	C	TL	0.007137	3.179	0.903*	521	—	7.0	21.5	—	Bilecenoglu & Kaya (2002)
<i>Solea solea</i>	İskenderun Bay	F	2000–01	C	TL	0.0093	3.077	0.960	530	—	20.5	28.2	—	Türkmen (2003)
<i>Solea solea</i>	İskenderun Bay	M	2000–01	C	TL	0.0117	2.998	0.973	553	—	8.8	25.0	—	Türkmen (2003)
<i>Solea</i> sp.	Black Sea	C	1988–94	C	TL	0.0019	3.580	0.941*	19	—	—	—	—	Erkoyuncu et al. (1994)
<i>Solea solea</i>	İzmir Bay	C	1989–90	C	TL	0.00514	3.135	0.815	335	26.2	11.0	34.5	0.6	Hossucu (1992)
<i>Solea solea</i>	İzmir Bay	C	1989	AUT	FL	0.00054	3.640	—	9	23.61	22.50	25.83	—	Kara & Gurbet (1990)
<i>Solea solea</i>	Aliağa-Çan-darlı Bay	C	1989	AUT	FL	0.00182	3.458	—	11	25.31	20.50	30.10	—	Kara & Gurbet (1990)
<i>Solea solea</i>	Edremit Bay	C	1989	AUT	FL	0.20137	1.951	—	14	26.21	21.50	31.00	—	Kara & Gurbet (1990)
<i>Solea vulgaris</i>	Yumurtalık	C	1992–93	C	TL	-2.52566	3.250	0.992*	603	—	12.1	34.5	—	Altun et al. (1997)
<i>Solea vulgaris</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000003	3.562	0.951*	8	—	23.8	34.2	—	Anonymus (1993)
<i>Solea vulgaris</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0000026	3.183	0.937*	66	—	13.9	27.4	—	Anonymus (1993)
<i>Solea vulgaris</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0001367	2.479	0.740*	23	—	19.2	33.5	—	Anonymus (1993)
<i>Solea vulgaris</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000098	2.958	0.960*	11	—	—	—	—	Anonymus (1993)
<i>Sparus aurata</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000025	2.941	0.960*	24	—	15.5	24.1	—	Anonymus (1993)
<i>Sparus aurata</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0000693	2.743	0.984*	9	—	15.3	22.2	—	Anonymus (1993)
<i>Sparus aurata</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000017	3.470	0.982*	31	—	14.2	22.3	—	Anonymus (1993)
<i>Sparus aurata</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000149	3.023	0.878*	14	—	16.0	19.6	—	Anonymus (1993)

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Sphyraena chrysotaenia</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000053	3.049	0.895*	6	—	21.0	25.2	—	Anonymus (1993)
<i>Sphyraena chrysotaenia</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000049	3.056	0.964*	22	—	14.0	25.6	—	Anonymus (1993)
<i>Sphyraena sphyraena</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0001309	2.428	0.925*	9	—	24.0	34.0	—	Anonymus (1993)
<i>Sphyraena sphyraena</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0002720	2.290	0.927*	13	—	22.0	35.0	—	Anonymus (1993)
<i>Spicara flexuosa</i>	İzmir Bay	F	1998–99	C	FL	0.1156	2.163	0.689*	240	—	9.20	14.90	—	Mater et al. (2001)
<i>Spicara flexuosa</i>	İzmir Bay	M	1998–99	C	FL	0.0452	2.566	0.740*	172	—	11.30	15.50	—	Mater et al. (2001)
<i>Spicara flexuosa</i>	İzmir Bay	C	1998–99	C	FL	0.0411	2.594	0.846*	412	—	9.20	15.50	—	Mater et al. (2001)
<i>Spicara smaris</i>	Eastern Black Sea	F	1989	C	TL	0.0510	3.229	0.958*	280	—	11.1	22.5	—	Sahin & Genc (1999)
<i>Spicara smaris</i>	Eastern Black Sea	M	1989	C	TL	0.0650	3.126	0.880*	176	—	11.3	22.0	—	Sahin & Genc (1999)
<i>Spicara smaris</i>	Eastern Black Sea	C	1991–92	C	TL	0.005	3.260	—	—	—	—	—	—	Ismen (1995)
<i>Spicara smaris</i>	Black Sea	C	1988–94	C	TL	0.0061	3.215	0.941*	25	—	—	—	—	Erkoyunucu et al. (1994)
<i>Spicara smaris</i>	Mid Black Sea	C	—	—	—	0.0051	3.217	—	—	—	—	—	—	Anonymus (1991c)
<i>Sprattus sprattus phallericus</i>	Eastern Black Sea	F	1991	SP–SU	TL	0.00214	3.456	0.999*	214	—	8.45	14.09	—	Sahin (1999)
<i>Sprattus sprattus phallericus</i>	Eastern Black Sea	M	1991	SP–SU	TL	0.00204	3.474	0.994*	158	—	8.27	12.5	—	Sahin (1999)
<i>Sprattus sprattus phallericus</i>	Eastern Black Sea	C	1991	SP–SU	TL	0.00211	3.461	0.997*	372	—	8.35	13.20	—	Sahin (1999)
<i>Squalus acanthias</i>	Black Sea	F	1969–73	—	TL	0.0041	3.004	0.996	1840	85	30.0	140.0	—	Kutaygil & Bilecik (1998)
<i>Squalus acanthias</i>	Black Sea	M	1969–73	C	TL	0.0053	2.929	0.999	1780	75	30.0	120.0	—	Kutaygil & Bilecik (1998)
<i>Squalus acanthias</i>	Eastern Black Sea	C	1994–95	C	TL	0.0013	3.254	0.81*	159	—	35.5	141.5	—	Karacam et al. (1996)
<i>Squalus acanthias</i>	Sigacik trawl area	C	2003	WI, SU	TL	0.0031	3.11	0.98	32	—	27.0	70.5	—	Filiz & Bigle (2004)
<i>Syphodus tinca</i>	İskenderun Bay	C	2000	AUT	TL	0.0021	3.675	0.997	10	14.41	12.10	17.2	1.75	Can et al. (2000)
<i>Thunnus thynnus</i>	Aegean and Mediterranean Sea	C	1996–98	C	TL	0.0000181	2.988	0.994*	3577	118.40	59.0	275.0	37.0	Karakulak & Oray (2001)
<i>Thunnus thynnus</i>	Aegean and Mediterranean Sea	C	1992–94	C	FL <120	-21.780	0.409	0.931	145	—	—	—	—	Karakulak & Oray (1994)

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Thunnus thynnus</i>	Aegean and Mediterranean Sea	C	1992–94	C	FL >120	-186.521	1.633	0.780	1494	-	-	-	-	Karakulak & Oray (1994)
<i>Torpedo marmorata</i>	Siagacik trawl area	C	2003	WI, SU	TL	0.0273	2.91	0.98	27	-	9.2	34.0	-	Filiz & Bilge (2004)
<i>Trachurus mediterraneus</i>	Eastern Black Sea	C	1996–97	-	TL	0.0108	2.980	-	-	-	6.3	17.8	-	Kayalı (1998)
<i>Trachurus mediterraneus</i>	Mid Black sea	C	-	-	-	0.0089	2.950	-	-	-	-	-	-	Anonymus (1991b)
<i>Trachurus mediterraneus</i>	Black Sea	C	-	-	-	0.3612	1.160	-	-	-	-	-	-	Düzungünes & Karaçam (1991)
<i>Trachurus mediterraneus ponticus</i>	Black Sea	C	-	-	FL	0.004834	3.218	-	601	-	6.5	17.5	-	Sahin et al. (1997)
<i>Trachurus trachurus</i>	Mid Black Sea	C	1995–96	C	TL	0.00759	3.05	-	720	14.13	9.4	16.8	0.07	Yücel & Erkoyuncu (2000)
<i>Trachurus trachurus</i>	Black Sea	C	1988–94	C	TL	0.0290	2.485	0.960	77	-	-	-	-	Erkoyuncu et al. (1994)
<i>Trachurus trachurus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000288	3.832	0.976*	259	-	9.0	29.6	-	Anonymus (1993)
<i>Trachurus trachurus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000071	3.072	0.955*	496	-	8.1	38.5	-	Anonymus (1993)
<i>Trachurus trachurus</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000139	3.964	0.974*	311	-	10.7	27.1	-	Anonymus (1993)
<i>Trigla lucerna</i>	Yumurtalık	C	1992–93	-	TL	-2.18077	3.088	0.966*	348	-	14.4	26.9	-	Altun et al. (1997)
<i>Trigla lucerna</i>	Black Sea	C	-	-	-	0.0085	3.032	-	-	-	-	-	-	Erdem et al. (1994)
<i>Trigla lucerna</i>	Black Sea	C	1988–94	C	TL	0.0070	3.089	0.980*	55	-	-	-	-	Erkoyuncu et al. (1994)
<i>Trisopterus minutus</i>	Çandarlı Bay	C	1993	SP	TL	0.0979	3.017	0.951*	35	-	-	-	-	Benli et al. (2000)
<i>Trisopterus minutus</i>	Çandarlı Bay	C	1993	AUT	TL	0.00709	3.144	0.910*	117	-	-	-	-	Benli et al. (2000)
<i>Upeneus moluccensis</i>	Gökova Bay	C	1993	AUT	TL	0.00606	3.351	0.962*	39	-	-	-	-	Benli et al. (2000)
<i>Upeneus moluccensis</i>	Mediterranean Sea	F	1991–92	C	FL	0.01051	3.150	0.945*	535	13.69	8.6	17.8	1.635	Kaya et al. (1999)
<i>Upeneus moluccensis</i>	Mediterranean Sea	M	1991–92	C	FL	0.00607	3.352	0.922*	176	11.16	8.5	16.1	1.378	Kaya et al. (1999)
<i>Upeneus moluccensis</i>	Fethiye Bay	C	1990–93	-	FL	-1.2889	3.386	0.828*	536	-	9.0	17.0	-	Torcu & Mater (1997)
<i>Upeneus moluccensis</i>	Mersin Bay	C	1990–93	-	FL	-1.0554	2.903	0.723*	500	-	8.9	16.0	-	Torcu & Mater (1997)
<i>Upeneus moluccensis</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SP	FL	0.0000043	3.257	0.976*	152	-	81.0	167.0	-	Anonymus (1993)

Species	Area	Sex	Year	S	L	a	b	r^2	N	mean	min	max	SD	Source
<i>Upeneus moluccensis</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	SU	FL	0.0000036	3.300	0.947*	300	—	93.0	176.0	—	Anonymus (1993)
<i>Upeneus moluccensis</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	AUT	FL	0.0000257	2.896	0.956*	207	—	86.0	178.0	—	Anonymus (1993)
<i>Upeneus moluccensis</i>	Marmara, Aegean and Mediterranean Sea	C	1991–93	WI	FL	0.0000027	3.353	0.941*	120	—	93.0	173.0	—	Anonymus (1993)
<i>Upeneus pori</i>	Babadilli-manı bight	F	1999–00	C	TL	0.0073	3.120	0.963	461	—	6.5	15.5	—	Cicek et al. (2002)
<i>Upeneus pori</i>	Babadilli-manı bight	M	1999–00	C	TL	0.0103	2.976	0.949	534	—	6.3	14.7	—	Cicek et al. (2002)
<i>Zosterisessor ophiocephalus</i>	Izmir Bay	C	1999–00	C	TL	0.0086	3.060	0.950	1066	—	8.0	23.3	—	Akyol (2003)

* r converted into r^2

** length-weight relationship corresponding to mm, g

RAZMERJA MED DOLŽINO IN TEŽO RIB, ŽIVEČIH V TURŠKIH MORJIH: PREGLED

Hatice TORCU KOÇ & Zeliha AKA ERDOĞAN

University of Balikesir, Faculty of Science and Arts, Department of Biology, TR-10100 Balikesir, Turkey

Jakov DULČIĆ

Inštitut za oceanografijo in ribištvo, HR-21000 Split, P.O. BOX 500, Hrvatska

E-mail: dulcic@izor.hr

POVZETEK

Članek navaja 360 razmerij med dolžino in težo rib, in sicer glede na podatke, ki so bili zbrani iz obstoječe literatur in zadevajo 90 vrst iz 40 družin, živečih v turških morjih. Naklon krivulje (b) se je gibal med 0,409 pri vrsti *Thunnus thynnus* in 4,343 pri vrsti *Dentex macrophthalmus*. Povprečna vrednost b je bila 3,088 ($\pm SD = 0,898$) in se ni bistveno razlikovala od 3 (t-test, $p < 0,05$). Mediana koeficiente b je bila 3,062 in 50% vrednosti b se je gibalo med 2,941 in 3,190.

Ključne besede: razmerje med dolžino in težo, ribe, turška morja, pregled

REFERENCES

- Aka, Z. (2003):** The examination of morphologic and genetic structure of anchovy (*Engraulis encrasicolus* (L. 1758)) in Turkish Seas. Ph.D. Thesis. University of Balikesir, Turkey.
- Akyol, O. (1999):** Investigations on population characteristics of *Mugil cephalus* (Linnaeus, 1758) and *Liza saliens* (Risso, 1810) in the Homa Lagoon (Izmir Bay, Aegean Sea). E. U. J. Fish. Aquat. Sci., 16(3–4), 391–419.
- Akyol, O. (2003):** Age, growth, reproduction period of grass goby (*Zosterisessor ophiocephalus*, Pallas 1811) in the Bay of Izmir (Aegean Sea). Arc. Fish. Mar. Res., 50(2), 1–5.
- Akyol, O., A. Tokaç & S. Unsal (1996):** An investigation on growth and reproduction of sardine (*Sardina pilchardus* Walbaum, 1792) in Izmir Bay. E. U. J. Fish. Aquat. Sci., 13(3–4), 383–394.
- Alpbaz, A. & T. Kinacigil (1988):** Investigations on the population of bluefish (*Pomatomus saltator* L., 1758) in Izmir Bay. E. U. J. Fish. Aquat. Sci., 5(19–20), 36–55.
- Altun, O. (2000):** Some biological aspects of the sand smelt (*Atherina hepsetus* Linnaeus, 1758) from Eceabat in the Dardanelles. E. U. J. Biol., 62, 69–77.
- Altun, A., M. Z. L. Göksu, C. Türeli & U. Erdem (1997):** An investigation on some biological characteristics of solea (*Solea vulgaris*) (Quensel, 1806) and Mediterranean gurnard (*Trigla lucerna* L., 1758) in the Yumurtalik Bay. Proc. 13th National Biology Congress. University of Istanbul, Istanbul, 17–20 September, 1996, p. 147–158.
- Anonymous (1991a):** Investigations on mullet *Mullus barbatus* in mid Black Sea. Turkish Ministry of Agricultural and Rural, Research Institute of Fisheries and Water Products, Trabzon.
- Anonymous (1991b):** Investigations on mackerel *Trachurus mediterraneus* in mid Black Sea. Turkish Ministry of Agricultural and Rural, Research Institute of Fisheries and Water Products, Trabzon.
- Anonymous (1991c):** Investigations on picarel *Spicara smaris* in mid Black Sea. Turkish Ministry of Agricultural and Rural, Research Institute of Fisheries and Water Products, Trabzon.
- Anonymous (1993):** Final report of demersal fisheries resource survey in the Republic of Turkey. Sanyo-Techno-Marine Inc., by Japan Intern. Cooperation Agency, 254 pp.
- Aral, O. & R. Bircan (1997):** Some population and reproduction characteristic of picarel (*Spicara smaris* L., 1758) in the Bay of Sinop. Turk. J. Vet. Anim. Sci., 21, 277–282.
- Avsar, D. (1995):** Sex, age composition and growth of the *Sprattus sprattus phalericus* (Risso, 1826) along the Turkish Black Sea Coast. Turk. J. Zool., 19, 157–163.
- Avsar, D. (1996):** Sex, age and growth of the spurdog (*Squalus acanthias* L., 1758) in the southeastern Black Sea. Yugosl. J. Operat. Res., 6(2), 295–304.
- Avsar, D. (1999):** Identification of turbot *Scophthalmus maximus* (L., 1758) stock along the Turkish Black Sea coast. Turk. J. Zool., 23(1), 207–213.
- Bagenal, T. B. & F. W. Tesch (1978):** Age and growth. In: Bagenal, T. (ed.): Methods for assessment of fish production in fresh waters. IBP Handbook 3. Blackwell Scientific Publications, Oxford, UK, p. 101–136.
- Bascınar, N. & I. Okumus (1997):** Determination of some bio-ecological characteristic of Pacific-Mullet (*Mugil so-iuy* Barlewsky, 1855). Proc. 13th National Biology Congress. University of Istanbul, Istanbul, 17–20 September, 1996, p. 425–434.
- Bilecenoglu, M. & M. Kaya (2002):** Growth of marbled spinefoot *Siganus rivulatus* Forsskål, 1775 (Teleostei: Siganidae) introduced to Antalya Bay, the eastern Mediterranean Sea, Turkey. Fish. Res., 54, 279–285.
- Buhan, E., H. Yılmaz, Y. Morkan, E. Büke & A. Yüksek (1997):** A new catch potential for Güllük bay and Gökovalı bay *Scomberomorus commerson* (Lacepede, 1800) (Pisces-Teleostei). Proc. Mediterranean Fisheries Congress, Izmir, 9–11 April, 1997, p. 937–944.
- Cailliet, G. M., M. S. Love & A. W. Ebeling (1986):** Fishes: A field and laboratory manual on their structure, identification, and natural history. Wadsworth Publishing Company, Belmont, California.
- Can, M. F., N. Basusta & M. Cekic (2002):** Weight-length relationships for selected fish species of the small-scale fisheries off the south coast of Iskenderun Bay. Turk. J. Vet. Anim. Sci., 26, 1181–1183.
- Celik, O. & H. Torcu (2000):** Investigations on the biology of Red Mullet (*Mullus barbatus* Linnaeus, 1758) in Edremit Bay, Aegean Sea, Turkey. Turk. J. Vet. Anim. Sci., 24, 287–296.
- Cicek, E., D. Avsar, H. Yeldan & M. Ozütok (2002):** Population characteristics of the Por's goatfish (*U. pori* Ben Tuvia and Golani, 1989) inhabiting Babadillimanı bight (Northeastern Mediterranean). Proc. Lessepsian Migration Workshop, Gökceada, 20–21 July 2002. Turk. Mar. Res. Found., 9, p. 92–99.
- Cihangir, B., A. Ünlüoğlu & M. Tirasin (1997):** Distribution and some biological aspects of the lesser spotted dogfish (Chondrichthyes, *Scyliorhinus canicula*, Linnaeus, 1758) from the northern Aegean Sea. Proc. Mediterranean Fisheries Congress, Izmir, 9–11 April 1997, p. 585–603.
- Dulčić, J. & M. Kraljević (1996):** Weight-length relationships for 40 fish species in the eastern Adriatic (Croatian waters). Fish. Res., 28(3), 243–251.
- Düzgüneş, E. & H. Karacam (1989):** Studies on some population parameters and growth of the European anchovy (*Engraulis encrasicolus*, Linneaus, 1758) in the Black Sea. Doga Turk. J. Zool., 13(2), 77–83.
- Düzgüneş, E. & H. Karacam (1990):** Some population aspects, meat yield and biochemical composition of whiting, *Gadus euxinus* (Nord. 1840) in the eastern Black Sea. Doga Turk. J. Zool., 14, 345–352.

- Düzungünes, E. & H. Karacam (1991):** Some population aspects, meat yield and biochemical composition of Mediterranean hores mackerel, *Trachurus mediterraneus* (Steindachner, 1968) in the Black Sea. *Doga Turk. J. Zool.*, 15, 195–201.
- Düzungünes, E., C. Mutlu & C. Sahin (1995):** Population parameters of anchovy (*Engraulis encrasicolus*, Linnaeus, 1758) in the eastern Black Sea. Proc. 2nd International Conference on the Mediterranean Coastal Environment (Medcoast 95), Tarrosgona, Spain, 24–27 October 1995, p. 56–66.
- Erdem, Y. (2001):** An investigation on selectivity of simple stretching nets used in fishery of turbot (*Scophthalmus maeticus* Pallas, 1881). Proc. 11th National Fishery Symposium, Hatay, 4–6 September 2001.
- Erdem, M., I. Erkoyuncu, O. Samsun & E. Ozdamar (1994):** Investigations on relationships of age-length, age-weight, the first catching age, and length. Research Found., Project no. S. 005, University of Ondokuz Mayıs.
- Erdem, Y., S. Ozdemir & C. Sümer (2001):** An investigation on food content of ray (*R. clavata* L.). Proc. 11th National Fishery Symposium, Hatay, 4–6 September 2001.
- Erkoyuncu, I., M. Erdem, O. Samsun, E. Erdamar & Y. Kaya (1994):** A research on the determination of meat yields, chemical composition and weight-length relationship of some fish species caught in the Black Sea. *I. U. J. Aquat. Prod.*, 8(1–2), 181–191.
- Filiz, H. & G. Bilge (2004):** Length-weight relationships of 24 fish species from the north Aegean Sea, Turkey. *J. Appl. Ichthyol.*, 20, 431–432.
- Goncalves, J. M. S., L. Bentes, P. G. Lino, J. Ribeiro, A. V. M. Canairo & K. Erzini (1997):** Weight-length relationships for selected fish species of the small-scale demersal fisheries of the south and south-west coast of Portugal. *Fish. Res.*, 30, 253–256.
- Gözler, A. M. & E. Ciloglu (1998):** A research on some population parameters of European anchovy (*Engraulis encrasicolus* L. 1758) caught in the Rize-Hopa coasts. Proc. 3rd Fisheries Symposium, Erzurum, Turkey, 10–12 June 1998, p. 373–382.
- Hossucu, B. (1992):** Investigations of biological aspects and distributions of sole (*Solea solea* L.). *E. U. J. Fish. Aquat. Sci.*, 9(33–34–36), 98–113.
- Hossucu, B. (2001):** Some growth parameters of mullet species (*Mugil* sp.) living in Güllük Lagoon (Aegean Sea). *E. U. J. Fish. Aquat. Sci.*, 18(3–4), 421–435.
- Ismen, A. (1995):** Growth, mortality and yield per recruit model of picarel (*Spicara smaris* L.) on the eastern Turkish Black Sea coast. *Fish. Res.*, 22(3–4), 299–308.
- Ismen, A. (2002):** Age, growth, sex ratio, spawning season of lizardfish (*S. undosquamis* Richardson, 1848) in the İskenderun Bay, the eastern Mediterranean Sea. Proc. Lessepsian Migration Workshop, Gökçeada, 20–21 July 2002. *Turk. Mar. Res. Found.*, 9, p. 108–115.
- Ismen, A. (2002):** A priliminary study on the population dynamics parameters of whiting (*Merlangius merlangus euxinus*) in Turkish Black Sea coastal waters. *Turk. J. Zool.*, 26, 157–166.
- Ismen, A. (2003):** Age, growth, reproduction and food of common stingray (*Dasyatis pastinaca* L., 1758) in İskenderun Bay, the eastern Mediterranean. *Fish. Res.*, 60, 169–176.
- Ismen, A., P. Ismen & N. Basuta (2004):** Age, growth and reproduction of tub gurnard (*Chelidonichthyes lucerna* L., 1758) in the Bay of İskenderun in the eastern Mediterranean. *Turk. J. Vet. Anim. Sci.*, 28, 289–29.
- Kara, F. & R. Gurbet (1990):** Distribution and measurement of stock dimensions of economically important demersal fishes in the northern Aegean Sea. E. U. Research Found., Project no. 002, 43 pp.
- Karacam, H. & E. Düzungünes (1990):** Age, growth and meat yield of the European anchovy (*Engraulis encrasicolus*, L. 1758) in the Black Sea. *Fish. Res.*, 9, 181–186.
- Karacam, H., I. Okumus, M. Feyzioglu & N. Sivri (1997):** A study on the growth, reproduction and feeding characteristics of spiny dogfish (*Squalus acanthias*, L. 1758) from the eastern Black Sea. Proc. 13th National Biology Congress, 17–20 September 1996, p. 322–331.
- Karakayis, M. & M. Togulga (2000):** Study on the biology of the sardine (*Sardina pilchardus* Walbaum, 1792) in the İzmir Bay (Aegean Sea). *E. U. J. Fish. Aquat. Sci.*, 17(3–4), 59–69.
- Karakulak, F. S. & I. K. Oray (1994):** The length-weight relationship of the bluefin tuna (*Thunnus thynnus* L., 1758) caught in Turkish waters. *I.U. J. Aquat. Prod.*, 8(1–2), 159–171.
- Karakulak, F. S. & I. K. Oray (2001):** Age et croissance du thon rouge (*Thunnus thynnus* L. 1758) peche de la mer Egee la mer Méditerranée en Turquie. *Rapp. Comm. int. Mer Médit.*, 36, p. 284.
- Kaya, M. & O. Ozaydin (1996):** A preliminary investigation on biology of *Capros aper* (L. 1758) (Pisces, Caproidae). *Turk. J. Zool.*, 20, 51–55.
- Kaya, M., H. A. Benli, T. Katagan & O. Ozaydin (1999):** Age, growth, sex-ratio, spawning season and food of golden banded goatfish, *Upeneus moluccensis* Bleeker (1855) from the Mediterranean and south Aegean Sea coasts of Turkey. *Fish. Res.*, 41, 317–328.
- Kaya, M., O. Ozaydin & H. Avni Benli (2001):** Age and growth parameters of red bandfish (*Cepola rubescens* L., 1766) in Izmir Bay. *Turk. J. Zool.*, 25, 111–116.
- Kayali, E. (1998):** An investigation on bio-ecological aspects of anchovy (*Engraulis encrasicolus*) and mackerel (*Trachurus mediterraneus*) in the eastern Black Sea. M.Sc. Thesis. University of Black Sea Technical, Trabzon, Turkey.
- Kinacigil, T. & O. Akyol (2001):** Effects on trawl selectivity of growth and reproduction in *Diplodus annularis* L. of Izmir Bay (Aegean Sea). *Arch. Fish. Mar. Res.*, 49(1), 1–8.

- Kınacigil, T., A. İlkyaz, O. Akyol, G. Metin, E. Cira & A. Ayaz (2001):** Growth parameters of Red mullet (*Mullus barbatus* L.) and seasonal cod-end selectivity of traditional bottom trawl nets in Izmir Bay. *Acta Adriat.*, 42(1), 113–123.
- Kınıkaslan, N. (1972):** Some investigations on the growth index, annual growth of the red mullet (*Mullus barbatus* L.) in the Edremit Bay (Aegean Sea). *Publ. Hydrobiological Research Institute, Faculty of Science, University of Istanbul.*, no. 8, 1–10.
- Koc, H. T., D. T. Cakir & J. Dulčić (2004):** Age, growth and mortality of comber *Serranus cabrilla* (Serranidae) in the Edremit Bay (NW Aegean Sea, Turkey). *Cybium*, 28(1), 19–25.
- Koc, H. T., D. T. Cakir & Z. Aka (2002):** Age, growth, sex-ratio, spawning season and mortality of annular bream, *Diplodus annularis* Linnaeus (1758) (Pisces: Sparidae) in Edremit Gulf (Aegean Sea). *Pakistan J. Biol. Sci.*, 5(10), 1126–1130.
- Koca, H. U. (2002):** A study on the determination of some parameters of the scorpion fish (*Scorpaena porcus*, Linne, 1758) caught by bottom nets in the area of Sinop in terms of fishery biology. *Turk. J. Vet. Anim. Sci.*, 26(1), 65–69.
- Kutaygil, N. & N. Bilecik (1998):** Studies on a shark species, picked dogfish (*Squalus acanthias* L.) distributed along the Anatolian littoral zones in the Black Sea. Ministry of Agricultural and Rural Affairs, Fisheries and Water Product Research Institute Press, Series B, no. 2, 73 pp.
- Martin-Smith, K. H. (1996):** Length-weight relationships of fishes in a diverse tropical freshwater community, sabah, Malaysia. *J. Fish. Biol.*, 49, 731–734.
- Mater, S. & H. Torcu (1997):** An investigation on biological and ecological aspects of an Red Sea Lizard fish, *Saurida undosquamis* (Richardson, 1848) living in Fethiye and Mersin Gulfs. *Proc. 13th National Biology Congress, Istanbul, 17–20 September 1996*, p.178–189.
- Mater, S. & B. Bayhan (1999):** Growth of the sardine (*Sardina pilchardus* Walbaum, 1792) distributing in İzmir Bay (Aegean Sea). *E. U. J. Fish. Aquat. Sci.*, 16(3–4), 257–267.
- Mater, S. & B. Bayhan (2000):** Investigation on the morphology and length-weight relationship of the scaldfish (*Arnoglossus laterna* Walbaum, 1792) in İzmir Bay (Aegean Sea). *E. U. J. Fish. Aquat. Sci.*, 17(1–2), 1–7.
- Mater, S., S. Malkav & B. S. Bayhan (2001):** A study on some biological peculiarities of the picarel (*Spicara flexuosa* Rafinesque, 1810) distributed in the bay of İzmir (Aegean Sea). *E. U. J. Fish. Aquat. Sci.*, 18(1–2), 25–32.
- Morato, T., P. Afonso, P. Lourinho, J. P. Barreiros, R. S. Santos & R. D. M. Nash (2001):** Length-weight relationships for 21 coastal fish species of the Azores, north-eastern Atlantic. *Fish. Res.*, 50, 297–302.
- Mutlu, C. (1994):** An investigation on some population characters of anchovy (*Engraulis encrasicolus*, Linnaeus, 1758) in the eastern Black Sea. M.Sc. Thesis. University of Black Sea Technical, Trabzon, Turkey.
- Ozaydin, O. (1997):** Investigations on biology and feeding regimes of some species (Pisces: Sparidae) living in Aegean Sea. Ph.D. Thesis. University of Ege, Izmir, Turkey.
- Ozdamar, E., L. Erkoyuncu, M. Erdem & S. Chen (1991a):** Effects of fishing on decreasing of stocks of *Engraulis encrasicolus* L. 1758 in the Black Sea recently. *Proc. 1st Interuniversity Conference of Scientific, Cultural, Social and Economic Possibilities of Collaboration between Georgia and Turkey, Batumi, 9–14 December 1991*.
- Ozdamar, E., K. Kohei & I. Erkoyuncu (1991b):** Some biological characteristics of European anchovy *Engraulis encrasicolus* L. 1758 in the Black Sea. *J. Tokyo Univ. Fish.*, 78(1), 57–64.
- Petrakis, G. & K. I. Stergiou (1995):** Weight-length relationships for 33 fish species in Greek waters. *Fish. Res.*, 21, 465–469.
- Safran, P. (1992):** Theoretical analysis of the weight-length relationships in the juveniles. *Mar. Biol.*, 112, 545–551.
- Samsun, O. (1995):** The determination of some parameters about growth characteristics of garfish (*Belone belone euxini* Günther, 1866) caught in the area of Sinop (Black Sea). *E. U. J. Fish. Aquat. Sci.*, 12(3–4), 347–355.
- Samsun, O. & I. Erkoyuncu (1992):** Research on some characteristics of the red mullet (*Mullus barbatus ponticus* Ess. 1927) caught by the trawler in the mid Black Sea region from the point of view of fishery biology. *Proc. 11th National Biology Congress, University of Fırat, Elazığ*, 3, p.189–198.
- Samsun, O., E. Ozdamar & O. Aral (1993):** An investigation in the base of fishery biology of whiting *Gadus merlangus euxinus* caught by deep trol in mid Black Sea trawling areas. *Proc. 1st National Ecology and Environment Congress*.
- Samsun, N. & I. Erkoyuncu (1998):** The research on the estimation of some parameters of whiting (*Gadus merlangus euxinus* Nordmann, 1840) caught by the bottom trawlers in the area of Sinop (Black Sea) from the view point of fishery biology. *E. U. J. Fish. Aquat. Sci.*, 15(1–2), 19–31.
- Sahin, T. (1999):** Some biological characteristics of sprat (*Sprattus sprattus phalericus* Russo, 1826) on the eastern Black Sea coast. *Turk. J. Zool.*, 23(1), 249–255.
- Sahin, T. & B. Akbulut (1997):** Some population aspects of whiting (*Merlangius merlangius euxinus* Nordmann, 1840) in the eastern Black Sea coast of Turkey. *Turk. J. Zool.*, 21, 187–193.
- Sahin, T. & B. Akbulut (1997):** Some biologic characteristics of *Mullus barbatus ponticus* Essipov, 1927 in the eastern Black Sea coast of Turkey. *Turk. J. Zool.*, 21(2), 179–185.

- Sahin, T. & Y. Genc (1999):** Some biological characters of picarel (*Spicara smaris*, Linnaeus, 1758) in the eastern Black Sea coast of Turkey. Turk. J. Zool., 23(1), 149–155.
- Sahin, T., Y. Genc & H. Okur (1997):** Investigations of the growth and reproduction of horse mackerel (*T. mediterraneus ponticus* Aliev) population in Turkish Black Sea coast. Turk. J. Zool., 21, 321–328.
- Stergiou, K. I. & D. K. Moutopoulos (2001):** A review of length-weight relationships of fishes from Greek Marine Waters. Naga, The ICLARM Quarterly, 24(1–2), 23–39.
- Togulga, M. (1977):** Research on the biology and population dynamics of *Mullus barbatus* L., from the Izmir Bay. E. U. J. Fac. Sci., Ser. B, 1(2).
- Torcu, H. & S. Mater (1997):** An investigation on biological aspects of the golden-banded goat fish *Upeneus moluccensis* (Bleeker, 1859) living in Fethiye and Mersin Gulf. Proc. Mediterranean Fisheries Congress, 9–11 April 1996, p. 545–554.
- Torcu, H., D. Türker & S. Mater (1998):** A preliminary study on some biological aspects of the population of comber (*Serranus cabrilla*, Linnaeus, 1758) in Edremit bay, Northern Aegean Sea. The eastern Anatolian Region. Proc. 3rd Symposium on Water Products, University of Atatürk, Erzurum, p. 611–616.
- Torcu, H., Ö. Çelik, Z. Aka & D. Türker (1997):** Investigations on biological aspects of hake population (*Merluccius merluccius*, Linnaeus, 1758) in Edremit bay, Northern Aegean Sea. Populasyonunun Biyolojik Özellikleri Üzerine Araştırmalar, 1997. Proc. 9th Symposium on Water Products, Egirdir, Isparta, I, p. 152–162.
- Tosunoğlu, Z., O. Akyol, G. Metin, A. Tokaç & S. Ünsal (1997):** The study on the population characteristics of three sparid species in the Gülbahçe Bay. E. U. J. Fish. Aquat. Sci., 14(1–2), 127–143.
- Türeli, C. & Ü. Erdem (1997):** The growth performance of red mullet (*Mullus barbatus* Linneaus, 1758) and brushtooth lizardfish (*Saurida undosquamis* (Richardson, 1848)) from the costal region of Adana province (Iskenderun Bay, Turkey). Turk. J. Zool., 21, 329–334.
- Türkmen, M. (2003):** Investigation of some population parameters of common sole, *Solea solea* (L., 1758) from İskenderun Bay. Turk. J. Vet. Anim. Sci., 27, 317–323.
- Türkmen, M. & I. Akyurt (2003):** Growth characteristics, sex inversion and mortality rates of striped sea bream, *Lithognathus mormyrus* L., in İskenderun Bay. Turk. J. Zool., 27, 323–329.
- Uckun, D., M. Togulga & E. Taskavak (2000):** A preliminary study on the growth of the common hake (*Merluccius merluccius* L., 1758) in İzmir Bay, Aegean Sea. Acta Adriat., 41(2), 25–34.
- Ünsal, N. (1989):** A study on age-length-weight relationship and determination of the smallest catching size of anchovy, *Engraulis encrasicolus* (L.) in the Black Sea. I. U. J. Aquat. Prod., 3(1–2), 17–28.
- Üysal, A. (1990):** An investigation on biology and population dynamics whiting (*Merlangius merlangus euxinus* Nordman, 1840) in the eastern Black Sea, Sinop-Hopa. Ph.D. Thesis. University of Istanbul, Turkey.
- Yücel, S. & I. Erköyuncu (2000):** Population dynamics of horse mackerel (*Trachurus trachurus* L. 1758) stocks in the Mid Black Sea, Turkey. Turk. J. Biol., 24, 543–552.