

**IMFM**

INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS  
JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

**Preprint series**

**Vol. 51 (2013), 1189**

ISSN 2232-2094

**AVERAGE DISTANCE,  
SURFACE AREA, AND  
OTHER STRUCTURAL  
PROPERTIES OF  
EXCHANGED HYPERCUBES**

Sandi Klavžar      Meijie Ma

Ljubljana, August 23, 2013

# Average distance, surface area, and other structural properties of exchanged hypercubes

Sandi Klavžar

Faculty of Mathematics and Physics

University of Ljubljana, Slovenia

and

Faculty of Natural Sciences and Mathematics

University of Maribor, Slovenia

and

Institute of Mathematics, Physics and Mechanics, Ljubljana

`sandi.klavzar@fmf.uni-lj.si`

Meijie Ma

Department of Mathematics

Zhejiang Normal University

Jinhua, Zhejiang, 321004, China

`mameij@mail.ustc.edu.cn`

## Abstract

Exchanged hypercubes [Loh et al., IEEE Transactions on Parallel and Distributed Systems 16 (2005) 866–874] are spanning subgraphs of hypercubes with about one half of their edges but still with many desirable properties of hypercubes. In this paper it is shown that also distance properties of exchanged hypercubes are comparable to the corresponding properties of hypercubes. The average distance and the surface area of exchanged hypercubes are computed and it is shown that exchanged hypercubes have asymptotically the same average distance as hypercubes. Several additional metric and other properties are also deduced and it is proved that exchanged hypercubes are prime with respect to the Cartesian product of graphs.

**Key words:** interconnection network; exchanged hypercube; Wiener index; average distance; surface area; Cartesian product of graphs

**AMS Subj. Class.:** 05C12, 05C82, 68M10

## 1 Introduction

Hypercubes form one of the fundamental models for interconnection networks. They are universal in the sense that the binary strings are naturally encoded into their topology.

Consequently, they possess numerous fine network properties such as small diameter, small density, and high connectivity. At the same time, straightforward local routing is possible. Nevertheless, from several reasons different variations of hypercubes were proposed, let us mention here only Möbius cubes [5], folded hyper-star networks [16], Fibonacci cubes [17], folded cubes [19],  $k$ -ary  $n$ -cubes [28], and twisted-cubes [29]. An additional variation of hypercubes—exchanged hypercubes—is of our prime interest here.

The exchanged hypercubes  $EH(s, t)$ , proposed by Loh et al. [23], are graphs obtained by systematically removing edges from hypercubes. The number of vertices in  $EH(s, t)$  is equal to that of the  $(s + t + 1)$ -dimensional hypercube, but the ratio of the number of edges in  $EH(s, t)$  to that of the  $(s + t + 1)$ -dimensional hypercube is  $1/2 + 1/(2(s + t + 1))$  [4]. Different properties of exchanged hypercubes were investigated by now. The bipancyclicity of them was investigated in [24]. To measure the fault-tolerance of them, the connectivity and the super connectivity were determined in [20, 25, 26]. These results, in addition to those obtained in the seminal paper [23], indicate that the exchanged hypercubes keep numerous desirable properties of the hypercubes. In this paper we take a closer look to metric properties of exchanged hypercubes and some other related properties.

Graphs considered here are simple, finite, and connected. The *distance*  $d_G(u, v)$  between vertices  $u$  and  $v$  of a (connected) graph  $G$  is the usual shortest-path distance. If  $G$  will be clear from the context, we will simply write  $d(u, v)$ . The *Wiener index*  $W(G)$  of a graph  $G$  is the sum of the distances between all unordered pairs of vertices of  $G$ . For instance,  $W(K_n) = \binom{n}{2}$ , that is, the number of its edges. This graph invariant is one of the fundamental properties of (interconnection) networks, cf. [2, 4, 6, 18, 27]. On the other hand, the Wiener index of a graph is a central and one of the most studied invariants in mathematical chemistry, see for instance surveys [8, 9].

A concept closely related to the Wiener index is the surface area. If  $v$  is a vertex of a graph  $G$  and  $r$  a positive integer, then the  *$r$ -surface area*  $B_{G,v}(r)$  of  $G$  centered at  $v$  is the number of vertices at distance  $r$  from the fixed base vertex  $v$ . That is, the surface area is the size of the  $r$ -sphere around  $v$ . If  $G$  is vertex-transitive, then the surface area is independent of a selected vertex. The surface area of a network is interesting because many network properties and algorithms are directly related to it. Consequently, it has been studied for a variety of networks, cf. [6, 7, 14].

The paper is organized as follows. In the next section we introduce the exchanged hypercubes in two equivalent ways and recall or deduce some of their basic properties. Then, in Section 3, we first determine the Wiener index of exchanged hypercubes. As a consequence it is demonstrated that asymptotically, exchanged hypercubes and hypercubes have the same average distance. Then we determine the surface area and indicate that this result yields an alternative way to determine the Wiener index. In Section 4 we first consider several other distance-based invariants and then prove that the exchanged hypercubes are prime with respect to the Cartesian product operation. In the final section we point out that dual-cubes are particular instances of exchanged hypercubes and list some of consequences of our results for the dual-cubes.

## 2 Exchanged hypercubes: two definitions and some properties

In this section we first introduce exchanged hypercubes and list some of their properties. Then we equivalently describe these cubes as the graphs obtained by adding a perfect matching between two collections of hypercubes and show how this point of view can be used to infer additional properties.

Exchanged hypercubes are spanning subgraphs of hypercubes. Recall that if  $d$  is a positive integer, then the  $d$ -dimensional *hypercube* (or  $d$ -*cube*, for short)  $Q_d$  is the graph with vertex set  $\{0, 1\}^d$ , two vertices (strings) being adjacent if they differ in exactly one coordinate. The *Hamming distance*  $H(b, c)$  between binary strings  $b$  and  $c$  of the same length is the number of positions in which  $b$  and  $c$  differ. It is well known that  $d_{Q_d}(u, v) = H(u, v)$  holds for any two vertices (alias strings) of  $Q_d$ .

Let  $u = u_{d-1} \dots u_0 \in \{0, 1\}^d$  be a binary string,  $d \geq 1$ . If  $j \geq i$ , then we will use the notation  $u_{j:i}$  for the substring of  $u$  between  $u_j$  and  $u_i$ , that is,  $u_{j:i} = u_j \dots u_i$ . For any integers  $s \geq 1$  and  $t \geq 1$ , the *exchanged hypercube*  $EH(s, t)$  is the graph with the vertex set  $\{0, 1\}^{s+t+1}$ . Hence, if  $u \in V(EH(s, t))$ , then its coordinates are  $u_{s+t} \dots u_{t+1} u_t \dots u_1 u_0$ . Vertices  $u$  and  $v$  are adjacent if one of the following conditions is satisfied:

- (i)  $u_{s+t:1} = v_{s+t:1}, u_0 \neq v_0$ ,
- (ii)  $u_0 = v_0 = 1, H(u_{t:1}, v_{t:1}) = 1$ , and  $u_{s+t:t+1} = v_{s+t:t+1}$ ,
- (iii)  $u_0 = v_0 = 0, H(u_{s+t:t+1}, v_{s+t:t+1}) = 1$ , and  $u_{t:1} = v_{t:1}$ .

For instance,  $EH(1, 2)$  and  $EH(2, 2)$  are shown in Fig. 1.




Figure 1: Exchanged hypercubes  $EH(1, 2)$  and  $EH(2, 2)$

Clearly,  $EH(s, t)$  has  $2^{s+t+1}$  vertices. Note also that if  $u \in V(EH(s, t))$  and  $u_0 = 0$ , then the degree of  $u$  is  $s + 1$ , otherwise the degree of  $u$  is  $t + 1$ . It is also straightforward that for any  $s$  and  $t$ , the exchanged hypercube  $EH(s, t)$  is isomorphic to  $EH(t, s)$ .

For practical purposes, another description of the exchanged hypercubes is useful. Note that the edge set of  $EH(s, t)$  is the disjoint union of sets  $E_1, E_2, E_3$ , where

$$\begin{aligned}
 E_1 &= \{uv : u_{s+t:1} = v_{s+t:1}, u_0 \neq v_0\}, \\
 E_2 &= \{uv : u_{s+t:t+1} = v_{s+t:t+1}, H(u_{t:1}, v_{t:1}) = 1, u_0 = v_0 = 1\}, \\
 E_3 &= \{uv : u_{t:1} = v_{t:1}, H(u_{s+t:t+1}, v_{s+t:t+1}) = 1, u_0 = v_0 = 0\}.
 \end{aligned}$$

In Fig. 1 the edges from  $E_1, E_2$  and  $E_3$  are represented by dashed lines, thin lines, and thick lines, respectively.

Let  $EH_1(s, t)$  be the subgraph of  $EH(s, t)$  induced by the edges  $E_2$ . Then  $EH_1(s, t)$  is the disjoint union of  $2^s$  copies of  $Q_t$ . Indeed, fixing the leftmost  $s$  bits and fixing the rightmost bit to 1, the induced subgraph is isomorphic to  $Q_t$ . Moreover, there are no edges between two such induced subgraphs isomorphic to  $Q_t$ . Similarly, the subgraph  $EH_0(s, t)$  of  $EH(s, t)$  induced by the edges  $E_3$  consists of  $2^t$  subgraphs isomorphic to  $Q_s$ . Finally, the edges from  $E_1$  form a perfect matching of  $EH(s, t)$ , it is a matching between  $EH_0(s, t)$  and  $EH_1(s, t)$ . More precisely, let  $Q$  be an arbitrary copy of  $Q_t$  from  $EH_1(s, t)$ . Then each vertex of  $Q$  has exactly one neighbor in  $EH_0(s, t)$ , each of these neighbors belongs to a different copy of  $Q_s$  from  $EH_0(s, t)$ . For instance, in the graph

$EH(2, 2)$  in Fig. 1, the subgraph  $EH_1(2, 2)$  consists of 4 copies of  $Q_2$  drawn with thin lines, the subgraph  $EH_0(2, 2)$  consists of 4 copies of  $Q_2$  drawn with thick lines, while the matching edges (that is, the edges from  $E_1$ ) are drawn with dashed lines.

Using the above representation of the exchanged hypercubes it is immediate that  $|E(EH(s, t))| = 2^t(s2^{s-1}) + 2^s(t2^{t-1}) + 2^t2^s = 2^{s+t-1}(s + t + 2)$ , cf. [4]. Since  $Q_{s+t+1}$  has  $(s + t + 1)2^{s+t}$  edges, it thus follows that  $EH(s, t)$  has about one half of the edges of the hypercube of the same dimension. As another property we have:

**Proposition 2.1** *If  $s, t \geq 1$ , then the number of 4-cycles of  $EH(s, t)$  is*

$$2^{s+t-2} \left( \binom{s}{2} + \binom{t}{2} \right).$$

**Proof.** From the above representation we infer that a 4-cycle of  $EH(s, t)$  contains no edge between  $EH_0(s, t)$  and  $EH_1(s, t)$ . It follows that any 4-cycle lies either in  $EH_0(s, t)$  or in  $EH_1(s, t)$ . As these two subgraphs are disjoint unions of hypercubes  $Q_s$  and  $Q_t$ , respectively, and since  $Q_n$  contains exactly  $2^{n-2} \binom{n}{2}$  4-cycles, cf. [15, Exercise 2.4], the assertion follows immediately.  $\square$

We have noted that the edges from  $E_1$  form a perfect matching of  $EH(s, t)$ . But exchanged hypercubes contain many additional perfect matchings. Indeed, hypercubes contain a huge number of perfect matchings, cf. [11], hence any combination of separate perfect matchings in copies of  $Q_s$  and  $Q_t$  gives a perfect matching in  $EH(s, t)$ . Moreover, Fink [10] proved that every perfect matching of  $Q_d$ ,  $d \geq 2$ , is contained in a Hamiltonian cycle. This result together with the second description of the exchanged hypercubes can be used to construct many Hamiltonian cycles in exchanged hypercubes  $EH(s, s)$ , see [3].

The *domination number*  $\gamma(G)$  of a graph  $G$  is the order of a smallest set  $X \subseteq V(G)$  such that any vertex  $u \in V(G) \setminus X$  has at least one neighbor in  $X$ . The exact value of  $\gamma(Q_d)$  is known only for  $d \leq 6$  and for  $d = 2^k - 1$  or  $d = 2^k$ , see [15, p. 90]. Using the second representation of  $EH(s, t)$  it follows immediately that  $\gamma(EH(s, t)) \leq 2^s\gamma(Q_t) + 2^t\gamma(Q_s)$ . With a little effort we can say a bit more:

**Proposition 2.2** *If  $s, t \geq 1$  and  $s \leq t$ , then  $\gamma(EH(s, t)) \leq 2^s\gamma(Q_t) + (2^t - 1)\gamma(Q_s)$ .*

**Proof.** Select an arbitrary  $s$ -cube  $Q$  from  $EH_0(s, t)$ . Let  $Q_t^{(i)}$ ,  $1 \leq i \leq 2^s$ , be the  $t$ -cubes from  $EH_1(s, t)$  that have a neighbor in  $Q$ . In each of the cubes  $Q_t^{(i)}$  select

a minimum dominating set such that it contains the vertex that has a neighbor in  $Q$ . (Such a dominating set exists since hypercubes are vertex-transitive graphs.) In this way  $Q$  is dominated by vertices from  $EH_1(s, t)$ . Selecting an arbitrary minimum dominating set in each of the remaining  $s$ -cubes and  $t$ -cubes, the result follows.  $\square$

The bound of Proposition 2.2 is (of course) not optimal. For instance,  $EH(2, 2)$  contains a perfect code and consequently  $\gamma(EH(2, 2)) = 8$ . It can also be shown that  $\gamma(EH(2, 3)) \leq 16$ . We believe that to determine the exact value of  $\gamma(EH(s, t))$  is a difficult problem which is worth of an independent study.

### 3 Average distance and surface area

Using the second description of exchanged hypercubes from the previous section we prove the following key lemma. It is implicitly given in [23, Table 2], but since it is the key lemma, we give a formal proof of it.

**Lemma 3.1** *If  $s, t \geq 1$  and  $u, v \in V(EH(s, t))$ , then*

$$d(u, v) = \begin{cases} H(u, v) + 2, & u_0 = v_0 = 0, u_{t:1} \neq v_{t:1}, \text{ or} \\ & u_0 = v_0 = 1, u_{s+t:t+1} \neq v_{s+t:t+1}; \\ H(u, v), & \text{otherwise.} \end{cases}$$

**Proof.** Note first that  $d(u, v) \geq H(u, v)$  because  $EH(s, t)$  is a spanning subgraph of  $Q_{s+t+1}$  and  $d_{Q_{s+t+1}}(u, v) = H(u, v)$ .

Suppose first that  $u_0 = v_0 = 0$  and  $u_{t:1} = v_{t:1}$ . Then  $u$  and  $v$  belong to the same subgraph  $Q_s$  of  $EH_0(s, t)$  and hence  $d(u, v) \leq H(u, v)$ . Thus  $d(u, v) = H(u, v)$ . Analogously we get the same conclusion when  $u_0 = v_0 = 1$  and  $u_{s+t:t+1} = v_{s+t:t+1}$ . Let next  $u_0 \neq v_0$  and assume without loss of generality that  $u_0 = 0$  and  $v_0 = 1$ . Then a  $u, v$ -path of length  $H(u, v)$  can be constructed as follows. First change one by one the bits of  $u$  between  $u_{s+t}$  and  $u_{t+1}$  in which  $u$  differs from  $v$ . Then change the rightmost bit to 1, and finally change the remaining bits in which  $u$  differs from  $v$ . Hence in all these cases we have  $d(u, v) = H(u, v)$ .

Assume now that  $u_0 = v_0 = 0$  and  $u_{t:1} \neq v_{t:1}$ . Then  $u$  and  $v$  belong to different copies of  $Q_s$  from  $EH_0(s, t)$ . Hence any  $u, v$ -path necessarily contains at least two matching edges between  $EH_0(s, t)$  and  $EH_1(s, t)$ . Then it follows that  $d(u, v) \geq H(u, v) + 2$ . We can find a  $u, v$ -path of length  $H(u, v) + 2$  as follows: change one by one

the bits of  $u$  between  $u_{s+t}$  and  $u_{t+1}$  in which  $u$  differs from  $v$ , then change the rightmost bit to 1, change the remaining bits in which  $u$  differs from  $v$ , and finally change the rightmost bit to 0. We conclude that  $d(u, v) = H(u, v) + 2$ . The case  $u_0 = v_0 = 1$  and  $u_{s+t:t+1} \neq v_{s+t:t+1}$  is treated analogously.  $\square$

**Theorem 3.2** *If  $s, t \geq 1$ , then*

$$W(EH(s, t)) = (s + t + 3)2^{2(s+t)} - 2^{2s+t} - 2^{s+2t}.$$

**Proof.** By Lemma 3.1, the contribution of a pair  $\{u, v\} \in \binom{V(EH(s, t))}{2}$  to  $W(EH(s, t))$  is either  $H(u, v)$  or  $H(u, v) + 2$ . Thus  $W(EH(s, t))$  is the sum of  $W(Q_{s+t+1})$  and twice the number of pairs of vertices  $\{u, v\}$  with  $d(u, v) = H(u, v) + 2$ .

Consider first pairs with  $u_0 = v_0 = 0$  and  $u_{t:1} \neq v_{t:1}$ . In the subcase when  $u_{s+t:t+1} = v_{s+t:t+1}$ , there are  $2^s$  possible substrings for the first  $s$  coordinates and hence the number of (unordered) pairs of vertices with  $u_0 = v_0 = 0$ ,  $u_{t:1} \neq v_{t:1}$ , and  $u_{s+t:t+1} = v_{s+t:t+1}$ , is

$$\binom{2^t}{2} 2^s.$$

If on the other hand  $u_{s+t:t+1} \neq v_{s+t:t+1}$ , we have  $\binom{2^s}{2}$  pairs with respect to the first  $s$  coordinates and  $\binom{2^t}{2}$  for the consecutive  $t$  coordinates. As such pairs can be combined in two ways, the number of such (unordered) pairs of vertices is

$$2 \binom{2^s}{2} \binom{2^t}{2}.$$

Similarly, the number of pairs  $\{u, v\}$  with  $u_0 = v_0 = 1$  and  $u_{s+t:t+1} \neq v_{s+t:t+1}$  is equal to

$$\binom{2^s}{2} 2^t + 2 \binom{2^s}{2} \binom{2^t}{2}.$$

It is well-known (cf. [12, Exercise 19.3]) that  $W(Q_{s+t+1}) = (s + t + 1)2^{2(s+t)}$ . Putting all this together we obtain that

$$W(EH(s, t)) = (s + t + 1)2^{2(s+t)} + 2 \left( \binom{2^t}{2} 2^s + \binom{2^s}{2} 2^t + 4 \binom{2^s}{2} \binom{2^t}{2} \right),$$

which, after a routine computation, reduces to the claimed expression.  $\square$

The *average distance*  $\mu(G)$  of a graph  $G$  is defined with

$$\mu(G) = \frac{1}{\binom{|V(G)|}{2}} W(G).$$

We note that some authors prefer to define the average distance as  $2W(G)/|V(G)|^2$ . The definitions are almost equivalent and are in fact equivalent in the asymptotic sense, which we consider next.

**Corollary 3.3**

$$\lim_{s,t \rightarrow \infty} \frac{\mu(EH(s,t))}{s+t+1} = \lim_{d \rightarrow \infty} \frac{\mu(Q_d)}{d} = \frac{1}{2}.$$

**Proof.** Using Theorem 3.2 we find that

$$\frac{W(EH(s,t))}{W(Q_{s+t+1})} = \frac{1}{s+t+1} (s+t+3 - 2^{-s} - 2^{-t}),$$

hence  $\lim_{s,t \rightarrow \infty} \frac{\mu(EH(s,t))}{s+t+1} = \lim_{s,t \rightarrow \infty} \frac{\mu(Q_{s+t+1})}{s+t+1}$ . For the latter limit we have

$$\lim_{d \rightarrow \infty} \frac{\mu(Q_d)}{d} = \lim_{d \rightarrow \infty} \frac{d 2^{2d-2}}{\binom{2^d}{2} d} = \frac{1}{2},$$

and we are done. □

We next determine the surface area of exchanged hypercubes:

**Theorem 3.4** *If  $s, t \geq 1$ , then*

$$B_{EH(s,t),v}(r) = \begin{cases} \binom{s+t+1}{r-1} + \binom{s}{r} - \binom{s}{r-2}, & v_0 = 0; \\ \binom{s+t+1}{r-1} + \binom{t}{r} - \binom{t}{r-2}, & v_0 = 1. \end{cases}$$

**Proof.** Assume  $v_0 = 0$  and  $d(u, v) = r$  ( $0 \leq r \leq s+t+2$ ). We count the number of  $u$  according to the three cases.

**Case 1:**  $u_0 = 0, u_{t:1} = v_{t:1}$ .

By Lemma 3.1, there are  $r$  different bits between  $u_{s+t:t+1}$  and  $v_{s+t:t+1}$ . The number of  $u$  is  $\binom{s}{r}$ . Note that  $r \leq s$  in this case.

**Case 2:**  $u_0 = 0, u_{t:1} \neq v_{t:1}$ .

By Lemma 3.1, there are  $r-2$  different bits between  $u_{s+t:t+1}$  and  $v_{s+t:t+1}$ . Note that  $u_{t:1} \neq v_{t:1}$  and  $r \geq 3$ . The number of  $u$  is  $\binom{s+t}{r-2} - \binom{s}{r-2}$ .

**Case 3:**  $u_0 = 1$ .

By Lemma 3.1, there are  $r-1$  different bits between  $u_{s+t:t+1}$  and  $v_{s+t:t+1}$ . The number of  $u$  is  $\binom{s+t}{r-1}$ .

Combining the above three cases we conclude that

$$B_{EH(s,t),v}(r) = \binom{s}{r} + \binom{s+t}{r-2} - \binom{s}{r-2} + \binom{s+t}{r-1} = \binom{s+t+1}{r-1} + \binom{s}{r} - \binom{s}{r-2}.$$

We proceed analogously when  $v_0 = 1$ . □

We conclude the section by noting that Theorem 3.4 yields another proof of Theorem 3.2. The computations are rather technical, so we state only the key steps. If  $v_0 = 0$ , the total distance of  $v$  is

$$\begin{aligned} \sum_{u \in V(EH(s,t))} d(u, v) &= \sum_{r=1}^{s+t+2} r \cdot B_{EH(s,t),v}(r) \\ &= \sum_{r=1}^{s+t+2} r \left[ \binom{s+t+1}{r-1} + \binom{s}{r} - \binom{s}{r-2} \right] \\ &= (s+t+3)2^{s+t} - 2^{s+1}. \end{aligned}$$

Analogously, if  $v_0 = 1$ , then the total distance of  $v$  is  $(s+t+3)2^{s+t} - 2^{t+1}$ . The number of vertices with  $v_0 = 0$  (or  $v_0 = 1$ ) is  $2^{s+t}$ . Hence, the Wiener index of  $EH(s, t)$  is  $2^{s+t}[(s+t+3)2^{s+t} - 2^{s+1} + (s+t+3)2^{s+t} - 2^{t+1}]/2 = (s+t+3)2^{2(s+t)} - 2^{2s+t} - 2^{s+2t}$ .

## 4 Additional properties

Recall that the *eccentricity*  $\text{ecc}(u)$  of a vertex  $u \in V(G)$  is the maximum distance between  $u$  and the vertices of  $G$ . The *radius*  $\text{rad}(G)$  and the *diameter*  $\text{diam}(G)$  of  $G$  are the minimum and the maximum eccentricity in  $G$ , respectively.

From Lemma 3.1 we can also determine the eccentricity of vertices of  $EH(s, t)$ . Let  $u$  be an arbitrary vertex, then its binary complement  $\bar{u}$  is the unique vertex with respect to  $u$  with the property  $H(u, \bar{u}) = s+t+1$ . By Lemma 3.1 we infer that  $d(u, \bar{u}) = s+t+1$ . On the other hand, the lemma implies that  $d(u, \bar{u}') = s+t+2$ , where  $\bar{u}'$  is the vertex obtained from  $\bar{u}$  by complementing  $\bar{u}_0$ . Moreover, using Lemma 3.1 again, we also observe that the vertex  $\bar{u}'$  is the unique vertex to  $u$  such that  $\text{ecc}(u) = d(u, \bar{u}')$ . We collect these facts into the following result, where the average eccentricity (cf. [13]) is defined in the natural way. (We note that the diameter of  $EH(s, t)$  was earlier determined in [23, Theorem 6].)

**Proposition 4.1** *If  $u \in V(EH(s, t))$ , then  $\text{ecc}(u) = s + t + 2$  and hence the average eccentricity of  $EH(s, t)$  is  $s + t + 2$ . Moreover, every vertex has a unique antipodal vertex.*

**Corollary 4.2** *If  $s, t \geq 1$ , then  $\text{diam}(EH(s, t)) = \text{rad}(EH(s, t)) = s + t + 2$ .*

This should be compared with the fact that  $\text{diam}(Q_{s+t+1}) = \text{rad}(Q_{s+t+1}) = s+t+1$ . Hence again the exchanged hypercubes keep these fine properties of hypercubes.

The *Cartesian product*  $G \square H$  of graphs  $G$  and  $H$  is the graph with vertex set  $V(G) \times V(H)$ , vertices  $(g, h)$  and  $(g', h')$  being adjacent whenever  $gg' \in E(G)$  and  $h = h'$ , or  $g = g'$  and  $hh' \in E(H)$ . For more information on this graph operation see [12]. For our purpose it is essential to recall that  $Q_d$  can be represented as the  $d$ -fold Cartesian product of  $K_2$ . Hence hypercubes are in a way the simplest possible Cartesian product graphs. A graph that cannot be represented as the Cartesian product of graphs on at least two vertices, is called *prime (with respect to the Cartesian product)*. We conclude this note with the following results that contrasts exchanged hypercubes from hypercubes.

**Proposition 4.3** *If  $s, t \geq 1$ , then  $EH(s, t)$  is a prime graph.*

**Proof.** Consider vertices  $u = 00 \dots 00$  and  $v = 00 \dots 01$ . Clearly,  $uv \in E(EH(s, t))$ . The open neighborhood  $N(u)$  of  $u$  is  $\{v, u^{(t+1)}, \dots, u^{(s+t)}\}$ , where  $u_i^{(i)} = 1$  for  $t+1 \leq i \leq s+t$ , and  $u_j^{(i)} = 0$  for  $t+1 \leq i \leq s+t$ ,  $0 \leq j \leq s+t$ ,  $j \neq i$ . Similarly,  $N(v) = \{u, v^{(1)}, \dots, v^{(t)}\}$ , where  $v_i^{(i)} = v_0^{(i)} = 1$  for  $1 \leq i \leq t$ , and  $v_j^{(i)} = 0$  for  $1 \leq i \leq t$ ,  $1 \leq j \leq s+t$ ,  $j \neq i$ . It follows that if  $x \in N(u) - \{v\}$  and  $y \in N(v) - \{u\}$ , then  $d(x, y) = 3$ . Consequently, the edge  $uv$  is contained in no 4-cycle of  $EH(s, t)$ . Since in the Cartesian product of two nontrivial graphs every edge is contained in at least one 4-cycle, the result is proved.  $\square$

## 5 Concluding remarks

In this paper we have obtained several properties of exchanged hypercubes with respect to the distance function. Another closely related class of cubes studied in the literature is formed by dual-cubes  $D_n$ ,  $n \geq 1$ . These cubes were by now well studied, see for instance [1, 3, 21, 22]. Here we follow the notation used in [1]. As it turns out,  $D_n$  is

isomorphic to  $EH(n-1, n-1)$ . Hence all our results can be directly applied to the dual-cubes. For instance:

**Corollary 5.1** *If  $n \geq 1$ , then  $W(D_{n+1}) = (2n+3)2^{4n} - 2^{3n+1}$ .*

Dual-cubes are vertex-transitive graphs. For such graphs  $G$  the surface area is independent of a selected vertex  $u$ , hence the notation  $B_{G,u}(r)$  can be simplified to  $B_G(r)$ . Then we have:

**Corollary 5.2** *If  $n \geq 1$ , then  $B_{D_{n+1}}(r) = \binom{2n+1}{r-1} + \binom{n+1}{r} - \binom{n+1}{r-1}$ .*

**Proof.** Apply Theorem 3.4 together with the facts that  $D_{n+1}$  is isomorphic to  $EH(n, n)$  and that  $\binom{n}{r} - \binom{n}{r-2} = \binom{n+1}{r} - \binom{n+1}{r-1}$ .  $\square$

Very recently, dual-cubes  $D_n$  were generalized to *dual-cube-like networks*  $DC_n$  in [1].  $DC_n$  consists of  $2^n$  disjoint copies of  $Q_{n-1}$ . In addition, it contains a perfect matching, where the endpoints belong to different copies of  $Q_{n-1}$ , and between any two copies of  $Q_{n-1}$  there is at most one edge of the perfect matching. This generalization of dual-cubes to dual-cube-like networks can be naturally extended to generalize extended hypercubes to extended hypercube-like networks. We think it would be worth studying the extended hypercube-like networks.

## Acknowledgements

This work was supported in part by ARRS Slovenia under the grant P1-0297, the China-Slovenia bilateral grant BI-CN/11-13-001, and the national natural science foundation of China 11101378.

## References

- [1] A. Angjeli, E. Cheng, L. Lipták, Linearly many faults in dual-cube-like networks, *Theoret. Comput. Sci.* 472 (2013) 1–8.
- [2] H. Braun, F.C. Stephan, On optimizing diameter and average distance of directed interconnected networks, *IEEE Trans. Comput.* 42 (1993) 353–358.
- [3] J.-C. Chen, C.-H. Tsai, Conditional edge-fault-tolerant Hamiltonicity of dual-cubes, *Inform. Sci.* 181 (2011) 620–627.

- [4] Y.W. Chen, A comment on “The exchanged hypercube”, *IEEE Trans. Parallel Distrib. Syst.* 18 (2007) 576.
- [5] B. Cheng, J. Fan, X. Jia, J. Jia, Parallel construction of independent spanning trees and an application in diagnosis on Möbius cubes, *J. Supercomput.* (2013) DOI: 10.1007/s11227-013-0883-1.
- [6] E. Cheng, K. Qiu, Z. Shen, On the surface areas and average distances of meshes and tori, *Parallel Process. Lett.* 21 (2011) 61–75.
- [7] E. Cheng, K. Qiu, Z. Shen, On the surface area of the augmented cubes, *J. Supercomput.* 61 (2012) 856–868.
- [8] A.A. Dobrynin, R. Entringer, I. Gutman, Wiener index of trees: theory and applications, *Acta Appl. Math.* 66 (2001) 211–249.
- [9] A.A. Dobrynin, I. Gutman, S. Klavžar, P. Žigert, Wiener index of hexagonal systems, *Acta Appl. Math.* (2002) 247–294.
- [10] J. Fink, Perfect matchings extend to Hamilton cycles in hypercubes, *J. Combin. Theory Ser. B* 97 (2007) 1074–1076.
- [11] N. Graham, F. Harary, The number of perfect matchings in a hypercube, *Appl. Math. Lett.* 1 (1988) 45–48.
- [12] R. Hammack, W. Imrich, S. Klavžar, *Handbook of Product Graphs*, Second Edition, CRC Press, Boca Raton, FL, 2011.
- [13] A.M. Hinz, D. Parisse, The average eccentricity of Sierpiński graphs, *Graphs Combin.* 28 (2012) 671–686.
- [14] N. Imani, H. Sarbazi-Azad, S.G. Akl, Some topological properties of star graphs: The surface area and volume, *Discrete Math.* 309 (2009) 560–569.
- [15] W. Imrich, S. Klavžar, D.F. Rall, *Topics in Graph Theory: Graphs and Their Cartesian Products*, A K Peters, Wellesley, 2008.
- [16] J.-S. Kim, S.W. Kim, E. Cheng, L. Lipták, Topological properties of folded hyperstar networks, *J. Supercomput.* 59 (2012) 1336–1347.
- [17] S. Klavžar, Structure of Fibonacci cubes: a survey, *J. Comb. Optim.* 25 (2013) 505–522.
- [18] S. Klavžar, M.J. Nadjafi-Arani, Wiener index versus Szeged index in networks, *Discrete Appl. Math.* 161 (2013) 1150–1153.
- [19] C.-N. Kuo, H.-H. Chou, N.-W. Chang, S.-Y. Hsieh, Fault-tolerant path embedding in folded hypercubes with both node and edge faults, *Theoret. Comput. Sci.* 475 (2013) 82–91.

- [20] X.-J. Li, J.-M. Xu, Generalized measures of fault tolerance in exchanged hypercubes, *Inform. Process. Lett.* 113 (2013) 533–537.
- [21] Y. Li, S. Peng, Dual-cubes: a new interconnection network for high-performance computer clusters, in: *Proceedings of the 2000 International Computer Architecture*, 2000, pp. 51–57.
- [22] Y. Li, S. Peng, W. Chu, Efficient collective communications in dual-cube, *J. Supercomput.* 28 (2004) 71–90.
- [23] P.K.K. Loh, W.J. Hsu, Y. Pan, The exchanged hypercube, *IEEE Trans. Parallel Distrib. Syst.* 16 (2005) 866–874.
- [24] M. Ma, B. Liu, Cycles embedding in exchanged hypercubes, *Inform. Process. Lett.* 110 (2009) 71–76.
- [25] M. Ma, The connectivity of exchanged hypercubes, *Discrete Math. Algorithms Appl.* 2 (2010) 213–220.
- [26] M. Ma, L. Zhu, The super connectivity of exchanged hypercubes, *Inform. Process. Lett.* 111 (2011) 360–364.
- [27] I. Pesek, M. Rotovnik, D. Vukičević, J. Žerovnik, Wiener number of directed graphs and its relation to the oriented network design problem, *MATCH Commun. Math. Comput. Chem.* 64 (2010) 727–742.
- [28] S. Wang, G. Zhang, K. Feng, Fault tolerance in  $k$ -ary  $n$ -cube networks, *Theoret. Comput. Sci.* 460 (2012) 34–41.
- [29] X. Wang, J. Fan, X. Jia, S. Zhang, J. Yu, Embedding meshes into twisted-cubes, *Inform. Sci.* 181 (2011) 3085–3099.